

114

INTERNATIONAL
JOURNAL
OF BIOCLIMATOLOGY
AND BIOMETEOROLOGY

VOLUME 3
1959

U. of ILL. LIBRARY
AUG 10 1971
CHICAGO CIRCLE



International Journal of Bioclimatology and Biometeorology

Journal of the
„International Society of
Bioclimatology and Biometeorology“

**VOLUME III
(1959)**

Reprinted with the permission of the International Society of
Biometeorology

JOHNSON REPRINT CORPORATION
111 Fifth Avenue, New York, N.Y. 10003

JOHNSON REPRINT COMPANY LIMITED
Berkeley Square House, London, W.1

International Journal
of
Bioclimatology and Biometeorology

Journal of the
International Society of
Bioclimatology and Biometeorology

VOLUME III
(1969)

First reprinting, 1965, Johnson Reprint Corporation

Printed in the United States of America

EDITORIAL BOARD

ASSOCIATE EDITORS

Phytological Bioclimatology

Dr. H. Boyko (Israel)
Prof. L. Emberger (France)
Prof. Dr. A. Mäde (D.D.R.)
Prof. A. de Philippis (Italy)
Prof. R. B. Platt (U.S.A.)

Zoological Bioclimatology

Prof. J. Benoit (France)
Dr. J. L. Cloudsley-Thompson
(Gr. Britain)
Prof. K. Göszwald (Germany)
Dr. W. G. Wellington (Canada)

Human Bioclimatology

Prof. W. Amelung (Germany)
Prof. K. J. Buettner (U.S.A.)
Prof. M. Fontaine (France)
Dr. E. M. Glaser (Gr. Britain)

MANAGING EDITOR

Dr. S. W. Tromp (Netherlands)

GENERAL INFORMATION

The Journal is published by the "International Society of Bioclimatology and Biometeorology" and is printed in the Netherlands in off-set type.

PURPOSE OF THE JOURNAL

The "International Society of Bioclimatology and Biometeorology" covers in its journal the following subjects:

1. Summaries of completed bioclimatological studies which can be published in full by the authors in any other scientific journal;
2. Short reports on research in progress in order to stimulate team work between research workers in different parts of the world;
3. Critical reviews of special fields of bioclimatology;
4. A complete survey of world literature. Titles, names of authors and their addresses (if possible) are given; where possible, abstracts of important articles are also provided;
5. Summaries of Symposia or Congresses of National and International Organizations dealing with bioclimatological subjects;
6. Information on dates and programmes of Symposia and Congresses related to bioclimatological subjects;
7. Reports from the chairmen of the special scientific committees of the Society:
 - a. Committee for the study of bioclimatological aspects of ALLERGIC DISEASES: secretary: Dr. D. Ordman (S. Africa). Two sub-committees:
 - (1) Study of the indirect effect of meteorological factors on Allergic Diseases through pollen, spores and other allergens (Organized by Dr. D. Ordman, S. Africa);
 - (2) Study of the possible direct effect of meteorological factors on Hay fever, Asthma and other Allergic Diseases (Organized by Dr. S.W. Tromp, Netherlands).
 - b. Committee for ECOLOGICAL CLIMATOGRAPHY: Chairman Dr. H. Boyko (Israel). This committee is concerned with the definition of climate on the basis of the plant and animal associations of a region.
 - c. Committee for INSTRUMENTATION: Chairman Dr. J.F. Griffiths (Great Britain). This committee compiles all available technical and economic data on instruments used in bioclimatological research.
 - d. Committee for the study of possible biological effects of various types of IONISATION OF THE AIR: Chairman Dr. J.H. Kornbluh (USA); secretary Mr. John C. Beckett (U.S.A.).
 - e. Committee of NAUTICAL BIOCLIMATOLOGY: Chairman Prof. F. Molino (Italy); secretary Prof. V. Parodi (Italy). Two sub-committees:
 - (1) Study of the physiological and pathological phenomena observed by ship surgeons (both naval and mercantile marine) in man and animals in various climates at sea.
 - (2) Study of the influence of climate at sea on the living cargo in ships, so called CARGO BIOCLIMATOLOGY.
 - f. Committee for the study of CHEMICAL TESTS, used in bioclimatological research in general and cosmic bioclimatology in particular: Chairman Prof. G. Piccardi (Italy); secretary Mrs. Eng. C. Capel-Boute (Belgium).
 - g. Committee for the study of TROPICAL BIOCLIMATOLOGY: Chairman Dr. W.S.S. Ladell (Great Britain); secretary Mr. J.P. Nicolas (France).
 - h. Committee on SOLAR RADIATION, in relation to biometeorology and bioclimatology: Chairman Prof. N. Robinson (Israel); secretary Miss Dr. Inge Dirmhirn (Austria)
8. Requests from both members and non-members for scientific information, about studies abroad, etc.;
9. Advertisements of private scientific firms of good standing.

Points 1 - 4 are dealt with in parts I - VI of each volume. Points 5 - 9 are dealt with in part VII.

Each year a new volume will be published, each volume being divided into seven parts and each part containing several sections. Depending on the material received and the funds of the I.S.B.B.

a number of sheets referring to various sections or parts of a volume will be sent at intervals to members. On top of each article the part and section number will be clearly indicated.

Each member will receive at the beginning of the year the cover of the volume for that year and the various coloured sheets separating the different parts.

The volume will be prepared in loose-leaved form, in order to facilitate the insertion of the sheets received in the corresponding parts or sections of the volume. At the end of the year, the member will be able either to keep the volume as one unit or alternatively the various parts can be separated and bound separately.

For this purpose a special printed white sheet, with the name of the Journal is supplied to members.

SUBSCRIPTIONS

Members of the Society receive the journal against payment of their yearly membership fee.

Non-members can obtain the journal against payment of 30 Dutch guilders per year, postage included.

EDITORIAL CORRESPONDENCE

Manuscripts submitted for consideration and all correspondence relating to editorial matters should be addressed to "the Executive Editor, International Journal of Bioclimatology and Biometeorology, Hofbrouckerlaan 54, Oegstgeest (Leiden), Holland".

MANUSCRIPTS

For preparation of manuscripts see "Instructions to Contributors" attached.

The Editors reserve the right to refuse any manuscript submitted, whether on invitation or otherwise, and to make suggestions and modifications before publication.

Articles accepted by the Editors remain the property of the International Society of Bioclimatology and Biometeorology, but may be reprinted in other scientific journals with the consent of the Editors of the Journal.

ADVERTISING

Companies in good standing may advertise in the Journal.

All correspondence should be addressed to the Executive Editor of the Journal, Hofbrouckerlaan 54, Oegstgeest (Leiden), Holland.

CONTENTS :

Each volume is divided into seven parts each of which is again subdivided into a number of sections as indicated below. Reports appertaining to one or more of these parts or sections will be sent to members at intervals according to the amount of data received by the Editor.

Part I: GENERAL BIOCLIMATOLOGY (yellow cover)

- Sect. A : History and biography
- Sect. B : Bioclimatological teaching
- Sect. C : Instrumentation
 - 1. General
 - 2. Phytological bioclimatology
 - 3. Zoological bioclimatology
 - 4. Human bioclimatology
- Sect. D : Aerosols and chemical aspects of bioclimatology
- Sect. E : Statistical methods in bioclimatology
- Sect. F : Miscellaneous data (classification of climates, air masses, human typology, etc., as far as they affect bioclimatological observations)
- Sect. G : World literature

Part II: PHYTOLOGICAL BIOCLIMATOLOGY (green cover)

- Sect. A : General phytological bioclimatology
- Sect. B : Agricultural bioclimatology
 - 1. General agricultural bioclimatology
 - 2. Agricultural phenology
- Sect. C : Forest bioclimatology
- Sect. D : Physiological phyto-bioclimatology
- Sect. E : Pathological phyto-bioclimatology
- Sect. F : World literature

Part III: ZOOLOGICAL BIOCLIMATOLOGY (red cover)

- Sect. A : General zoological bioclimatology
- Sect. B : Entomological bioclimatology
- Sect. C : Veterinary bioclimatology
- Sect. D : World literature

Part IV: HUMAN BIOCLIMATOLOGY (blue cover)

- Sect. A : Physiological bioclimatology
 - 1. General physiological bioclimatology
 - 2. Geographical bioclimatology
 - 3. Ethnological bioclimatology
 - 4. Acclimatisation bioclimatology
- Sect. B : Social bioclimatology
 - 1. General social bioclimatology
 - 2. Psychological bioclimatology (including aestheto-bioclimatology)
 - 3. Archeological bioclimatology
- Sect. C : Pathological bioclimatology
 - 1. General pathological bioclimatology
 - 2. Meteorological pathology
 - 3. Climatological pathology
 - 4. Air pollution pathology
 - a. Pollution with organic particles (pollen, fungi, etc.)
 - b. Pollution with inorganic particles (dust, etc.)
 - c. Chemical pollution
 - 5. Geographical climatopathology
 - 6. Climatotherapy
 - a. General climatotherapy and therapeutic climates
 - b. Thalassotherapy (climatological-)
 - c. Heliotherapy
 - d. Thermotherapy
 - e. Aerosol (and ionisation) therapy
 - f. Socio-climatotherapy
 - g. Climatic health resorts
 - h. Other therapeutic methods

- Sect. D : Urban bioclimatology
 - 1. General urban bioclimatology
 - 2. Architectural bioclimatology
 - 3. Sanatorium bioclimatology

Sect. E : World literature

Part V : COSMIC BIOCLIMATOLOGY (grey cover)

- Sect. A : General cosmic bioclimatology
- Sect. B : Special cosmic bioclimatology
- Sect. C : World literature

Part VI : PALEO-BIOCLIMATOLOGY (brown cover)

- Sect. A : General paleo-bioclimatology
- Sect. B : World literature

Part VII : MISCELLANEOUS BIOCLIMATOLOGICAL DATA (White cover)

Sect. A : Scientific committees of the Society

1. Reports

- a. Allergic diseases
- b. Ecological climatology
- c. Instrumentation
- d. Ionisation of the air
- e. Nautical bioclimatology (General-, cargo bioclimatology)
- f. Chemical tests
- g. Tropical bioclimatology
- h. Solar Radiation

2. Literature

- Sect. B : Symposia or congresses of national bioclimatological societies
(dates, programmes, summaries of important lectures, decisions, etc.)
- Sect. C : Bioclimatological stations and institutions
- Sect. D : Requests from members
- Sect. E : Requests from non-members
- Sect. F : Book reviews
- Sect. G : International Organizations (WMO, FAO, WHO, etc.)
- Sect. H : Advertisements

DEFINITIONS:

As not all members may be acquainted with the various bioclimatological divisions outside their own speciality, the following list of definitions has been prepared:

BIOClimATOLOGY and BIOMETEOROLOGY (in general): comprises the study of the direct and indirect interrelations between the geo-physical and geo-chemical environment and living organisms, plants, animals and man. The term "environment" is broadly conceived and includes micro-, macro- and cosmic environments and the diverse physical and chemical factors which comprise these environments.

Investigations in these disciplines are conducted in nature or in the laboratory under as rigidly controlled conditions as possible to describe measurable and reproducible physical, chemical and biological factors which show a sufficiently high statistical correlation with measurable physiological and pathological processes to suggest a valid cause and effect relationship between organism and environment.

AEROSOL BIOCLIMATOLOGY: Study of the biological effects of aerosols (i.e. gaseous, liquid or solid aggregates floating in the air, with diameters of $1/1000$ to 10 micron and consisting of hundreds to millions of molecules, often with either positive or negative electrical charges).

PHYTOLOGICAL BIOCLIMATOLOGY: Study of the influence of climate, weather and cosmic factors on plants.

ZOOLOGICAL BIOCLIMATOLOGY: Study of the influence of climate, weather and cosmic factors on animals.

ENTOMOLOGICAL BIOCLIMATOLOGY: Study of the influence of climate, weather and cosmic factors on insects and other terrestrial Arthropoda.

VETERINARY BIOCLIMATOLOGY: Study of the influence of climate, weather and cosmic factors on domestic and farm animals and birds and on animal products such as eggs, wool, milk, etc.

HUMAN BIOCLIMATOLOGY: Study of the influence of climate, weather and cosmic factors on man.

GENERAL PHYSIOLOGICAL BIOCLIMATOLOGY: Study of the influence of specific single or groups of meteorological components, of different climates (mountain-, marine-, forest climate, etc.) and of their seasonal variations on the various physiological processes of normal, healthy man.

GEOGRAPHICAL BIOCLIMATOLOGY: Study of the influence of geographical differences on the physiological processes of normal, healthy man.

ETHNOLOGICAL BIOCLIMATOLOGY: Study of the influence of climate and weather on race and body structure of man.

ACCLIMATISATION BIOCLIMATOLOGY: Study of the adaptation of the human body to extreme climatological conditions.

SOCIAL BIOCLIMATOLOGY: Study of the influence of climate and weather on the social habits of man.

PSYCHOLOGICAL BIOCLIMATOLOGY: Study of the influence of climate and weather on the mental processes of man.

AESTHETIC BIOCLIMATOLOGY: Study of the influence of climate and weather on the aesthetic expression of man.

ARCHEOLOGICAL BIOCLIMATOLOGY: Study of the influence of climate and weather on the origin, distribution and disappearance of past civilizations.

PATHOLOGICAL BIOCLIMATOLOGY: Study of the influence of climate and weather on the various physiological and pathological phenomena associated with the diseases of man.

METEOROLOGICAL PATHOLOGY: Study of the influence of specific single meteorological components (temperature, humidity, etc.) or groups of components on the origin and frequency of diseases and on the physiological phenomena of the diseases of man.

CLIMATOLOGICAL PATHOLOGY: Study of the influence of different climates (marine-, forest-, mountain-, etc.) and their seasonal variations on the origin and frequency of diseases and on the physiological phenomena of the diseases of man.

AIR POLLUTION PATHOLOGY: Study of the pathological influences of air pollution (either organic or inorganic particles or chemical substances) on man.

GEOGRAPHICAL CLIMATOPATHOLOGY: Study of the geographical distribution of diseases as a result of geographical differences in climate and in single or in groups of meteorological factors.

URBAN BIOCLIMATOLOGY: Study of the micro climates in houses and cities, their influence on the health of man, and of the methods of eliminating unfavourable influences and of increasing favourable biological effects in certain types of architectural construction and town planning.

SANATORIUM BIOCLIMATOLOGY: Study of the best location and construction methods of sanatoria from the point of view of climate and weather.

CLIMATOTHERAPY: Study of the therapeutic influence of certain climates and meteorological conditions on the diseases of man.

THALASSOTHERAPY (CLIMATOLOGICAL-): Study of the therapeutic influence of marine climates on man.

HELIOOTHERAPY: Study of the therapeutic influence of solar radiation on man.

THERMOTHERAPY: Study of the therapeutic influence of various forms of heat on man.

AEROSOTHERAPY: Study of the therapeutic influence of certain aerosols on man.

SOCIO-CLIMATOTHERAPY: Study of construction methods of schools, of location and construction of holiday camps for children and of other social aspects of life as a function of climate and weather and of the methods for improvement of the favourable climatological effects.

COSMIC BIOCLIMATOLOGY: Study of the biological effects of cosmic factors.

PALEO-BIOCLIMATOLOGY: Study of the influence of the climates of the past on the development and geographical distribution of animals and plants on earth.

ECOLOGICAL CLIMATOGRAPHY: Study of the definition of climate on the basis of plant and animal associations.

NAUTICAL BIOCLIMATOLOGY: Study of the physiological and pathological phenomena observed by ship surgeons (both naval and mercantile marine), in man and animals in various climates at sea.

CARGO BIOCLIMATOLOGY: Study of the influence of climate at sea on the living cargo in ships (plants, fruits and animals).

VII

INSTRUCTIONS TO CONTRIBUTORS

In the interests of uniformity, contributors are requested to adhere to the following instructions:

1. CORRESPONDENCE AND MANUSCRIPTS:

All correspondence should be addressed to Dr. S.W. Tromp, Hofbrouckerlaan 54, Oegstgeest (Leiden), Netherlands.

2. PAPERS SUBMITTED TO THE JOURNAL:

- (a) Papers should be written in English, French or German. Manuscripts should be typed, on one side of the paper only, double-spaced, quarto-size, leaving top and left hand margins at least 2.5 cm. (one inch) wide. No underlining should be used in the text for words, titles etc., but CAPITAL LETTERS, or CAPITAL LETTERS WITH DOUBLE SPACING between the letters should be employed. Manuscripts should be packed flat.
- (b) Dates should be in the form "5 February 1957" not "February 5th 1957".
- (c) All Latin specific names should be followed by the authority when the name is first mentioned in the text or tables.
- (d) Accurate details of experimental controls should be given in all cases.
- (e) Each article should begin with a short abstract of not more than 50 words, immediately after the title and author's name, preferably in 3 languages (English, French and German). It would be appreciated if each article could be concluded with a short summary containing the main conclusions.
- (f) Four copies of the complete text of each manuscript should be submitted.
- (g) Graphs and other illustrative material which can be reproduced in off-set print without additional expenses (i.e. graphs and diagrams drawn in jet-black Indian ink on strong white or transparent paper) are permitted without charge. However, additional material of this kind which requires special cliché work, e.g. photographs (see point 6), may be allowed at cost, at the discretion of the editors.
- (h) Proof-reading is done by the Editors.

3. REFERENCES

Bibliographic references should be listed in alphabetic order at the end of the paper and continuous numbering should be used. After the title of the Journal quoted, the following particulars should be supplied: the number of the vol., part (if known), first and last page and finally, the year of publication.

For example: 1. BROWN, A.B. : Insects and Weather - Proc. R. Ent. Soc. London, 54, 7, 3-10, 1954.

2. LEE, D.H.K. : "Climate and Economic development in the Tropics" - Harper & Brothers Publ. Co, New York, 182 p., 1957.

3. SMITH, P.O. : Influence of Aerosols on health - Int. J. Biocl. Biomet., 1, IV, Sect. C 6e, 20 - 25, 1958.

References should be referred to in text as (1), (2) etc.

Titles of Journals should be abbreviated, as much as possible, according to the World List of Scientific Periodicals, except that the English usage of capitals for adjectives should be followed in the titles of British and American periodicals.

4. TABLES:

Tables should be set out on separate sheets and numbered in Arabic numerals. They should be supplied with headings and kept as simple as possible. They should be referred to in the text as "Table 2", etc.

5. GRAPHS AND DIAGRAMS:

These should be drawn in jet-black Indian ink on strong white or transparent paper. They should be numbered in succession in Arabic numerals and should be referred to in the text as "Fig. 3" etc.

Graphs and diagrams should be large enough to permit reduction to a size of 10 x 10 cm. (4

VIII

x 4 inches) without affecting legibility.

In exceptional cases graphs or diagrams not complying to these rules can be redrawn for the author in the office of the Executive Editor at cost price.

6. PHOTOGRAPHS:

Photographs can be easily reproduced in off-set print but should be unmounted, glossy prints showing good detail and contrast, permitting reduction to a size 10 x 10 cm (4 x 4 inches) without affecting legibility. They should be numbered separately from the tables, graphs and diagrams and should be referred to in the text as "Photo 4" etc.

7. REPRINTS:

50 copies of articles accepted by the Editors will be supplied gratis. Further copies may be obtained at cost price:

If ordered before printing, the prices are as follows:

100 reprints: 3.00 guilders/page, each additional 50 copies at 1.15 guilder/page.

For example 200 reprints of a 5 page article will cost $5 \times 5.30 = 26.50$ guilders + postage.

If ordered after printing is completed the price will be:

100 reprints: 4.30 guilders/page, each additional 50 copies 1.15 guilder/page.

Thus 200 reprints of a 5 page article will cost $5 \times 6.60 = 33$ guilders + postage.

C L E A R I N G H O U S E

1. Requests for reprints of articles mentioned in the bibliographic summaries should first be addressed directly to the authors. If no reprints are available the International Journal of Bioclimatology and Biometeorology can prepare photocopies at cost price (abt. 40 Dutch cents = 11 dollarcents per photocopy of $4\frac{1}{4} \times 5\frac{3}{4}$ inches; or 60 Dutch cents = 16 dollarcents per photocopy of $5\frac{3}{4} \times 8\frac{1}{2}$ inches; or microfilms per frame of 2 pages = 8 dollarcents. These prices may change during 1958 and do not include postage). The forms enclosed, duly signed, should be sent to the Executive Editor of the Journal, Hofbrouckerlaan 54, Oegstgeest (Leiden), Holland.
2. Requests from members concerning literature relating to research projects should be sent to the Executive Editor of the Journal who will try to compile all available references concerning the subject and will forward these to the person requesting the information. If considerable administrative work is involved in compiling the data requested by a member a special service charge may have to be paid by the requesting member. Requests from non-members are only handled against payment of a service charge. The Editorial Staff may deviate from this rule in special circumstances.

R E Q U E S T F O R P H O T O C O P I E S

INTERNATIONAL SOCIETY OF BIOCLIMATOLOGY AND BIOMETEOROLOGY

The undersigned : NAME (printed) :

ADDRESS (printed) :

would like to receive photo copy(ies) of the following article:

Author :

Title :

Journal :
(only complete references can be dealt with)

On receipt of this bill he agrees to pay immediately the amount required for reproducing the article, either to the account of the Int. Soc. of Bioclimatology and Biometeorology at the Twentsche Bank, Breestraat 81, Leiden, Holland, or to B.W. Blijdenstein & Co., (Bank), 54 - 56 Threadneedle Street, London E.C. 2, Great Britain. On the payment order the number of the bill should be mentioned. The Society will mail the copy (ies) immediately upon receipt of payment.

Date:

Signature:

INTERNATIONAL JOURNAL OF BIOCLIMATOLOGY
AND BIOMETEOROLOGY

INDEX TO VOLUME III, 1959 *

AUTHOR INDEX

- BORTELS, H. : Das Problem meteorobiologischer Beziehungen im Lichte mikrobiologischer Forschung - I.J.B.B., III, I F, 1959.
- BOYKO, H., BOYKO, E. : Seawater irrigation a new line of research on a bioclimatic plant-soil complex - I.J.B.B., III, II B 1, 1959.
- BOURKE, P.M.A. : Meteorology and the timing of fungicide applications against potato blight - I.J.B.B., III, II E, 1959.
- BUETTNER, K.J.K. : Program of symposium on bioclimatology during the 176th National Meeting of the American Meteorological Society held jointly with the section of Meteorology, American Geophysical Union at Washington, D.C., U.S.A. on 5 May 1959, I.J.B.B. III, VII B, 1959.
- CLOUDSLEY-THOMPSON, J.L. : Microclimate, diurnal rhythms and the conquest of the land by arthropods - I.J.B.B., III, III B, 1959.
- CLOUDSLEY-THOMPSON, J.L. : The effects of pollution on living material - I.J.B.B., III, VII B, 1959.
- ERBAN, L. : A study of biochemical and haematological changes under the application of ionized air - I.J.B.B., III, IV C 6 e, 1959.
- ITO, S., TSUJIOKA, K., SAITO, H. : Blood clotting time under metal cover (biological P-test) - I.J.B.B., III, V A, 1959.
- KREČMER, I. : Program of the Bioclimatological Conference at Liblice, (Czechoslovakia) (3-5 November 1958) - I.J.B.B., III, VII B, 1959.
- LIMAN, ST. : Die Bioklimatologie des Menschen in Polen - I.J.B.B., III, VII C, 1959.
- MADE, A. : Ueber den Stand der agrarmeteorologischen Forschung in der Deutschen Demokratischen Republik - I.J.B.B., III, VII G, 1959.
- NOFFSINGER, T.L., ANDREWS, F.N. : Correlation of body temperature and of respiration rate of swine with environmental temperature - I.J.B.B., III, III C, 1959.
- OOSTERLEE, C.C. : The influence of climatic factors on the heat balance and milk yield of the cow in relation to the design of farm buildings in the Netherlands - I.J.B.B., III, III C, 1959.
- OPDYKE, N.D. : The impact of paleomagnetism on paleoclimatic studies - I.J.B.B., III, VI A, 1959.
- ORTMEYER, : Program Bioclimatological Conference on Climatological and Balneological Health Resorts at E. Berlin, D.D.R. (3-5 March 1959) - I.J.B.B., III, VII B, 1959.
- PASZYNSKI, J. : Preliminary research on the transparency of the atmosphere at Bydgoszcz - I.J.B.B., III, I F, 1959.
- PICCARDI, G. : Symposium International sur les Relations entre Phénomènes Solaires et Terrestres en Chimie-physique et en Biologie - I.J.B.B., III, VII B, 1959.
- PIEŚLAK, Z. : Report of the Second Session of the Commission for Agricultural Meteorology of the World Meteorological Organization at Warsaw (Poland) (29 Sept.- 17 Oct. 1958) - I.J.B.B., III, VII G, 1959.

* "International Journal of Bioclimatology and Biometeorology" in references briefly indicated as I.J.B.B.

- RENBURN, E.T. : A contribution to the physiology of damp cold an aspect of temperate climate, Part I. The chills of a damp cold climate an historical survey - I.J.B.B., III, IV A 4, 1959.
- RENBURN, E.T., REID, A.M., GILLING, D.R.: A contribution to the physiology of damp cold an aspect of temperate climate, Part II. An investigation on clothed men under indoor conditions - I.J.B.B., III, IV A 4, 1959.
- SARGENT, F. : World Meteorological Organization and International Society of Bioclimatology and Biometeorology to collaborate - I.J.B.B., III, VII G, 1959.
- SCHOLZ, : Probleme der Bioklimatologie und des Stallklimas bei der Rinder- und Schweinehaltung, Program of Symposium Pörsneck, Thüringen, D.D.R. (28-29 May 1959) - I.J.B.B., III, VII B, 1959.
- SVORAD, D., WELLNEROVA, J. : Temperature and air humidity as factors influencing sleep and wakefulness - I.J.B.B., III, IV B 2, 1959.
- TYCZKA, S. : World literature - human Bioclimatology (Polish references) - I.J.B.B., III, IV E, 1959.
- WELLINGTON, W.G. : World literature - Entomological Bioclimatology - I.J.B.B., III, III D, 1959.
- WELLNEROVÁ, J., SVORAD, D.: Participation of some meteorological factors on the determination of emotional reactivity - I.J.B.B., III, IV B 2, 1959.
- WÓJTOWICZ, W. : Report on the Bioclimatological Conference in Ciechocinek (Poland) (30-31 October 1958) - I.J.B.B., III, VII B, 1959.
- WORLD LITERATURE : General phytological bioclimatology; general agricultural bioclimatology; agricultural phenology; forest bioclimatology; physiological bioclimatology - I.J.B.B., III, II F, 1959.
- WORLD LITERATURE : General zoological bioclimatology; physiological zoo-bioclimatology; entomological bioclimatology; veterinary bioclimatology - I.J.B.B., III, III D, 1959.
- WORLD LITERATURE : General human bioclimatology; gen. physiological bioclimatology; acclimatization bioclimatology - I.J.B.B., III, IV E, 1959.

INTERNATIONAL JOURNAL OF BIOCLIMATOLOGY
AND BIOMETEOROLOGY

INDEX TO VOLUME III, 1959

SUBJECT INDEX *

PART I : GENERAL BIOCLIMATOLOGY

SECT. A : HISTORY AND BIOGRAPHY : ---

SECT. B : BIOCLIMATOLOGICAL TEACHING : ---

SECT. C : INSTRUMENTATION :

1. General : ---

2. Phytological bioclimatology : ---

3. Zoological bioclimatology : ---

4. Human bioclimatology : ---

SECT. D : AEROSOLS AND CHEMICAL ASPECTS OF BIOCLIMATOLOGY : ---

SECT. E : STATISTICAL METHODS IN BIOCLIMATOLOGY : ---

SECT. F : MISCELLANEOUS DATA (CLASSIFICATION OF CLIMATES, AIR MASSES, HUMAN TYPOLOGY, ETC., AS FAR AS THEY AFFECT BIOCLIMATOLOGICAL OBSERVATIONS): Bortels (H.), Paszyński (J.).

SECT. G : WORLD LITERATURE : ---

PART II : PHYTOLOGICAL BIOCLIMATOLOGY

SECT. A : GENERAL PHYTOLOGICAL BIOCLIMATOLOGY : ---

SECT. B : AGRICULTURAL BIOCLIMATOLOGY :

1. General agricultural bioclimatology: Boyko (H.) and Boyko (E.).

2. Agricultural phenology : ---

SECT. C : FOREST BIOCLIMATOLOGY : ---

SECT. D : PHYSIOLOGICAL PHYTO-BIOCLIMATOLOGY : ---

SECT. E : PATHOLOGICAL PHYTO-BIOCLIMATOLOGY : Bourke (P.M.A.)

SECT. F : WORLD LITERATURE : General phytological bioclimatology (compiled by secretariat), general agricultural bioclimatology (comp.secr.), agricultural phenology (comp. secr.), forest bioclimatology (comp. secr.), physiological phyto-bioclimatology (comp. secr.).

PART III : ZOOLOGICAL BIOCLIMATOLOGY

SECT. A : GENERAL ZOOLOGICAL BIOCLIMATOLOGY : ---

SECT. B : ENTOMOLOGICAL BIOCLIMATOLOGY : Cloudsley-Thompson (J.L.)

SECT. C : VETERINARY BIOCLIMATOLOGY : Noffsinger (T.L.) and Andrews (F.N.), Oosterlee (C.C.).

SECT. D : WORLD LITERATURE : Entomological bioclimatology (Wellington. W.G.), general zoological bioclimatology (comp. secr.), physiological zoo-bioclimatology (comp. secr.), entomological bioclimatology (comp. secr.), veterinary bioclimatology (comp. secr.).

* --- : No publications received.

PART IV : HUMAN BIOCLIMATOLOGY

SECT. A : PHYSIOLOGICAL BIOCLIMATOLOGY :

1. General physiological bioclimatology : ---
2. Geographical bioclimatology : ---
3. Ethnological bioclimatology : ---
4. Acclimatisation bioclimatology : Renbourn (E.T.).

SECT. B : SOCIAL BIOCLIMATOLOGY :

1. Social bioclimatology (general) : ---
2. Psychological bioclimatology (including aestheto-bioclimatology):
Svorad (D.) and Wellnerová (J.), Wellnerová (J.) and Svorad (D.).
3. Archeological bioclimatology: ---

SECT. C : PATHOLOGICAL BIOCLIMATOLOGY :

1. General pathological bioclimatology : ---
2. Meteorological pathology : ---
3. Climatological pathology : ---
4. Air pollution pathology :
 - a. Pollution with organic particles (pollen, fungi, etc.): ---
 - b. Pollution with inorganic particles (dust, etc.) : ---
 - c. Chemical pollution : ---
5. Geographical climatopathology : ---
6. Climatotherapy :
 - a. General Climatotherapy and Therapeutic climates: ---
 - b. Thalassotherapy (climatological -) : ---
 - c. Heliotherapy : ---
 - d. Thermotherapy : ---
 - e. Aerosol therapy (and Iono therapy) : Erban (L.).
 - f. Socio-climatotherapy : ---
 - g. Climatic health resorts : ---
 - h. Other therapeutic methods : ---

SECT. D : URBAN BIOCLIMATOLOGY :

1. General urban bioclimatology : ---
2. Architectural bioclimatology : ---
3. Sanatorium bioclimatology : ---

SECT. E : WORLD LITERATURE: Human bioclimatology (Tyczka, S.), general phys. bioclimatology (comp. secr.), acclimatisation bioclimatology (comp. secr.).

PART V : COSMIC BIOCLIMATOLOGY

SECT. A : GENERAL COSMIC BIOCLIMATOLOGY : Itoh (S.) and Tsujioke (K.) and Saito (H.)

SECT. B : SPECIAL COSMIC BIOCLIMATOLOGY : ---

SECT. C : WORLD LITERATURE : ---

PART VI : PALEO - BIOCLIMATOLOGY

SECT. A : GENERAL PALEO-BIOCLIMATOLOGY : Opdyke (N.D.)

SECT. B : WORLD LITERATURE : ---

PART VII : MISCELLANEOUS BIOCLIMATOLOGICAL DATA

SECT. A : SCIENTIFIC COMMITTEES OF THE SOCIETY :

1. Reports :
 - a. Allergic diseases : ---
 - b. Ecological climatology : ---
 - c. Instrumentation : ---
 - d. Ionisation of the air : ---
 - e. Nautical bioclimatology (general-, cargo bioclimatology) : ---
 - f. Chemical tests : ---
 - g. Tropical bioclimatology : ---
 - h. Solar radiation : ---
2. Literature : ---

- SECT. B : SYMPOSIA OR CONGRESSES OF INTERNATIONAL ORGANIZATIONS AND OF NATIONAL BIOCLIMATOLOGICAL SOCIETIES (DATES, PROGRAMMES, SUMMARIES OF IMPORTANT LECTURES, DECISIONS, ETC.) : Preliminary Programmes 2nd International Bioclimatological Congress; Symposium International sur les relations entre phénomènes solaires et terrestres en chimie-physique et en biologie (Piccardi, G.); Symposium on Bioclimatology (Buettner, K.J.K.); Symposium on the effects of pollution on living material (Cloudsley-Thompson, J.L.); Report on the Bioclimatological Conference in Ciechocinek, Poland (Wojtowicz, W.); Program of the Bioclimatological Conference at Liblice, Czechoslovakia (Krečmer, V.); Program bioclimatological Conference on Climatological and Balneological Health Resorts at E. Berlin (Ortmeyer); Probleme der Bioklimatologie und des Stallklimas bei der Rinder- und Schweinehaltung at Pössneck, D.D.R. (Scholz).
- SECT. C : BIOCLIMATOLOGICAL STATIONS AND INSTITUTIONS : Liman (St.), Mäde (A.).
- SECT. D : REQUESTS FROM MEMBERS AND GENERAL INFORMATION : List of regular Members and List of members according to 44 countries on January 1st, 1960 (for members only).
- SECT. E : REQUEST FROM NON-MEMBERS : ---
- SECT. F : BOOK REVIEWS : Recent Bioclimatological publications
- SECT. G : INTERNATIONAL ORGANIZATIONS (WMO, FAO, WHO, etc.) : Third Session of the World Meteorological Congress at Geneva, Switzerland; Reports of the Second Session of the Commission for Agricultural Meteorology of the World Meteorological Organization at Warsaw, Poland (Piéslak, Z.); Recent Arid Zone Projects (UNESCO); World Meteorological Organization and International Society of Bioclimatology and Biometeorology to collaborate (Sargent, F.)
- SECT. H : ADVERTISEMENTS : ---

PART I

GENERAL BIOCLIMATOLOGY

(1959)

Section A : History and biography

Section B : Bioclimatological teaching

Section C : Instrumentation

1. General
2. Phytological bioclimatology
3. Zoological bioclimatology
4. Human bioclimatology

**Section D : Aerosols and chemical aspects of
bioclimatology**

Section E : Statistical methods in bioclimatology

**Section F : Miscellaneous data (classification of climates,
air masses, human typology, etc., as far as they
affect bioclimatological observations)**

GENERAL BIOCLIMATOLOGY
Section F: Miscellaneous data
(Possible causes of Bioclimatological effects)

DAS PROBLEM METEORO-BIOLOGISCHER BEZIEHUNGEN
IM LICHT E MIKROBIOLOGISCHER FORSCHUNG

von

Dr.H. Bortels (Germany) *

Seit einigen Jahrzehnten finden die schon im Altertum bekannten, heute als Wetterföhligkeit und Wetterempfindlichkeit oft erwöhlten Beziehungen zwischen physiologischen und psychologischen Reaktionen des Menschen einerseits und meteorologischen und solaren Prozessen andererseits wieder allgemeine Beachtung und intensive wissenschaftliche Bearbeitung. Ausführliche Berichte wurden u.a. von PETERSEN (43), DE RUDDER (53), DÜLL (26), BERG (4), CURRY (25) und ASSMANN (2) veröffentlicht. Der Mensch steht im Mittelpunkt dieser Forschung. Deshalb wird in fast allen zu diesem Problem aufgestellten und grösstenteils wieder verworfenen Theorien über die Kausalkette das Nervensystem als Rezeptor eines mit Wetteränderungen verknöhlften, nicht näher bekannten und verschieden gedeuteten Reizes genannt. Hierzu und zu anderen Teilen des Gesamtproblems soll im folgenden aus mikrobiologischer Sicht Stellung genommen werden. Die Mikrobiologie genieusst dabei den Vorzug, dass ihre verhältnismässig einfach gebauten Objekte im Experiment leichter und auch eindeutiger befragt werden können als höher organisierte Lebewesen wie der Mensch.

Vor etwa 30 Jahren wurde von BORTELS (7) festgestellt, dass Einzeller auf bevorstehende oder gleichzeitig ablaufende Wetteränderungen reagieren. Die ersten Versuchsreihen über mehrere Jahre ergaben auffallende, aber in ihrer Signifikanz damals nicht mit statistischen Methoden geprüfte Korrelationen zwischen verschiedenen Lebensäusserungen einiger Bakterien und Pilze und meteorologischen Ereignissen wie Änderungen der Bewölkung und vor allem des Luftdrucks (8). Bei abnehmender Bewölkungsdichte nahm die Beweglichkeit mehrerer verschiedener Bakterienarten gleichzeitig zu und bei zunehmender Bewölkungsdichte und Niederschlägen nahm sie bis zur völligen Ruhe ab. Bei antizyklonaler Wetterentwicklung vermehrten sich z.B. aerobe Bakterien schneller, leuchteten Kulturen eines Leuchtbakteriums stärker, kopulierten verschiedene Mikroorganismen zahlreicher und liessen Sporangien von Phytophthora infestans weit mehr Zoosporen schlöpfen als bei zyklonaler Wetterentwicklung. Anaerobe Mikroorganismen reagierten dagegen umgekehrt, indem die von ihnen ausgelösten Reduktionen und Gärungen, z.B. die CO₂-Bildung durch Hefe, bei zyklonalen Wetterentwicklungen intensiver verliefen als bei antizyklonalen. Dabei befanden sich die Objekte bei konstanter Temperatur und Feuchtigkeit und zum Teil auch in hermetisch verschlossenen Gefässen im dunklen Thermostaten.

Soweit sich Ergebnisse dieser Art bestätigen und sichern liessen (16,17,18), sind also irgendwelche Reizwirkungen, ausgehend von Änderungen bekannter Faktoren wie Licht, Temperatur, Feuchtigkeit und Druck, zumindest für die betreffenden Organismen von vornherein aus dem Kreis der Betrachtungen auszuschliessen. Ferner kann das Nervensystem des Menschen oder der Tiere nicht länger als erster und einziger Resonanzort für das unbekannte exogene Agens angesehen werden. Ausser bei Bakterien wurden wetterbezogene Reaktionen bei Pilzen (32) und Algen (37, 38) auch von anderen Autoren festgestellt. Hier handelt es sich aber ebenfalls um nicht gesicherte Feststellungen. Immerhin liegt schon eine gewisse Sicherung in der Bestätigung von Einzelbefunden durch verschiedene Autoren an verschiedenen Objekten. BORTELS (10,11) hatte aus seinen Versuchsergebnissen geschlossen, dass Maxima von Oxydationen, aerober Atmung, generativer Vermehrung und Beweglichkeit bei Bakterien und Pilzen im Laboratorium meistens mit steigendem, Minima mit fallendem Luftdruck verbunden sind und umgekehrt Maxima von Reduktionen, Gärungen, Aggregationen und vegetativem Wachstum mit fallendem und ihre Minima mit steigendem Luftdruck. Entsprechende Feststellungen machte später KISS (37,38) bezüglich des Verhaltens von Mikroorganismen der sog. Wasser- und Schneeböhlen. Die vegetative Massenvermehrung der diese Erscheinung hervorrufenden Algen auf Grund der Assimilation, also Reduktion der Kohlensäure, erfolgte bei fallendem und die Gametenbildung bei steigendem Luftdruck. In jüngster Zeit haben BROWN und Mitarbeiter (20) be-

* Institut f. Bakteriologie,* Biologische Bundesanstalt f. Land- und Forstwirtschaft, Berlin - Dahlem, Germany.

richtet, dass verschiedene Arten höherer und niederer Tiere und Pflanzen unter weitgehend konstanten Versuchsbedingungen einschliesslich Druck in ihrem Sauerstoffverbrauch und ihrer Beweglichkeit parallel zu Luftdruckänderungen exogen beeinflusst werden.

Es gibt jedoch auch Arbeiten, die diese bisher genannten Feststellungen zu widerlegen scheinen. So hat JENSEN (34) die von BORTELS (7) festgestellte Beziehung zwischen AZOTOBACTER-Entwicklung und Luftdruckänderung, nämlich intensive Stickstoffbindung mit entsprechender Entwicklung vor und bei steigendem, schwache oder gar keine vor und bei fallendem Luftdruck, nachgeprüft und keine derartige gesicherte Beziehung gefunden. Dasselbe gilt für die Nachprüfung der von BORTELS (10) behaupteten Abhängigkeit der Nitrifikation vom Wetter durch STÜVEN und ENGEL (54), die nur eine Korrelation zwischen Oxydationsmaxima und steigendem Luftdruck, nicht aber auch zwischen Minima und fallendem Luftdruck feststellen konnten. Möglicherweise war jedoch in diesem Falle die Versuchszeit zu kurz (3 Monate), und in beiden Fällen die Untersuchungsmethode nicht sehr geeignet, worauf die Autoren z.T. selbst hingewiesen haben. Mindestens ebenso wichtig für die Deutung dieser Widersprüche ist die Tatsache, dass JENSEN 1948/49 experimentiert hat, als auch BORTELS (z.B. 11) keine deutlichen Beziehungen der AZOTOBACTER-Entwicklung und anderer mikrobiologischer Reaktionen zu Luftdruckänderungen feststellen konnte. Dagegen hat er sie 1950/51 wieder nachweisen können. Das hat sehr wahrscheinlich seinen Grund in der veränderlichen Sonnenaktivität. Schon die ersten Reihenexperimente, die BORTELS (7) mit AZOTOBACTER als Versuchsobjekt angestellt hat, lassen deutlich erkennen, dass die jährliche Durchschnittsmenge gebundenen Luftstickstoffs von 1933 = 15,8 mg über 1934 = 13,0 mg und 1935 = 11,6 mg bis 1. Vierteljahr 1936 = 11,6 mg bei zunehmender Sonnenaktivität abnahm. Ueberdies wurden die kurzfristigen Schwankungen in der Menge des gebundenen Stickstoffs kleiner und standen seit Winter 1934/35 mit zunehmender Zyklonentätigkeit nicht mehr in so deutlicher Korrelation zu Wetteränderungen wie 1933. Später wurden solche Überlagerungen des Wettereinflusses durch den Sonneneinfluss in bestimmten Jahren und Jahreszeiten auch bei anderen Objekten festgestellt, nämlich beim Gefrieren unterkühlten Wassers und beim Erkranken von Tabak nach Infektion mit PSEUDOMONAS TABACI (11,13,17). Dass die periodischen Schwankungen der Sonnenaktivität mit Lebensvorgängen unmittelbar in Beziehung zu bringen sind, ist an mancherlei Objekten von verschiedenen Autoren gezeigt worden, obwohl auch hier vieles nicht ausreichend gesichert ist. Es genügt, wenn in diesem Zusammenhang auf die kritischen Betrachtungen von BERG (5) verwiesen wird. Auf Grund der allerdings noch mangelhaften Erfahrungen auf diesem Gebiet scheinen die Beziehungen mikrobiologischer Reaktionen zu Luftdruckänderungen immer dann am deutlichsten zu sein, wenn die Fleckenaktivität der Sonne langsam abnimmt oder ein Minimum erreicht, und am undeutlichsten bei der raschen Zunahme und im Maximum der Fleckentätigkeit.

Inzwischen hat sich gezeigt, dass die zeitliche Abstände zwischen biologischen und zugehörigen meteorologischen Ereignissen nicht immer konstant sind, also z.B. bei der Verwendung der Synchronisations- oder "n"-Methode in den mittleren Luftdruckdifferenzkurven um die biologischen Stichtage "n" manchmal Phasenverschiebungen vorkommen (16,17). Deshalb können bei der statistischen Auswertung einer grösseren Zahl von Einzelfällen mit einigen zeitlich abweichenden Reaktionsabständen tatsächlich vorhandene meteorobiologische Beziehungen durch Interferenz ausgelöscht werden. Da sich während des steilen Anstiegs der Sonnenfleckenkurve die Verhältnisse entsprechend schnell ändern, ist es schon deshalb verständlich, warum in dieser Phase des Fleckenzyklus die Beziehungen biologischer Reaktionen beispielsweise zu Luftdruckänderungen im Synchronisationsverfahren weniger deutlich in Erscheinung treten als während des allmählichen Abfalls der Fleckenkurve. Ausserdem tritt bei dem betont zyklonal turbulenten Wetter vor und in einem Fleckenmaximum während der ganzen Zeit oft nur ausschliesslich und gleichbleibend der zum Tiefdruckwetter gehörige Typ einer biologischen Reaktion auf, wie auch umgekehrt der zu Hochdruckwetter gehörende Typ nach dem Fleckenmaximum während des Maximums der erdmagnetischen Aktivität in gleichbleibender Stärke vorherrscht, weil weitere Steigerungen bei der Natur des Objektes nicht möglich sind (12,15).

Neben diesen Störungen in den genannten Korrelationen zwischen den Reaktionen der Versuchsobjekte und meteorologischen Vorgängen soll noch eine weitere Erwähnung finden, die besonders wichtig erscheint. Zuvor aber ist es nötig, ausdrücklich darauf hinzuweisen, dass nicht nur lebende Zellen, sondern auch unbelebte physikalisch-chemische Systeme von dem Wetteragens in entsprechend verschiedener Weise je nach zyklonaler oder antizyklonaler Wetterentwicklung beeinflusst werden. BORTELS (12) hat beim Ansetzen von Nährlösungen bemerkt, dass Niederschläge von Calciumphosphat bei zyklonalen Wetterentwicklungen schnell und grossflockig ausfallen, bei antizyklonalen dagegen langsamer, in feinkörnigen Streifen. Ähnlich verhält es sich mit der Synthese von Agar, Serum (10), Arsentrisulfid (31,41), Blut (23,44) und anderen hydrophilen Kolloiden (42, 45,47), deren Teilchen bei zyklonalem Wetter leichter aggregieren als bei antizyklonalem. Beziehungen zur Sonne sind besonders deutlich gesichert von PICCARDI und anderen (3,22,47) an der wechselnden Sedimentationsgeschwindigkeit bei der Wismutoxychlorid-Fällung nachgewiesen worden. Es kann folglich auch nicht überraschen, dass Kristallisationen aus übersättigten Lösungen und das Gefrieren kleiner Mengen unterkühlten Wassers gleichfalls solche Wetterabhängigkeiten zeigen. 5 ml in Ampullen eingeschmolzenen Wassers gefrieren bei -6° C vor und bei steigendem Luftdruck leichter als vor und bei fallendem Luftdruck (12,13). Aus allen diesen Feststellungen muss

geschlossen werden, dass das unbekannte meteorobiologische Agens über mikrophysikalische Reaktionen in der Zelle oder ihrer nächsten Umgebung zur Wirkung kommt. Diese Wirkungen sind nicht nur biologisch von Bedeutung, sondern spielen auch im Bereich der Technik eine Rolle (18,40).

Bei den Versuchen mit unbelebten, also extrem einfachen Systemen, liessen sich deren Beziehungen zu meteorologischen Vorgängen durchweg signifikant nachweisen. Hier hat sich auch besonders klar herausgestellt, welche Bedeutung verschiedener, räumlich über den Versuchsobjekten vorhandener Materie zukommt. Nach TZSCHASCHEL und BERGTER (56) ist die Wirkung der über dem Versuchsobjekt vorhandenen, vermutlich als Absorber oder Reflektor wirkenden Materie abhängig von ihrer Masse nach Art einer ROSSI-Kurve. PICCARDI (46) dagegen macht die Wirkung nicht von der Masse, sondern von der spezifischen elektrischen Leitfähigkeit des Materials abhängig. Er findet dünnes Kupfer wirksamer als dickes Blei. Doch ist noch nicht sicher, ob das immer zutrifft. Jedenfalls müssen bei allen Experimenten etwa vorhandene Absorber oder Reflektoren gebührende Beachtung finden. In nächster Nähe des Objektes sind sie kontrollierbar. Andere aber sind es nicht immer, z.B. die Dunstglocke über einer Stadt (16) oder sonstige atmosphärische Qualitäten und ihre Veränderungen. Sie kommen als Störungsursachen für synchrone meteorobiologische Untersuchungen an verschiedenen Orten in Betracht.

Das also das Wetteragens nachweislich nicht nur lebende, sondern auch unbelebte Materie in wässriger kolloidaler oder echter Lösung beeinflusst, war nicht nur eine unmittelbare Wirkung auf Mikroorganismen, sondern auch eine mittelbare über das Nährsubstrat von vornherein anzunehmen. Tatsächlich liess sich diese Annahme verschiedentlich bestätigen (14,15,16,17,18). Physiologische und morphologische Eigenschaften von Mikroorganismen werden dadurch verändert, dass das für sie bestimmte Nährsubstrat bei seiner Auflösung in Wasser vom Wetteragens in charakteristischer Weise beeinflusst wird. Dieser Einfluss ist wie der unmittelbar auf die Zellen erfolgende so, dass die Lösung unter zyklonalem Wettereinfluss umgekehrte Wirkungen auf die sich in solchem Substrat entwickelnden Mikroorganismen zur Folge hat wie die Lösung unter antizyklonalem Wettereinfluss. Die Wirkung kommt wahrscheinlich so zustande, dass entsprechende Stoffwechseltypen einer Bakterienpopulation auf oder in dem durch das Wetteragens in der einen oder anderen Weise veränderten Substrat einen Selektionsversprung erhalten und so die Eigenschaften der ganzen Kultur prägen. Natürlich können auch bei Nichtbeachtung dieser Wirkungen vorhandene meteorobiologische Beziehungen durch Überlagerung der beiden verschiedenen Einflussmöglichkeiten infolge Interferenz mehr oder weniger ausgelöscht werden. Hierin liegt eine weitere und, wie mir scheint, eine der wichtigsten Störungsursachen, die bisher keine Beachtung gefunden und zu Kontroversen über die Realität meteoromikrobiologischer Beziehungen geführt hat.

Der Substrateinfluss des Wetteragens auf die Farbstoffbildung einiger CHROMOBACTERIUM-Arten und anderer Mikroorganismen konnte so weit geklärt werden, dass nunmehr in diesem mikrobiologisch wichtigen Prozess nahezu experimentelle Unabhängigkeit vom Wetteragens besteht. Das bei den einschlägigen Versuchen verwendete Nährsubstrat enthielt ausser Glucose und Pepton Dikaliumphosphat, Magnesiumsulfat, Ferrosulfat und Calciumcarbonat. Die noch nicht veröffentlichten Versuchsergebnisse haben bewiesen, dass der Einfluss des Wetteragens auf CHROMOBACTERIUM VIOLACEUM und anderen Mikroorganismen über den Nährboden nur dann sehr deutlich zu erkennen ist, wenn die Salze für jeden Versuch frisch in Wasser gelöst werden. Diese Forderung gilt nicht in gleichem Masse für die organischen Komponenten des Substrats. Die Wetterentwicklung muss also vornehmlich über die anorganischen Bestandteile verlaufen. Eine getrennte entsprechende Untersuchung der 4 Salze ergab, dass das Phosphat vor Magnesium- und Ferrosulfat den wesentlichsten Faktor darstellt. Es sei daran erinnert, dass Calciumphosphat je nach Wetterentwicklung schneller oder langsamer, grobflockig oder feinteilig streifig ausfällt. Es ist anzunehmen, dass auch mit Magnesium- und Eisenphosphat ähnliches geschieht. Da unter zyklonalem Einfluss, wenn Phosphat grobflockig ausfällt, CHROMOBACTERIUM VIOLACEUM auf entsprechendem Nährboden rau und tief violett wächst, wäre theoretisch dasselbe infolge einer Verminderung der Phosphatkonzentration und damit der Zuckerphosphorylierung und Atmung zu erwarten. Tatsächlich trifft das auch bis zu einem gewissen Grade zu. Farbstoffbildung ist oft eine Folge gehemmter Kohlehydrat- und verstärkter Eiweissatmung (Autolyse). Die Kulturen wachsen seit einigen Jahren mit sehr wenig Phosphat rau und tief violett und mit viel Phosphat glatt und heller violett, jedoch nicht farblos. Das farblose Extrem wird zur Zeit bei den reduktionsbegünstigenden starken Tiefdruckwirkungen (15) erst dann erreicht, wenn dem Nährboden ausser viel Phosphat ein Oxydationsmittel wie etwa Chinon in geeigneter Konzentration zugesetzt wird. Entsprechend können bei starken oxydationsbegünstigenden Hochdruckwirkungen tief violetten Kulturen auf demselben Substrat nur dann erzielt werden, wenn ihm ausser extrem wenig Phosphat ein Reduktionsmittel wie etwa Sulfid zugesetzt wird. Dieselben Beziehungen zwischen Populationswechsel (Selektion morphologisch und physiologisch verschiedener Typen) und Atmungs-stoffwechsel haben in ganz anderem Zusammenhang auch ALTENBERN und Mitarbeiter (1) gefunden. Die Substratwirkungen des Wetteragens scheinen also ebenfalls sowohl über Fällungen als auch über Oxydationen und Reduktionen, und zwar über Beeinflussungen der Atmung zu verlaufen, entsprechend dem eingangs erwähnten Wirkungsunterschied zwischen zyklonalen und antizyklonalen Wetterentwicklungen. Die sich aufdrängende Frage, ob dieser Wirkungsmechanismus auch für die unmittelbare Beeinflussung des Zellinneren durch das unbekannte Wetteragens gilt, bedarf noch weiterer Untersuchung und Klärung.

Nachdem somit im Verlauf mikrobiologischer und physikalisch-chemischer Experimente eine Reihe von Störungsfaktoren bekannt geworden sind, wird es unter Beachtung dieser Erkenntnisse nunmehr möglich sein, noch stichhaltigere Beweise für die Existenz meteorologisch-biologischer Beziehungen zu liefern. Das ist unter anderem auch für die epidemiologische Forschung besonders wichtig. Hier sei ein Wort zum Grundsätzlichen im meteoropathologischen Geschehen erlaubt: Unter dem Einfluss zyklonaler Wetterentwicklung können aerobe Organismen, auch der Mensch, in ihrem Stoffwechsel und der Funktion ihrer Organe gehemmt werden. Diese Organe können dann auch erkranken (4, 25, 26, 43, 53). Fällungen, z.B. im Blut, sind dann stark begünstigt, die Gefahr von Thrombosen ist erhöht (23, 44). Folgt nun antizyklonales Wetter, dann tritt in jeder Beziehung das Gegenteil ein. Der Stoffwechsel wird intensiver, die Organe erholen sich, Thromben lösen sich und können Embolien hervorrufen. Das wird durch zahlreiche korrelationsstatistische Untersuchungen belegt (23, 33). Diese Abhängigkeit der Atmung und Stoffwechselintensität gilt aber auch z.B. für aerobe parasitische Bakterien, mit dem Unterschied allerdings, dass sie als Einzeller anscheinend schneller reagieren. So treffen beim Uebergang von zyklonalem zu antizyklonalem Wetter schon aktivierte Bakterien auf noch nicht abwehrbereite Wirtsorganismen und machen diese krank. Wenn dann aber wieder der Uebergang zum zyklonalen Wetter erfolgt, können sich die nun schon geschwächten Bakterien des inzwischen stark gewordenen und noch anhaltend starken Gegenangriffs des Wirtsorganismus nicht mehr erwehren, und die Infektionskrankheit kann heilen, die Seuche erlöschen. Diese Erkenntnisse wurden aus den Ergebnissen von Modell-experimenten mit menschlichem Speichel und einem saprophytischen Bakterium abgeleitet (9) und fanden bis jetzt sowohl durch Versuche mit pflanzenpathogenen Bakterien (17) als auch durch statistische Untersuchungen an umfangreichem Material von Infektionskrankheiten des Menschen (28) weitgehende Bestätigung. Ebenso fügt sich die Feststellung von TSCHESCHE und Mitarbeitern (55) über die Bildung von Peroxyden als Bakterieninhibine im menschlichen Speichel gut in den allgemeinen Rahmen der gegebenen Theorie ein. Anaerobe pathogene Bakterien und die von ihnen hervorgerufenen Krankheiten stehen selbstverständlich in anderer Beziehung zum Wetterverlauf als die Aeroben, und auch Viren scheinen sich anders zu verhalten. PRIMAULT (49, 50) hat z.B. festgestellt, dass die spontanen Ersterkrankungen an Maul- und Klauenseuche unter Berücksichtigung der Inkubationszeit mit einem markanten Luftdruckfall verbunden sind, und CORDES (24) fand dasselbe für die Grippe. Wenn alle diese verschiedenen Wirkungen des unbekannten Agens auf Wirt und Parasit berücksichtigt werden, dann können auch die Beziehungen des Seuchengeschehens zu meteorologischen Ereignissen deutlicher hervortreten.

Natürlich hat trotz Klärung so mancher Teilfragen vieles noch ungeklärt bleiben müssen. Auch die Kernfrage nach dem wirksamen Agens kann z. Zt. noch nicht beantwortet werden. Alles, was hierzu von mikrobiologischer und physikalisch-chemischer Seite festgestellt werden konnte, deutet auf Strahlung, deren totalen Abschirmung bisher nicht gelungen ist, auch nicht 1000 m unter der Erdoberfläche. Beziehungen zur Sonnenaktivität, zeitlich weltweite Uebereinstimmung bestimmter Reaktionen (12, 47, 57), Sonnenrichtungs- und Abschirmeffekte (12, 44, 46), Tages- und Jahreszeitenrhythmen (7, 12, 47) und manches andere sprechen für eine variable Sonnenstrahlung, die nach DÜLL (27) und KÖPPE (39) sowohl das Wetter (Luftdruck) als auch nach mancherlei Umwandlung in der Atmosphäre das Leben auf der Erde beeinflusst. Auch haben u.a. EUGSTER und HESS (30), TZSCHASCHEL und BERGTER (56) sowie KELENTEY und KELETI (35) biologische Wirkungen der kosmischen Strahlung und ihrer Schauer hinter Blei nachgewiesen. Die Schwankungen des P-Testes nach PICCARDI (45) korrespondieren mit denen der kosmischen Strahlung. Bakteriennährböden können nach WYSS und Mitarbeitern (58) durch UV- und Röntgenstrahlen in ähnlicher Weise beeinflusst werden wie durch antizyklonale Wetterentwicklung. Nach BROWN und Mitarbeitern (19) bestehen gesicherte Beziehungen zwischen der Atmungsgrösse und Intensität der Umgebungsstrahlung, die nach BORTELS (15) z.B. auch zum Phosphat-Fällungstyp in Beziehung steht. Es ist nun nötig, dass in dieser Richtung weiter vorgestossen und das Problem noch enger eingekreist wird. Möglichkeiten hierzu sind beim Experimentieren mit mikrobiologischen und physikalisch-chemischen Objekten durchaus vorhanden. Man sollte versuchen, von Einzelmessungen zu laufenden Registrierungen überzugehen, wie es NEUWIRTH (41) schon mit Erfolg versucht hat, damit auch kurzfristige Ereignisse und das Verhalten während der Nacht erfasst werden. BERGTER und NOACK (6) halten das noch unbekannte meteorologisch-biologische Agens auf Grund ihrer Experimentalergebnisse für Korpuskularstrahlung. Sie nehmen zwei verschiedene, positiv und negativ geladene Teilchen mit verschieden grosser Durchdringungsfähigkeit an. Die negativen Korpuskeln könnten für die zyklonalen und die positiven für die antizyklonalen Wirkungen verantwortlich gemacht werden. Jedoch gibt es noch kein physikalisches Messverfahren für diese Teilchen, so dass also ihre Existenz noch nicht sicher bewiesen ist. Manche andere Versuchsergebnisse, z.B. zeitliche Uebereinstimmungen mit magnetischen Stürmen, von TAKATA (57), BORTELS (11, 15), PICCARDI (45, 46) und BURKARD (22) sprechen aber ebenfalls dafür. DÜLL (26), KIEPENHEUER und Mitarbeiter (36), BRÜZEK (21) und REITER (52) nehmen dagegen Wellenstrahlung als Ursache an. Korrelationen zwischen dem Auftreten von Infralangwellen (51) und bestimmten biologischen Reaktionen oder die Wirkung elektrischer Felder im Experiment (48, 52) sind aber noch kein schlüssiger Beweis dafür, dass Feldstrahlung auch immer die meteorologischen Reaktionen unmittelbar auslöst. ELMERT und Mitarbeiter (29) haben die Versuchsergebnisse von KIEPENHEUER und Mitarbeitern nicht bestätigen können, und auch BORTELS konnte bisher keine Wirkung von cm- und m-Wellen auf die von ihm geprüften Objekte feststellen. Möglicherweise

könnte es daran liegen, dass das Auftreten bestimmter Wellenstrahlen nur ein dem Agens parallel geschaltetes Symptom wäre, oder dass die genannte Teilchenstrahlung als Agens durch elektrische Felder verändert wird, oder auch dass für verschiedene Versuchsobjekte und Reaktionen verschiedene Agentien, teils Wellen, teils Korpuskeln, in Betracht kommen. Alle diese grundsätzlichen Fragen sind jetzt einer erfolversprechenden Bearbeitung mit Hilfe mikrobiologischer und physikalisch-chemischer Versuchsobjekte besser zugänglich geworden, und es ist zu hoffen, dass auf diesem Wege bald weitere Ergebnisse erzielt werden.

L I T E R A T U R V E R Z E I C H N I S

1. ALTENBERN, R.A., WILLIAMS, D.R., KELSH, J.M. and MAUZY, W.L.: Metabolism and population changes in *Brucella abortus*. II. Terminal oxydation and oxygen tension in population changes - J. Bact. 73, 697-702, 1957.
2. ASSMANN, D. : Die Wetterfühligkeit des Menschen - Jena 1955.
3. BECKER, U. : Ein Zusammenhang zwischen der Fällung von Wismutoxychlorid (anorganischer Test von Piccardi) und der Sonnentätigkeit - Arch.f. Meteorol., Geophys. u. Biokl. Beiblätter 10, 1-4, 1943.
4. BERG, H. : Wetter und Krankheiten - Bonn 1948.
5. BERG, H. : Solar-terrestrische Beziehungen in Meteorologie und Biologie- Leipzig 1957.
6. BERGTER, F und NOACK, D.: Über eine auf biologische und physikalische Systeme wirksame Komponente der kosmischen Strahlung (KS) - Die Naturwiss. 43, 208, 1956.
7. BORTELS, H. : Meteorobiologische Untersuchungen an *Azotobacter* - Zentralbl. für Bakt. II. 102, 129-153, 1940.
8. BORTELS, H. : Meteorobiologische Reaktionen einiger Mikroorganismen - Zentralbl. f. Bakt. II. 105, 305-325, 1942.
9. BORTELS, H. : Ueber Beziehungen zwischen epidemiologischem und meteorologischem Geschehen, unter besonderer experimenteller Berücksichtigung der Inhibinwirkung - Zentralbl. f. Bakt. II. 104, 289-325, 1942.
10. BORTELS, H. : Mikrobiologischer Beitrag zur Klärung der Ursachenfrage in der Meteorobiologie - Arch. f. Mikrobiol. 14, 450-508, 1948.
11. BORTELS, H. : Beziehungen zwischen Witterungsablauf, physikalisch-chemischen Reaktionen, biologischem Geschehen und Sonnenaktivität - Die Naturwiss. 38, 165-176, 1951.
12. BORTELS, H. : Zwei einfache Modelle meteorobiologischer Reaktionen in ihren Beziehungen zu Luftdruckänderungen und zur Solaraktivität - Arch.f. Meteorol., Geophys. u. Bioklim. 5, 234-257, 1954.
13. BORTELS, H. : Das Gefrieren unterkühlten Wassers in Beziehung zu interdiurnen Luftdruckänderungen und zur Solaraktivität - Arch.f. Meteorol., Geophys. u. Bioklim. B. 7, 269-276, 1956.
14. BORTELS, H. : Beziehungen zwischen Wetterwechsel, Atmung und Farbstoffbildung bei Mikroorganismen - Zentralbl. f. Bakt. II. 109, 330-335, 1956.
15. BORTELS, H. : Die hypothetische Wetterstrahlung als vermutliches Agens kosmo-meteoro-biologischer Reaktionen - Wiss. Zeitschr. d. Humboldt-Univers. Berlin, 2, 116-123, 1956/57.
16. BORTELS, H. : Synchronbeobachtungen über Beziehungen physikalisch-chemischer und mikrobiologischer Reaktionen zu Luftdruckänderungen in und bei Braunschweig 1953/54 - Arch. f. Meteorol., Geophys. u. Bioklim. Im Druck.
17. BORTELS, H. : Ueber Beziehungen zwischen der durch *Pseudomonas tabaci* (Wolf u. Foster) Stevens hervorgerufenen "Wildfeuer"-Erkrankung des Tabaks, Luftdruckänderungen und Solaraktivität - Phytopath. Zeitschr., 33, 403-425, 1958.
18. BORTELS, H. und KOPPE, H.: Ueber die Beziehungen zwischen den Wirkungsgraden von Jute-Bandspinnmaschinen und der Farbstoffbildung durch *Chromobacterium violaceum* sowie der interdiurnen Luftdruckänderung, der erdmagnetischen Aktivität und der Grenzfrequenz der sporadischen E-Schicht - Melland Textilber. 37, 1011-1016, 1956.

19. BROWN, Fr.A.Jr., SHRINER, J. and WEBB, H.M.: Similarities between daily fluctuations in background radiation and O_2 -consumption in the living organism. - Biol. Bull. 113, 103-111, 1957.
20. BROWN, Fr.A.Jr., WEBB, H.M. and MACEY E.J.: Lag-leadcorrelations of barometric pressure and biological activity. - Biol. Bull. 113, 112-119, 1957.
21. BRUZEK, A. : Die veränderlichen Strahlungen der Sonne. - Wetter und Leben, 3, 33-36, 1951.
22. BURKARD, O. : Einige statistische Versuche zur Piccardischen Fällungsreaktion. - Arch. f. Meteorol., Geophys. u. Bioklim. B. 6, 506-510, 1955.
23. CAROLI, G. und PICHOTKA, J.: Weitere Untersuchungen zur Beziehung zwischen Blutgerinnung und Wetter. - Arch. f. Meteorol., Geophys. u. Bioklim. B. 5, 403-412, 1954.
24. CORDES, H. : Die Grippe und ihre Beziehung zu atmosphärischen Einflüssen. - Arch. f. Meteorol., Geophys. u. Bioklim. B. 6, 462, 1955.
25. CURRY, M. : Bioklimatik. - Riederau / Ammersee 1946.
26. DULL, B. : Wetter und Gesundheit. I. Wissenschaftliche Forschungsberichte-54, Dresden und Leipzig 1941.
27. DULL, G. and B. : The behaviour of barometric pressure during and after solar particle invasions and solar ultraviolet invasions. - Smith. Misc. Coll. 110, 1948.
28. DONLE, W. : Jahreszeit und Witterung im Seuchengeschehen. - Stuttgart 1956.
29. ELMERT, G., REINERT, J. und MITLACHER, W.: Zur Frage des Einflusses von Meterwellen auf Keimungs- und Wachstumsvorgänge bei Pflanzen. - Zeitschr. f. Naturforschg. 7b, 109-112, 1952.
30. EUGSTER, J. und HESS, V.F.: Die Weltraumstrahlung (Kosmische Strahlung) und ihre biologische Wirkung. - Zürich u. Leipzig 1940.
31. FINDEISEN, E. : Experimentelle Untersuchungen über den Einfluss des Witterungsablaufs auf die Beständigkeit eines Kolloids. - Bioklim. Beibl. 10, 23-34, 1943.
32. FISCHER, F. und GAUMANN, E.: Biologie der pflanzenbewohnenden, parasitischen Pilze. - Jena 1929.
33. HALSE, Th. und LOSSNITZER, H.: Untersuchungen über den Einfluss der Witterung auf das fibrinolytische Potential als Beitrag zur Frage einer Meteorotropie der Thrombosegefährdung. - Dtsch. med. Wschr. 74, 790-793, 1949.
34. JENSEN, H.L. : Periodical variations in nitrogen fixation by Azotobacter. - Plant and Soil. 2, 301-310, 1950.
35. KELENTEY, B. and KELETI, T.: Effect of cascade showers on the respiration of Staphylococcus aureus Duncan. - Zentralbl. f. Bakt. I. Orig. 171, 457-460, 1958.
36. KIEPENHEUER, K.O., HARTE, C. und BRAUER, I.: Über die Wirkung von Meterwellen auf das Teilungswachstum der Pflanzen. - Naturwiss. 36, 27-28, 1949.
37. KISS, I. : Die meteorobiologische Analyse der periodischen Massenproduktion einiger vegetabilischer Mikroorganismen, Bakterien und Chlorobacterien. - Ann. Biol. Univers. Hungariae. Pars Szegediensis. 387-395, 1951 (Deutsche Zusammenfassg.).
38. KISS, I. : Überprüfungsuntersuchungen der meteoropathischen Erscheinungen im Leben pflanzlicher Mikroorganismen. - Időjárás 61, 15-21, 1957 (Deutsche Zusammenfassg.).
39. KOPPE, H. : Sonnenaktivität, Grosswetter und wetterbezogene Reaktionen. - Ann. Meteorol. 4, 87-96, 1951.
40. LUYKEN, E. und WEGENER, W.: Der Witterungseinfluss auf die Belastbarkeit gedehnter Kunstseiden. - Melliand Textilber. 33, 767-770 und 876-878, 1952.
41. NEUWIRTH, R. : Atmosphärische Einflüsse auf die Alterung eines anorganischen Sol.-Meteorol. Rundsch. 7/8, 108-113, 1955.
42. NEUWIRTH, R. und HUMMEL, K.: Wettereinflüsse auf das Lichtstreuvermögen von kolloidalen Lösungen (Polyvinylpyrrolidon). - Arch. f. Meteorol., Geophys. u. Bioklim. B. 5, 388-402, 1954.

43. PETERSEN, W.F. : Man - Weather - Sun. - Springfield, Ill., 1947.
44. PICCARDI, G. und DAN'TI, M.: Wirkung einer Metallabschirmung, ein biologischer Test. - Wetter und Leben 7, 7, 1955.
45. PICCARDI, G. : Neue Resultate mit dem anorganischen Test zum Studium von Umwelt-einflüssen - Ein vereinfachter Test (P-Test). - Wetter und Leben, 7, 7, 1955.
46. PICCARDI, G. : Sopra un test inorganico per lo studio degli influssi atmosferici e cosmici. - Geofis. e Meteorol. 3, Nr. 1/2, 1955.
47. PICCARDI, G. : Über einen Zusammenhang zwischen atmosphärischen und kosmischen Phänomenen und physikalisch - chemischen Prozessen. - Arch. f. Meteorol., Geophys. u. Bioklim. B. 6, 486-505, 1955.
48. PICCARDI, G. und CINI, R.: Azione di un campo elettromagnetico della frequenza di 10 KHz sul test chimico: Ol problema dell'influsso degli atmosferici. - Geophisica e Meteorologia 6, Nr. 1/2, 1956.
49. PRIMAULT, B. : De l'influence des variations de la pression atmospherique sur l'apparition de la fièvre aphteuse. - Schweizer Arch. f. Tierheilkunde. 97, 412-422, 1955.
50. PRIMAULT, B. : Eléments météorologiques sur l'apparition et l'extension de la fièvre aphteuse. - Schweizer Arch. f. Tierheilkunde. 100, 383-399, 1958.
51. REITER, R. : Neue Untersuchungen zum Problem der Wetterabhängigkeit des Menschen. - Arch. f. Meteorol., Geophys. u. Bioklim. B. 4, 327-377, 1953.
52. REITER, R. : Gibt es einen direkten Kausalzusammenhang zwischen raschen elektrischen oder elektromagnetischen Feldänderungen und biometeorologischen Reaktionen ? - Medizin-Meteorologische Hefte, 2, 35-59, 1954.
53. DE RUDDER, B. : Grundriss einer Meteorobiologie des Menschen. - 3. Aufl. Berlin-Göttingen-Heidelberg 1952.
54. STUVEN, K. und ENGEL, H.: Die Nitritoxydation durch Nitrobacter winogradskyi Buch. in ihrer Beziehung zu einigen Wetterfaktoren. - Arch. f. Mikrobiol. 24, 347-361, 1956.
55. TSCHESCHE, R., KORTE, Fr. und BETHGE, J.: Über die Rolle des Wasserstoffperoxyds bei der Inhibition nach Dold. - Zeitschr. f. Naturforschg. 6b, 22-24, 1951.
56. TZSCHASCHEL, R. und BERGTER, F. : Untersuchungen zur Wirksamkeit der kosmischen Ultrastrahlung (KS) auf Mikroorganismen. - Naturwiss. 42, 649, 1955.
57. TAKATA, M. : Über eine neue biologische wirksame Komponente der Sonnenstrahlung. Beitrag zu einer experimentellen Grundlage der Heliobiologie. - Arch. f. Meteorol., Geophys. u. Bioklim. B. 2, 486-508, 1951.
58. WYSS, O., HAAS, O.F., CLARK, J.B. and STONE, W.S.: J. Comp. Physiol. 35. Suppl. I. 133-140, 1950. Nach KAPLAN, R.W., Auslösung von Farbsektor - und anderen Mutationen bei *Bacterium prodigiosum* durch monochromatisches Ultraviolett verschiedener Wellenlängen. - Zeitschr. f. Naturforschg. 7b, 291-304, 1952.

GENERAL BIOCLIMATOLOGICAL DATA

Section F: Miscellaneous data

PRELIMINARY RESEARCH ON THE TRANSPARENCY OF THE ATMOSPHERE AT BYDGOSZCZ

by

Dr. J. Paszyński (Poland)*

During the latter part of 1955 the intensity of direct solar radiation was investigated at Bydgoszcz.

The object of this study was to determine the influence of the urban environment upon the transparency of the atmosphere. Measurements were carried out simultaneously in several parts of the town by means of Górczyński actinometers. The results obtained were the basis for calculation of the value of the new turbidity factor Θ (according to F. Linke).

In addition to the differentiation of atmospheric transparency in the town area, resulting from air pollution by industrial smoke, the existence of a characteristic diurnal variation of the turbidity factor Θ was also found. The highest values of Θ were observed in the morning and evening, whereas a minimum was reached at noon, as is clearly shown in the diagram attached.

The curve shown differs distinctly from the variation diagram of Θ quoted by F. Linke ("Handbuch der Geophysik", VIII) for Frankfurt a/M., which is situated in similar conditions to those at Bydgoszcz. However, it should be mentioned that Linke's diagram relates to the summer months.


On the basis of W.G. Kastrow's data and according to the formula of F. Möller, the share in atmospheric extinction of the following factors has also been calculated: 1. molecular scattering, 2. selective absorption due to water vapour and 3. scattering and absorption by aerosols. The following table shows the relative share of each of the three above mentioned factors at noon only on a few related days during the period under discussion. The value of the molecular scattering has always been accepted as 1.

Relative Share of Water Vapour (d_{vI}) and Aerosols (d_{aI})
in Atmospheric Extinction at Bydgoszcz, compared with
Molecular Scattering (d_{mI})

date	d_{vI}	d_{aI}
9 September 1955	1.04	1.50
12 September 1955	0.93	1.70
21 September 1955	0.86	1.82
25 September 1955	0.93	0.90
30 September 1955	0.83	0.57
2 October 1955	0.80	0.97
3 October 1955	0.74	0.77
4 October 1955	0.84	0.77
12 October 1955	0.79	0.61
19 October 1955	0.69	0.74

* Institute of Geography, Polish Academy of Sciences, Warszawa, Polska.

This article could be classified also under part IV, Section C4c: Chemical pollution.



Average Diurnal Variation of the Turbidity Factor [H]
at Bydgoszcz in September and October 1955.

Section G : World literature

PART II

PHYTOLOGICAL BIOCLIMATOLOGY

(1959)

Section A : General phytological bioclimatology

Section B : Agricultural bioclimatology

- 1. General agricultural bioclimatology**
- 2. Agricultural phenology**

PHYTOLOGICAL BIOCLIMATOLOGY

Section B1: General agricultural bioclimatology

SEAWATER IRRIGATION

A NEW LINE OF RESEARCH ON A BIOCLIMATIC PLANT- SOIL COMPLEX

by

Dr. Hugo and Elisabeth Boyko (Israel) *

I. REPORT ON PRINCIPLES
AND PRELIMINARY EXPERIMENTS

In the following paper an experiment is described in which two plant species with extremely different ecological requirements were successfully grown on dune sand under direct irrigation, with various dilutions of seawater, during a period of approximately $1\frac{1}{2}$ years. This experimental period included two seasons of drought, one of about three months before the winter rains of 1957/58, and one of almost nine months between the winter rains of 1957/58 and 1958/59.

This experiment was carried out in order to test our theory that the direct use of seawater for irrigation is possible in many cases, provided that accumulation of salts, particularly NaCl, or the domination of sodium and chloride ions in the root layer can be prevented.

According to an extensive review organized by Unesco on "Utilization of saline water" (23) based on a bibliography of hundreds of papers and books, this appears to be the first experiment of its kind. In this review, in the chapter on "Regional Climatic Problems", water of marine origin is also mentioned. One sentence is all that is said; namely: that in this case the real problem is not how to use, but how to get rid of this saline water (23a, p. 21).

Before describing our experiment, it is therefore justified for us to deal not only with the underlying scientific principles, but also to show their gradual development and stages.

2. HISTORY

Many trials have been carried out with salt water, possibly since prehistoric times, when man first started to irrigate crops in the neighbourhood of salty waters by flooding fields in arid regions. Archeological records excavated by H. Helboek in Irak and numerous other data elucidate the gradual salination of alluvial soils there during a period of about 2000 years.

It is worthwhile quoting in this connection, one of the "parallel lines of evidence".

Pottery excavated from sites in Southern Irak dating from about 3500 B.C. suggest that at that time the proportions of wheat to barley were nearly equal.

A little more than 1000 years later, the less salt-tolerant wheat accounted for only one-sixth of the crop - and in 2100 B.C. for even less than 2 per cent. By 1700 B.C. the cultivation of wheat had been abandoned completely. The average yield had shrunk from 2537 litres per hectare in about 2400 B.C. to an average of only 897 litres per hectare about 700 years later (1700 B.C.).

This information may throw new light also on the existence of the abandoned cities and settlements in the Negev. It also supports the suggestion that there have been no large climatic changes during the last 3000 years in these desert parts of Israel, now in process of reclamation.¹⁾

*) Dr. Hugo Boyko, Ecological Advisor, Ministry of Agriculture, Israel.

Dr. Elisabeth Boyko, Senior Research Officer, Agric.Res.Sta., Rehovot, Ministry of Agriculture, Israel.

For reasons of economy all photographs are printed at the end of this paper.

1) See also H. Boyko (6) and the discussion on climate fluctuations at the UNESCO-Symposium on Hydrology in Ankara, 1952.

In those times, salination of the once fertile soils was, in itself, enough to cause increasing poverty and depopulation, not to speak of the numerous wars to win new arable lands.

It is surprising enough that salination was already detected in those early times; but to prevent it was an unsolvable problem then. In such arid areas fields of wheat and barley are flooded up this day by collecting the run off waters and leading them onto terraced fields. Direct precipitation alone does not suffice. The damage caused to agricultural soils by flooding with seawater however was always catastrophic and known as such.

It is therefore understandable that, after these unfortunate experiences for thousands of years, man apparently never conceived the idea of using salt water in just those areas of the barren shifting sand dunes where, even without such dangerous irrigation, almost nothing could grow.

For this purpose a synoptic and synthetic approach, such as ecology has adopted during these last few decades, was essential.

We feel obliged however to stress the true importance of our simple experiment and deprecate the sensational storm that was created in the world press (see also 11).

We connected well-known geophysical principles with well-known phytobiological principles. We combined general climatic data and meteorological measurements of wind, spray-salt and sand movement. Furthermore the figures of temperature, drought and precipitation were linked with the physical phenomena of rapid percolation and good aeration in sand dunes on the one hand, and the recognized phytobiological phenomena of slower water intake by roots in saline solutions, higher osmotic pressure, etc., on the other.

Instead of proceeding step by step; increasing salinity and taking normal agricultural soil as our mainstay, we adopted a new approach in our experiments. We began from the other extreme; namely, with the highest amount of seawater, the highest available concentration through the longest possible dry period and on the most barren places, the shifting sand dunes.

3. FORMER STUDIES

The idea which prompted the following experiment was based on various previous studies. We had carried out similar and extensive studies in the "Sand- and Salt-Steppes" of the "Neusiedler-See" in Burgenland, Austria, during the years 1929 - 1935 (4,5,7), and supplemented them by many additional investigations on the coasts of the Baltic Sea, the North Sea, and on many parts of the Mediterranean Coast. Many observations on halophytic and psammophytic plant communities were also made in Germany, Southern France, Anatolia, South West Asia, Egypt and, since 1936, extensively in Israel. These have shown that islands of non-halophytic plant communities frequently thrive in the midst of areas of pure halophytic vegetation. Closer investigations has made it clear that the soil in these islands is of a sandy or a gravelly texture or else percolation is facilitated by other favourable circumstances (e.g. artificial drainage).

Apart from the marked permeability of sand and gravel, the idea of considering barren sand dunes was further influenced by the established fact that the main danger from sodium ions in irrigation water grows with the amount of colloidal clay particles in the soil, since these ions are adsorbed by the latter and make the soil impermeable to water by deflocculation. The sands of shifting dunes are obviously particularly poor in such clay particles and provided the best basis for our experiment.

In addition to these physical facts, certain phyto-physiological phenomena were also considered. Halophytic and drought resistant plants show a similarity in that, quite generally, both are assumed to need a relatively high environmental osmotic value. Another important fact taken into consideration was the slowing-down of water intake by the roots with higher concentrations. The influence of temperature and other climatic factors on all physiological processes played another and often decisive role.

For these reasons dunes and gravel soils gave the best prospects for experiments with certain plants to be grown and irrigated with water of high salt concentration. Economic aspects of the species chosen were also taken into account, and priority was given to species of higher economic value.

One of the first steps in this direction was the hydro-engineering experiment of A. Werber (27) on light agricultural soil without plants but with irrigation water of 800 mg/l Chlorine content, then thought to be of exceptionally high salinity (1936). In recent years a few cautious experiments were even made with diluted seawater.

Prof. E.R. Purvis (New Jersey), described his experiment in a letter to Prof. H. Heimann. During the dry summer 1954/55 he irrigated crops on a fertile soil with a dilution of oceanic

water (14,000 mg total salt content in four applications of 1 inch each, at weekly intervals, in addition to normal crop treatment, irrigation and rainfall) and found no practical damaging.

Experiments of a similar nature but on a bigger scale are being systematically worked out at the U.S. Salinity Laboratory in California (Correspondence of December 1958 with Director H.F. Hayward. See also ref. no. 23 a and 23 b). It is to be hoped that the details of all these important experiments will be published in the near future so that they can be combined with the results of our experiment to assist further progress.

More research is being carried out on Desalination (Demineralisation). An extensive review of the Advisory Committee on Arid Zone Research of UNESCO (24) "On Saline Water Conversion" reveals that 85 scientific Institutes are working on this subject. A short review and survey of the various methods of desalination is also to be found in an article by W.G.V. Balchin (1) in "The New Scientist".

4. THE DESERT GARDEN OF ELATH

Before we began to experiment with seawater and dune sand, we took the opportunity to make preliminary investigations (E. Boyko, 2,3).

In summer 1949 one of us (E.B.) was entrusted with the task of planning a desert garden at Elath, the southernmost point of Israel in the Negev on the Red Sea. The climate there is that of absolute desert, with less than 25 mm annual rainfall and its sparse vegetation is confined to Wadis and to the "Sabha", i.e. depressions near the very saline groundwater. These depressions have a more or less heavy soil, whereas the surrounding hills are built of gravel rock debris, and are completely devoid of vegetation.

We did not choose the depressions, the "Sabha", for planting, in spite of the temptation offered by the many halophytic bushes (of *Suaeda fruticosa*, *Tamarix* spp., of date palms, etc.) growing there already, but we chose the bare hills on account of their high permeability. This can be seen from their physical soil structure in Table I:

Table I. DESERT GARDEN OF ELATH: Mechanical soil analysis

Stones (Granite and Limestone)	61.0 %
Coarse Sand	23.0 %
Fine Sand	12.3 %
Silt	2.7 %
Clay	1.0 %
	<hr/> 100.0 %

Elath is situated in one of the hottest and driest areas of the world. The mean monthly temperature of July is about 33° C., of January 15° C., (mean yearly relative humidity is 39.1 %, with strong and persistent winds, mainly from the north, less frequently from the south). Day temperature in the shade frequently reaches 43° C., but the nights are mostly cool.

The great success achieved there with saline water of a fluctuating salt content of about 2000-6000 mg/litre (mainly sulphates) can be seen from the two pictures: Fig.1 and 2, showing the start of our experiment and Fig. 3, showing the same area nine years later. Today it is a splendid public garden in an absolute desert region, and it represents a most valuable recreation centre for the desert harbour town of Elath. It has been irrigated continuously with water containing up to 6000 mg/litre total salt content.

THE EXPERIMENT WITH SEAWATER: Encouraged by this success, we tried to obtain financial support for our main idea in order to experiment with much higher extremes on a bigger scale and under more exactly controlled conditions (Investigation of percolation and salt-accumulation by exact instrumentation (lysimeters, etc.), of solution intake by plants with isotopes, using as many species and varying treatments as possible). After years of unrewarding applications for the necessary support, we began to experiment without any specific budget. For this reason the investigation was simplified to the very minimum and involved personal sacrifices both of time and of money.

We used empty asphalt barrels, we cleaned them and made holes in the bottom, thus achieving quick percolation. The barrels were filled with sand from the dunes near Bat Yam, South of Tel Aviv: its composition can be seen from Table IV. We refrained from painting the dark barrels in order to save expense, but we measured the temperatures in the rootlayer 2 cm apart from the fringe on all sides and in the centre at a depth of 7 cm. There were well defined differences in the daily temperature curves, but no differences could be observed in the development of the plants in consequence of this factor. Another arrangement of plants and another scale of experi-

ment would have been necessary to achieve a statistical basis for such conclusions.

In order to imitate a certain dune height and a similar percolation we stood the 90 cm high barrels upright, after making many small holes of about 1 sq. cm in their bottoms. Later we took out the entire bottoms and put the barrels into furrows filled with sand, thus achieving a depth for percolation of 1.30 m. This enlarged by far the potential number of species to be experimented with. With these simple and cheap methods all species with a root system of about 1 meter or less (most of the cereals for instance, and many others) could be investigated.

In order to overcome to a certain degree the restriction in the choice of plant species, we chose two with very different natural habitats: the first (*Agropyrum junceum* Beauv.), a highly xerophytic pioneer of the lightest soils, the shifting sand dunes; and the other (*Juncus arabicus* Asch. et Buch.), a swamp plant and halophyte, known from the salty swamps on very heavy alluvial soils. We collected the seed material of *Juncus* from the Oasis Jotvata ("Ein Ghadian") in Wadi Araba and that of *Agropyrum* from the coastal dunes near Haifa.

This wide ecological range enables conclusions reached to be applied to a great number of other species, in the first instance to all xerophytes and to all species with a certain degree of salt tolerance. Our experience in Elath and the soil analyses made during this experiment with seawater show, however, that the potential range of species may be still wider. This is so, particularly for experiments with brackish waters, which frequently occur in desert regions with dune areas or with soils of a similar good permeability (e.g. some hammadas, gravelly hills, old Wadi beds, etc.).

Both species were also chosen from the point of view of their potential economic aspect.

Juncus arabicus is a valuable cellulose plant which has been shown to be excellent for making high quality writing and printing paper.

Agropyrum junceum is a fodderplant of high nutritive value (see Table V). In addition, this grass is closely related to wheat, and successful cross-fertilizations have often been made (N. I. Vavilov, 25). Hybrids between *Triticum* and *Agropyrum junceum* have also been found in nature. (J. Rousseau, 21).

This indicates that, once genetical work has begun, there may be good prospects for achieving a perennial, drought resistant and salt tolerant cereal for arid areas under saline irrigation. The number of chromosomes in our Mediterranean Ecotype of *Agropyrum Junceum* is 28, the same as in many wheat varieties (C.D. Darlington and E.K.J. Ammal, 12).

The length of the dry season during which seawater or brackish water can be used without damaging plants is decisive for certain species, since it is necessary to utilize the hot season for the development of most plants. In order not to risk a too lengthy dry season, but still to use that part of the hot season necessary for its development, we started the experiment with *Juncus* at the end of July 1957; that is, about two to four months before the end of the dry season. Thereafter it was expected that effective rains would wash away any accumulation of salt in the root layer (Compare Fig. 4).

Our main aim at this stage of the experiment was not to follow the specific development of particular plant species, but to discover what plants could exist at all in view of the small accumulation of salt expected.

The actual accumulation, following regular drenching with highly concentrated seawater solutions, was to be ascertained by soil-analyses after a specific period, and we wanted this to be influenced by root layers from the very start.

Agropyrum was sown on the 9th of February 1958, in a manner similar to that in a densely sown wheat-field. This date was chosen because we observed that, in its natural habitat on shifting dunes germination occurred during the rainy season approximately at that time.

The three dry months (see fig. 4) which we utilized in summer and autumn 1957 already indicate a possible application for many other less resistant species in large parts of the Mediterranean and other coastal areas of the globe.

Most areas of coastal dune however have a short season of complete drought. A much smaller proportion including Israel, experience a long dry period. The same relation applies approximately also for inland regions.

Rainfall distribution in coastal areas can be seen from the map (Fig. 5) and from Table II.

Table II. APPROXIMATE SEASONAL DISTRIBUTION OF RAINFALL (PRECIPITATION) IN COASTAL REGIONS (Expressed as a proportion of the total coastal length)

Coastal lines in	with a dry season of	
	about 3 months or less	more than 3 months
	approximately	approximately
Australia	1/2	1/2
Africa	1/2	1/2
Asia	3/5	2/5
America	4/5	1/5
Europe	6/7	1/7
All Continents except Antarctica	2/3	1/3
Coastal lines except the Arctic	5/8	3/8

Although providing a rough estimate only, the map and the table show that approximately two thirds of all coast lines have no dry season at all, or one of less than three completely dry months. Even if we exclude the Arctic coast lines as irrelevant, we still find 5/8 of the remainder with these more humid conditions. The results of the first three months of our experiment apply, therefore, to about two thirds of all coastal areas with highly permeable soils and to extensive inland regions. The second dry period of our experiment is about 9 months and, even if the number of species may be restricted by this, the principle may be applied to almost all arid regions as well as to other areas.

For logical reasons the coast line of the Antarctic continent and for technical reasons those of most of the islands and inland lakes are not taken into this rough account, although many of them (e.g. Lake Tsad, Lake Aral) are situated in arid regions with dune areas. On the other hand, in more humid places the need for saline water does not yet exist. If, however, we read articles such as that by C.J. Jackson (18) about the surprising rise of water use near great cities, then we see that this problem is not without significance even in humid regions (Areas of soils other than dunes, with high permeability; may also reach considerable dimensions and can be computed from the soil maps of each region).

The total area of coastal dunes cannot be estimated here. The total of all dune-areas is officially stated as 3,200 millions of acres; that is twice as much as the total area and seven times as much as the agricultural area of the United States of America. (Ch.Y. Whitfield and R. L. Brown, 28).

5. WATER - AND SOIL - ANALYSES

It may be remarked here that the total salt and NaCl concentration in the seawater along the coast of Israel is higher than that of many parts of the ocean. The following data illustrate this:

The specific weight of oceanic waters varies from approximately 1.023 to 1.031 according to the salt content. Near the coast this content may differ, mainly as a result of evaporation, precipitation and the influence of rivers. The average total salt content of oceanic water, according to G. Grillo (23a) is 35%.

The following figures give a rough idea of average seawater composition (see also E.D. Howe, 23c, for comparison).

NaCl	26,900 mg/litre
MgCl ₂	3,200 " "
Na ₂ SO ₄	2,200 " "
CaSO ₄	1,400 " "
KCl	600 " "
Various	100 " "
	<hr/> 34,400 mg/litre

Some other examples are: The total salt content of oceanic water taken near Point Pleasant, New Jersey is 28,000 mg/litre according to a letter from Prof. E.R. Purvis, New Jersey.

The North Sea near the dunes of Holland has approximately 17,500 mg/litre and near Denmark even

less. The Caspian Sea has a total salt content of about 13,000 mg/litre, decreasing from South to North (The Karabuga Bay with its extremely high salt concentration is almost completely disconnected from the Caspian Sea itself). The Aral Sea has about 10,000 mg/litre, the Baltic Sea only 6000 mg/litre total salt content.

The salt content in the Mediterranean Sea is much higher and increases from West to East. Its average total is about 37,000 mg/litre, that of the eastern parts near Crete 39,900 with a NaCl content of about 30,000 mg/litre. The NaCl content of the seawater actually used in these experiments is about 31,700 mg/litre, and the total salt content 43,600 mg/litre.

According to theoretical consideration and to our observation in nature we expected that, after several months' irrigation with seawater, a small salt-accumulation, particularly of NaCl not used by the plants and not washed down, would be found in the root layer. This would be eliminated from time to time by fresh water irrigation, or by the winter rains. The results of the analyses, however, exceeded our expectations. The figures of barrels 4 to 5a, show that even after 1½ years of the experiment and after a daily irrigation with 100% seawater during the dry season, there was no accumulation of salt whatsoever. The highest concentration of NaCl in the dry soil of the root layer, i.e. in barrel 5a, was necessarily much smaller (minimum 6160, maximum 10,480 mg NaCl/litre) than the amount added by the last single irrigation the preceding day (31,700 mg NaCl/litre). In this case, however, we still had a moisture content of 14.7% - 15.0% in the soil sample at the time of its analysis.

Four analyses of the seawater used, taken always from the same spot on the coast, gave almost identical results:

Table IIIA. SEAWATER-ANALYSES *)

Seawater from the Mediterranean Sea near Bat Yam, South of Tel-Aviv (Jaffa). The water from this spot was used in our experiment during the entire period (July 1957 up to date, 31. December 1958). Date of analysis: 7. August 1958.

Total Salt Content:	43,600 mg/litre		
Analysis			
	Na ⁺	: 12,500 mg/litre	or meq (= milliequivalent)
	K ⁺	: 455 " "	
	Ca ⁺⁺	: 725 " "	
	Mg ⁺⁺	: 1,230 " "	
	SO ₄ ⁻⁻	: 3,118 " "	
	Cl ⁻	: 22,750 " "	
		40,778 mg/litre	
Calculated amounts:	CaSO ₄	2,965 " "	or meq
	MgSO ₄	1,713 " "	
	MgCl	3,530 " "	
	KCl	870 " "	
	NaCl	31,700 " "	
	Various	2,822 " "	
		43,600 mg/litre	

Table IIIB. DILUTIONS USED IN THIS EXPERIMENT

Barrels	Dilution	Total salt-content	NaCl	comparable to:
1	Tapwater from the Citrus-			
1 a	grove at Beth Dagon			normal drinking-water
2	25 % Seawater	1.1 %	0.8%	Aral Sea (1%)
2 a	75 % Tapwater			Caspian Sea (1.3%)
				>Baltic Sea (0.6%)
				>most brackish desert waters
3	50 % Seawater			
3 a	50 % Tapwater	2.2 %	1.6%	North Sea (1.7%)
4	75 % Seawater			
4 a	25 % Tapwater	3.3 %	2.4%	Oceanic concentration
5				
5 a	100% Seawater	4.4 %	3.2%	East-Mediterranean Sea

*) The analyses were made at our request in the Chemical Laboratory of the Ministry of Agriculture, Agricultural Research Station, Rehovot.

These figures can be compared with the average salt content in the saline waters of oceanic and other origin mentioned previously.

Even the concentration of 75% seawater used successfully in the barrels No.4 and 4a of our experiment was, therefore, similar or higher than that of pure seawater in many parts of the ocean and much higher than that of many salty seas, salt lakes and other brackish waters; e.g. the North Sea, the Caspian Sea, the Baltic Sea, etc., and far higher than that of most of the underground-waters in deserts and other regions.

There are species which are able to absorb a part of the salt content of saline irrigation water. Others are not able to do this. Accumulation can be foreseen especially in experiments with the latter group of species in less permeable basins, beneath inland dunes, and in coastal regions where there is insufficient rainfall and/or no outlet for the underground-water to the Sea.

Geological research and regional hydrological and pilot hydro-engineering experiments must be combined in such cases with ecological investigation. Premature conclusions without such specific and regional experiments must be avoided.

But, after proving that accumulation in the root layer is not to be feared - the main research must now be aimed at achieving the most nutritive solution for each species. Pure sea-water in various concentrations was intentionally used in this first experiment. It should not be overlooked, in this connection, that seawater has been the medium for plant and animal life through most of geological time and that all essential trace substances are probably contained in it.

These facts, together with possibility of occasionally using partially fresh water for direct irrigation or combined with many other varied treatments, enlarges considerably the potential number of species and the applicability of the principle in general. When we have cheap energy at our disposal to divert seawater to inland dunes, then the prospects will appear very great indeed. In any case, even if this first experiment had been less satisfactory, it would have justified the recommendation of experiments along these lines in all countries with sand-dunes, gravelly or similar soils.

The most important consequence of the absence of any accumulation in the soil is that the decisive limiting factor for the growth of any plant species with salt water irrigation on dunes and similar soils, depends upon their specific ability successfully to use water of that chemical composition and specific concentration. The danger of accumulation which is the factor prohibiting the growing of field crops irrigated with even slightly brackish water on normal agricultural soils was completely prevented by the principles devised here.

According to existing formulae (as, for example, that quoted by G. Grillot, 23a) and the empirical experience, greater amounts of irrigation water must be used with higher concentrations; but we did not rely on formulae for the reason that with such unusually high concentrations the existing empiric formulae may be misleading.

For practical use the energy problem will in any case be an important factor. But, these problems and their solutions must be tackled by technicians after the results of theoretical research and adequate scientific experiments have paved the way.

The complete lack of accumulation in our experiments can be seen from Table IIIC, which indicates the results of soil-analyses after about a year's irrigation with various dilutions of seawater up to approximate oceanic concentration (75%) and to the still higher concentration of East Mediterranean Seawater (100%).

Table IIIC. SOIL-ANALYSES

Total Salt and NaCl content in the soil (dune sand) after irrigation with various seawater concentrations.

Beginning of irrigation: 31 July 1957.

Date of Analysis : 2 July 1957 (based on dry weight 105° C.)

Barrel No.	Daily irrigation (see Fig. 2.) with:	Depth of soil samples	Moisture %	Cl %	Calculated NaCl % (Coeff. 1.6)	Total sum of salts %
1	0 % SEA-WATER (Tap- water)	10 cm	6.7	0.004	0.0064	0.017
		30 cm	0.5	0.003	0.0048	0.015
		60 cm	1.0	0.004	0.0064	0.015
1		10 cm	6.8	0.013	0.0208	0.032
		30 cm	1.5	0.004	0.0064	0.015
		60 cm	1.0	0.004	0.0064	0.013
2	25 % SEA-WATER NaCl 0.793% Total 1.090%	10 cm	4.6	0.050	0.0800	0.100
		30 cm	4.3	0.061	0.0976	0.120
		60 cm	2.2	0.044	0.0704	0.090
2		10 cm	8.1	0.056	0.0896	0.115
		30 cm	3.8	0.061	0.0976	0.120
		60 cm	1.9	0.046	0.0736	0.095
3	50 % SEA-WATER NaCl 1.585% Total 2.180%	10 cm	1.8	0.097	0.1552	0.190
		30 cm	2.2	0.083	0.1328	0.180
		60 cm	1.5	0.058	0.0928	0.120
3		10 cm	5.8	0.115	0.1840	0.235
		30 cm	4.6	0.098	0.1568	0.190
		60 cm	3.3	0.110	0.1760	0.210
4	75 % SEA-WATER NaCl 2.378% Total 3.270%	10 cm	1.8	0.138	0.2208	0.280
		30 cm	1.8	0.115	0.1840	0.235
		60 cm	2.4	0.075	0.1200	0.145
4		10 cm	5.0	0.160	0.2560	0.300
		30 cm	2.4	0.107	0.1712	0.220
		60 cm	2.9	0.080	0.1280	0.255
5	100% SEA-WATER NaCl 3.170% Total 4.360%	10 cm	11.9*	0.358	0.5728	0.675
		30 cm	14.1	0.463	0.7408	0.950
		60 cm	15.9	0.460	0.7360	0.900
5		10 cm	10.6	0.510	0.8160	0.950
		30 cm	13.2	0.465	0.7440	0.900
		60 cm	15.3	0.540	0.8640	0.950

* The high moisture content of these six analyses of barrel 5 indicates a still lower salt content, if the samples were taken a few hours later and with the same low moisture content than those of the other barrels. The control barrels gave similar results.

6. EXPERIMENTAL DATA OF PLANT DEVELOPMENT

As mentioned above, cleaned asphalt barrels formerly used for road building were stood upright and adapted to allow percolation. Sand was taken from the dunes of Bat Yam in its natural composition in order to imitate natural conditions as far as possible. Table IV shows the composition of this dune sand.

Table IV. MECHANICAL SAND ANALYSIS

Physical soil-structure analysis in %					Total % of CaCO ₃	Water Solution 1:5		% Organic Matter	pH
>1.0 mm	1.0 -0.50 mm	0.50-0.25 mm	0.25-0.10 mm	<0.10 mm		Total Soluble Salts	Cl		
0.0	1.0	15.0	78.8	3.5	3.9	0.090	0.044	2.3	8.6

The reasons for choosing *Juncus arabicus* and *Agropyrum junceum* as our test plants are given above. Furthermore we had familiarised ourselves with their ecological amplitudes under natural conditions as well as with their potentialities for artificial propagation and cultivation. This was brought about by extensive observations during 30 years in many countries and by experiments over a period of 10 years in Israel (H. Boyko, E. Boyko and D. Tsurriel, 10).

Furthermore both are perennials and live for many decades. Observations on the same individuals can therefore be continued for a very long time, thus excluding unknown specific factors which might occur in the case of short-living plants which would have to be replaced during the course of long term experiments.

Our method of planting the *Juncus* seedlings allowed continuous observations on the individual plants, whereas *Agropyrum* was densely sown and successfully grown in imitation of a field crop.

AGROPYRUM JUNCEUM BEAUV.:

Detailed data of the experiments with individual *Agropyrum* plants as well as with some other species, will be dealt with in another paper. Figs. 6 and 7 indicate only the general behaviour of *Agropyrum junceum*. Nevertheless we are publishing these figures here because they demonstrate a practical application of fodder production in sand-dunes with a brackish water supply that may have a high economic value.

The photographs (Fig. 6a and 6b) and the graph (Fig. 7) show the development of *Agropyrum* in the same barrel. This barrel was irrigated from February 1958 with 11,000 mg/litre total salt content and 7,900 mg/litre NaCl. This corresponds with the concentration of many seas and of highly saline underground water in deserts.

In order to estimate the size of this foddercrop on 18.1.1959 we cut the stalks of 1 sq. dm (10 x 10 cm) at a height of 1 cm above the soil-surface. The yields of this square decimeter (100 square centimeters) weighed 116.80 grams freshweight and 75.62 grams air dry. If calculated to a field area this would be equivalent to the yield of a very rich fodder crop and/or pasture.

From the curve (Fig. 7) it can be seen that growth did not stop throughout the whole year.

The Chemical Laboratory of the Ministry of Agriculture kindly supplied us with the following chemical Analysis of *Agropyrum junceum*.

Table V. CHEMICAL FODDER ANALYSIS
OF AGROPYRUM JUNCEUM

Water	72.6 %
Fats	1.1 %
Cellulose	8.5 %
Protein	4.1 %
Ashes	3.3 %
Carbohydrates	10.4 %
	<hr/> 100.0 %

(Beta Carotin Provitamin A 1.8 mg)

Fieldexperiments under most unfavourable conditions on salty, sandy soils and on sand dunes with and without irrigation have already been carried out successfully in our dune station and in other places, and the plant could now be introduced into the economy of several settlements with great success (D. Tsurriel, 22. See also Ref. nr. 10).

The high nutritive value and palatability, the high yield in spite of most inadequate spacing, lack of fertilizing, etc., and the success achieved with this perennial in the first season permit us to recommend this strain of *Agropyrum* as a very promising fodderplant on dunes or similar sterile, permeable soils, particularly in latitudes of less than 40°, even in situations where irrigation with highly brackish waters only is available.

OSMOTIC VALUES:

A few remarks should be added here in respect to our measurements of osmotic values both for the dilutions and of the two species.

Osmotic values had to be measured with the strictest precautions and economy of material, as every measurement brought about the destruction of a considerable part of the total material available. We shall therefore draw no conclusions from the results owing to the necessarily small amounts of data.

But, owing to the most interesting figures which they represent, we reproduce here the results of the latest measurements (Table VI).

Table VI. OSMOTIC VALUES (measurements of 10.1.1959)

Dilu- tion		Crioscopic Temp.	Osmotic Value
	Distilled water	- 0.55 ° C	
1)	Tap-Water	- 0.6 " "	0.6 Atm.
2) 25 %	Sea-Water	- 1.1 " "	6.63 "
3) 50 %	" "	- 1.55 " "	12.06 "
4) 75 %	" "	- 2.05 " "	18.09 "
5) 100 %	" "	- 2.55 " "	24.12 "

Dilu- tion	Species	No. of barrel	Osmotic Value
1) 0 %	Agropyrum	1	5.43 "
2) 25 %	"	2	7.12 "
1) 0 %	Juncus	1	2.05 "
2) 25 %	"	2	2.41 "
3) 50 %	"	3	4.34 "
4) 75 %	"	4a	4.824 "

The xerophytic dune plant *Agropyrum junceum* shows a high osmotic value even when irrigated with freshwater and this value rises with higher concentrations.

The hydrophilous-halophyte *Juncus*, however, shows relatively low osmotic values and seems to withstand the high osmotic pressure and any loss of moisture induced thereby. Exact physiological experiments may bring about results of physiological and physical significance.

Obviously the influence and importance of the osmotic values in plant life need to be thoroughly re-investigated (compare H. Walter, 26, and various textbooks).

JUNCUS ARABICUS ASCH. ET BUCH.:

As for *Juncus*, 10 seedlings were planted in each barrel and all the seedlings in each barrel were mapped on a simple sketch and measured. Thus, the measurements of all plants could be followed individually and any unforeseen occurrences (dwarfing, dying, abnormal development of any part, etc.) could easily be checked in relation to the whole development and possibly their causes determined.

Fig. 8 shows as an example the sketch of one day's measurements.

Two barrels were equally treated with the same concentration and 10 seedlings were planted on July, 31st, 1957 in each of the eight barrels 1 to 4a. All the seedlings were two months old and had been grown in our own nursery, irrigated with fresh water. They were transplanted at the same age but of two different sizes:

5 plants in each barrel were about the size of 3 - 4 cm and 5 plants in each barrel the size of 7 - 10 cm, the seedlings of both groups having 3 - 4 leaves each (see also Fig. 8 and its text).

This was done in order to find out, in this first experimental step, if the taller or the smaller seedlings would be more apt to adapt themselves to high concentrations of seawater.

In the first stages there was ample space in each barrel for each of the ten tiny individual plants. The distance was 15 - 20 cm between the single seedlings. But, later on, when almost all roots and shoots grew surprisingly well there was great root competition, a factor which minimized a possibly greater success.

For 10 days after transplanting the seedlings were watered with tap-water in order to re-root them properly in their new habitat. Nevertheless, a few plants and also a few shoots were lost in the course of this transplantation. Of the 40 larger seedlings transplanted on July 31st, 1957, 3 seedlings or 7.5% were dead on 10. VIII. Of the 40 smaller seedlings transplanted on 31. July, 8 seedlings or 20% were dead on 10. VIII; that is, on the day before we began the seawater irrigation.

It can be assumed that the initial fresh-water irrigation after transplantation could be eliminated and substituted by rains if the time of transplanting coincides with rainfalls, either shortly before or after. In our case, however, (with *Juncus arabicus*) high temperatures seemed decisively favourable for re-rooting.

After these first 10 days all the barrels received an irrigation of sea-water or seawater-dilution according to the scale of concentrations mentioned before (Table IIIB) and marked in the graphs.

After we had observed that plants lived vigorously even with 75% Seawater irrigation (= 33,000 mg/litre total salt content), we began, at the end of October, to plant *Juncus* seedlings into barrels using 100% seawater irrigation. But, this time most of the plants died during the transplantation period, in spite of freshwater irrigation. The season seemed unfavourable for beginning this experiment. We continued, however, to irrigate the remaining plants with 100% Seawater. All the plants which survived the transplantation are growing slowly and are developing new shoots up to date but the results are not comparable and are therefore excluded from our conclusions regarding the plants in the other barrels. The only certain conclusion that can be drawn from this part of the experiment is that, even under irrigation with 100% seawater (that is, in this area, 43,600 mg/litre total salt content and 31,700 mg NaCl/litre) all plants that survived transplantation are also surviving this extreme salt concentration with daily irrigation from 1. November 1957 to 31. December 1958.

The further development of the other barrels up to oceanic concentration (barrels 4 and 4a) can be seen from the Figs. 9-11. The first graph (Fig. 4) shows that the first effective rainfall was 13. - 15. November, 1957, but we did not interrupt regular seawater irrigation before the end of November when effective rainfalls occurred over a period of about 8 days. The last effective rainfall occurred on 8. and 26. February, 1958. From then on, seawater irrigation was again the only source of irrigation, apart from a light rain of 0.5 mm on 21. April and of 1 mm on 22. April, 1958, and a momentary sprinkle with tapwater of less than 1 mm on 20. May, 1958, erroneously applied by the gardener.

It was observed with *Juncus* that if all the seedlings of this species were transplanted at the same age, the growth of smaller seedlings (3-4 cm on 31. July, 1957) was slower than that of taller seedlings (7-10 cm on 31. July 1957). Better results can therefore be expected from taller seedlings (For comparison see also Fig. 8).

The number of shoots (leaves) was the same for all seedlings at the time of their transplanting, namely 3 - 4. Nevertheless, the increase in number of shoots was very variable. It may, therefore, be assumed that the tendency to produce many shoots can be successfully augmented by selection. This maximum number of 196 shoots in barrel 4a, irrigated with water of oceanic salt concentration was less than that of the barrels 1 - 3a, but was the same as that of well-developed plants of the same age raised in a field experiment with manuring, spacing, etc. on heavy soil in Ein Hamifrats. The latter were irrigated with water from the river Naaman, south of Acre, containing about 700 mg/litre chloride only.

The maximum number of shoots developed from single seedlings, all of them bearing 3 - 4 shoots (leaves) at the time of transplantation, can be seen from Table VII.

Table VII. *JUNCUS ARABICUS*: MAXIMUM NUMBER OF SHOOTS (LEAVES) OF SINGLE INDIVIDUALS

Sea - Water	0%	25%	50%	75%
Barrel No	1 and 1a	2 and 2a	3 and 3a	4 and 4a
Total Salt	(tapwater)	10,900	21,800	32,700
NaCl		7,900	15,860	23,800
		mg/l	mg/l	mg/l
Date of measurement:				
10 August 1957	3 - 4	3 - 4	3 - 4	3 - 4
3 November 1957	32	39	16	20
12 January 1958	51	40	34	24
2 March 1958	81	93	42	35
2 April 1958	82	119	53	44
18 May 1958	151	124	94	60
25 August 1958	228	177	167	83
25 September 1958	237	246	205	160
31 December 1958	241	257	227	196

The development from 3-4 leaves (shoots) in each individual to the maximum on 31. December 1958 of 257 shoots in one single individual under irrigation with approx. 11,000 mg/l salt content with 8000 mg NaCl/l was certainly remarkable. But, still more remarkable to us was the development of 196 healthy shoots out of such a tiny seedling under irrigation with approx. 33,000 mg/l total salt content and with approx. 24,000 mg NaCl/l; that is, the 75% dilution of the East-Mediterranean seawater which is equivalent to a concentration of oceanic seawater.

It should be added that the measurements described here were concluded with the end of the year

1958, but that all the plants are continuing to show healthy growth in respect of height and number of shoots. This whole development indicates the possibility of achieving much higher yields by breeding, or even by simple selection.

It seems reasonable also to expect still better results if seawater irrigation is interrupted by part-time irrigation with freshwater where feasible. This applies to areas where fresh-water resources - either below or above soil surface - are available though not in sufficient amounts to allow a full irrigation; also, wherever fresh-water reserves must be saved or stored - for example, for town-development; or where there is danger of salination of freshwater by over-charging the underground resources. The latter conditions, for instance, occur frequently in coastal areas.

The fact that we purposely chose the worst possible conditions and achieved positive results shows that improvement in yields will be possible. Such improvements, apart from selection, may include optimal spacing, treatment with growth stimulating substances like gibberellic acid, and the addition of fertilizers.

We have, however, intentionally refrained in this first experiment from any fertilizing - either with organic manure or chemical fertilizers - and from any other additional treatment, in order to eliminate as much as possible influences other than seawater. We have also refrained from the addition of Potash or Calcium to neutralise sodium-ions in order to simplify this first experiment. The lack of financial resources made it imperative for us to restrict this first experiment to the minimum of barrels and technical work and to adhere exclusively to investigation of the principle, or, more specially, to the combination of the respective physical, geophysical and phytobioclimatological principles.

FOR ALL THESE REASONS, THEREFORE, THIS EXPERIMENT REPRESENTS THE MINIMUM OF SUCCESS *)

In experiments of this kind it may be necessary for economic reasons to know the time-retarding effect of higher salt concentrations on plant growth. This may be of less importance in the case of long living plants such as *Juncus arabicus* or *Agropyrum junceum*, for example, but, even with these, it may change the optimal times and/or frequency of harvesting, or the time of the first harvest in general.

In our experiment with *Juncus* this time-retarding effect on minimum height and maximum number of shoots can vaguely be seen in Table VIII. We say "vaguely" because the small size of this experiment indicates only the trend of development, and does not give the desirable statistical basis.

Equally, we have not as yet touched upon the question of the optimal irrigation-quantities with the respective dilutions. We used 5 litres per barrel for each irrigation because with this amount we were certain that water would percolate to and through the bottom of the barrel. We intend to decrease this amount gradually, in order to discover the optimum amount for each species with regard to its various development stages.

It was also clear to us from the beginning that the time of percolation was an important factor for each species, but we did not intend to include different and controlled permeabilities for the reasons mentioned above. It is therefore only by chance that we can now observe the considerable effect of this factor, if only in a general way, particularly in the barrels with higher concentrations, i.e. in barrels 3, 3a, 4, 4a (see Table IX).

*) After this paper was written we received a manuscript from Professor H. Heimann ("Irrigation with saline water and the balance of the ionic environment"), in which are discussed the possibilities of creating balanced composition in the solution. His studies open up further ways for application of our principles. In spite of this, we would not go so far as to say that experiments on demineralisation should be stopped but research work on the utilisation of oceanic water should be approached from both sides. The economical one explained here, as well as the costly experiments on demineralisation should be continued. Our method may lead to success with many species, but with others success may be achieved only if combined with desalination. In this case even desalination could prove to be of economic value, particularly if combined with the methods of counter-balancing the sodium ions, described in the paper of H. Heimann (16) mentioned above.

Table VIII. TIMETABLE OF GROWTH

TIMETABLE OF MAXIMUM NUMBER OF SHOOTS ¹⁾ Period 31.7.1957 till 31.12.1958									
Number of shoots:	4	40	60	80	100	120	150	200	240
Achieved with:	on	on	on	on	on	on	on	on	on
0% seawater (tapwater)	31.7.57	1.12.57	27.1.58	25.2.58	24.4.58	28.4.58	17.5.58	22.7.58	6.12.58
25% seawater	"	12. 1.58	31.1.58	17.2.58	12.5.58	17.4.58	5.7.58	5.9.58	20. 9.58
50% "	"	17. 2.58	10.4.58	4.5.58	25.5.58	22.6.58	1.8.58	20.9.58	not yet
75% "	"	22. 3.58	18.5.58	31.7.58	30.8.58	6.9.58	18.9.58	not yet (10.1.59)	not yet

1) Compare Fig. 10.

TIMETABLE OF MAXIMUM HEIGHT ²⁾ Period 31.7.57 till 31.12.1958								
Max.Height of leaves ³⁾	10 cm	20 cm	30 cm	40 cm	50 cm	60 cm	90 cm	100 cm
Achieved with:	on	on	on	on	on	on	on	on
0% seawater (tapwater)	31.7.57	1.9.57	17.10.57	13.12.57	12. 4.58	2. 5.58	1.10.58	30.12.58
25% seawater	"	1.9.57	17.10.57	15. 2.58	21. 4.58	20. 5.58	2.12.58	not yet
50% "	"	24.9.57	not yet	5. 4.58	8. 5.58	15.11.58	not yet	not yet
75% "	"	15.9.57	20.10.57	higher leaves broken by birds and not comparable			?	?
							(see Fig. 9)	

2) Compare Fig. 9

3) Flowering stems not included (they are much higher)

Table IX. GROWTH DIFFERENCES CAUSED BY DIFFERENT PERMEABILITY

Measurement of 18 May 1958	Irrigation with 50% seawater		Irrigation with 75% seawater	
Barrel No.	3	3a	4a	4
Percolation	slower	quicker	slower	quicker
Average height in cm (of the 4 highest individuals in each barrel on that date)	48.6	33.5	35.4	20.0

We refrained from giving the times of percolation for two reasons. First, we made these measurements with the primitive methods at our disposal, namely by pouring the water in the same measure of time (about 10 seconds) and counting the seconds till the entire surface was again visible. Secondly, conclusions could not be drawn from only two barrels. It should be noted, however, that barrels 3 and 4a always had a slower rate of percolation than the control barrels 3a and 4.

Possible reasons for the different rates of percolation may be chemical or physical or simply that the holes at the bottom varied in size as some of them may have been blocked by a stone or other obstruction. But, with an adequate instrumentation, percolation and the factors affecting it, could easily be controlled in future experiments and the optimal percolation determined.

Part of the explanation of this difference in growth may also be physiological (slower intake with higher concentration). In other words, the rootsystems in the more slowly percolating water may somehow take in the saltwater better and evaluate its meagre nutritient matter, than root systems in the too-quickly percolating water. Irrespective of any additional nutrition given, manure would most probably reduce too rapid percolation. This could, however, also be achieved by other methods, such as by plowing in seaweed, by regulating nutritient solutions, or by increasing the amount of irrigation. These problems are of secondary importance in this initial experiment, but it is worthwhile remembering them in forthcoming experiments with *Juncus* or any other species.

The 90 cm high asphalt-barrels had a diameter of 46 cm ($r^2\pi = 1662 \text{ cm}^2$). The sand-surface was therefore equivalent to about $1/6 \text{ m}^2$ and irrigation with 5 litre was equivalent quantitatively to 30 mm of rain. The exact amount of sand varied with the barrel and was between $1/7$ to $1/6,5 \text{ m}^3$, the volume of the root layer about $1/12 - 1/10 \text{ m}^3$. The sand-surface was kept about 5 cm below the fringe of the barrel; not higher, in order to enable higher rainfalls to percolate and as in nature. They were not kept lower, in order to avoid too much shadow, creating adverse conditions.

The barrels were placed more than 7 m from a white wall, 4 m high, in order to prevent influence by reflex radiation (H. Boyko, 8).

Certain irregularities in the growth rate of the plants in the various barrels, particularly in the development of new shoots, led us to raise the question of whether the cooler months, or quite generally lower temperatures, may not be more favourable for plantgrowth under highly saline than under freshwater irrigation. It may be a general law, or a specific problem for each species. Only the teamwork of bioclimatological, physiological, and bio-chemical research can give the answer. It is possible, for instance, that certain chemical processes connected with growth-retarding may be intensified by higher temperatures. These lines of research, however, transgress the limits of our experiments and of our abilities *).

We hope to continue this work, but in any case the results encourage similar experiments elsewhere. New lines of research in other countries too may result in a rise in the production of potential food and industrial plant raw materials far beyond present expectation.

7. CONCLUSIONS AND SUMMARY

It can be concluded in general, that the experiment described above has proved the soundness of the theory on which it was based.

If highly permeable soils extend deeper than the root layers of the plant species growing in them, then no accumulation of salt is to be feared, even if oceanic water is used for irrigation. Accumulation of salt in the deeper layers of these soils is washed away into even deeper depressions in the desert or into the sea by irrigation, rainfall, or the flow of underground water.

Highly saline water in deserts and sea-water can be used for the cultivation of a number of economic plants if used, not on the usual agricultural soils but ON SAND-DUNES OR SIMILAR SOILS WITH HIGH PERMEABILITY.

In consequence of these basic results the experiment justifies the undertaking of further extensive research. The main lines indicated by these results, include experiments as follows:

- 1) Appertaining to problems of minimal, maximal, and optimal amounts of sea- and saltwater of varied quality, varied mixing, alternating with freshwater, varied timing, control of percolation, aeration, accumulation in correlation with the varied climatic and meteorological factors.
- 2) With numerous prospective species of economic value, with priority for halophytes (salt tolerant plants) and xerophytes (drought resistant plants), under various climatic conditions and under varied treatment in many countries, in coastal regions as well as under continental conditions.
- 3) Bioclimatic, agronomic, agrotechnical and hydrotechnical experiments must be undertaken after the basic ecological results have been achieved with the species or ecotypes to be investigated.
- 4) Genetic experiments (selection, crossing, etc.) to achieve luxuriant plant material of ever increasing salt- and drought-tolerance and shockresistance.

*) With great pleasure we express out thanks to the devoted collaboration of our coworkers in this experiment, to Mr. Arnold Douwes, Chief Gardener, who did the technical work of irrigation and recorded the rainfall on the spot, to Miss Ilse Herzberg, who made the drawings and to Mrs. Adiva Ilan, who carried out the measurements of osmotic values and is now starting experiments with giberellinic acid as growth-stimulants. We owe also particular thanks to the Meteorological Service of the Government for supplying us with a raingauge and with the data of daily Maxima and Minima of temperatures.

- 5) Finally, economic investigation and field experiments on a large scale must be undertaken in order to transfer the results into the economic sphere.

As a result of such experiments, we see good prospects for putting into production enormous stretches of dune lands and other desert areas irrigated with practically limitless amounts of available highly saline water, of many seas up to oceanic concentration and brackish underground waters in all areas, but particularly in arid and semi-arid regions.

The ultimate development will also depend on the development of cheap energy of any kind, particularly of atomic energy and of sun-radiation, for these natural resources are at our disposal in just as limitless quantities as are the waters of the ocean.

8. REFERENCES

1. BALCHIN, W.G.V. : The Purification of Saline Water - The New Scientist, London, 4, 31, 1198-1200, 1958.
2. BOYKO, E. : The Building of a Desert Garden - Journal Royal Horticultural Society, London, 76, 4, March 1952.
3. BOYKO, E. : Plant List of the Desert Garden in Eilat - Div. of Ecology, Ministry of Agric., Israel, Hakirya, Tel Aviv, 1951.
4. BOYKO, H. : Ein Beitrag zur Oekologie von *Cynodon dactylon* Pers. und *Astragalus exscapus* L. - Sitzungsberichte der Akademie der Wissenschaften, Wien, Math.-naturwiss.Klasse, Wien, 140, I, 9 & 10, 1931.
5. BOYKO, H. : Die Vegetationsverhältnisse im Seewinkel, Versuch einer pflanzensoziologischen Monographie des Sand- und Salzsteppengebietes östlich vom Neusiedlersee: II. A) Allgemeines, B) Die Gesellschaften der Sandsuccession - Beihefte, Bot. Zentralbl., Prag-Dresden, 51, 11, 600-747, 1934.
6. BOYKO, H. : On the Climax-Vegetation of the Negev, with special reference to arid pasture-problems - Palest. J. of Bot., Rehovoth, Series, 7, 17-35, 1949. (See also corrections and remarks to on sand-dunes in the same volume inside the 2. coverpage).
7. BOYKO, H. : On the ecological foundations of the regeneration of vegetation in arid zones - Int. Union of Biol.Sciences, Paris, B, 9, 28-31, 62-80, 142-147, 1951. (Excavation of sand dunes with *Haloxylon persicum* on p.68).
8. BOYKO, H. : Introductory lecture to Section II. Climatic Ecoclimatic and Hydrological Influences on Vegetation - Arid Zone Research, Proceedings of the Unesco Symposium in Montpellier, Nov. 1953, Paris, 39-48, 1955.
9. BOYKO, H. : On the role of plants as quantitative climate indicators and the "Geo-Ecological Law of Distribution - J. of Ecol., Cambridge, 35, 138-157, 1947.
10. BOYKO, H., BOYKO, E. and TSURIEL, D.: Ecology of Sand Dunes, Final report to Ford Foundation, Sub-project C -1e - Department of Ecology, Agric.Res.Sta., Rehovoth, Jerusalem, 353-399, 1957.
11. BOYKO, H. : "Statement" in order to rectify certain press-errors - Ministry of Agric., Israel, Nov. 1958.
12. DARLINGTON, C.D., and JANAKI AMMAL, E.K.: Chromosome Atlas of Cultivated Plants - London, 397, 1945.
13. GRILLOT, G. : see UNESCO, 23'a.
14. HALPERIN, RATNER, A.R., and HEIMANN, H.: Sodium and Potassium in the water sources of Israel - Hebrew Institute of Technology, Haifa, 1-18 (and 12 tables) , Nov. 1955.
15. HAYWARD, H.E. : see UNESCO, 23 b.
16. HEIMANN, H. : Irrigation with Saline Water and the Balance of the Ionic Environment - Lecture, Int. Potassium Symposium, Madrid, Sept. 1958.
17. HOWE, E.D. : See UNESCO, 23 c.
18. JACKSON, C.J. : Water Supply and Demand - The New Scientist, London, 4, 82, 150-152, 1958.

19. JACOBSEN, T. and ADAMS, R.M.: Salt and Silt in Ancient Mesopotamian Agriculture - Science, Washington, D.C., 128, 3334, 1251-1258, Nov. 1958.
20. RICHARDS, L.A. ET ALIAS (U.S. SALINITY LABOR. STAFF): Diagnosis and Improvement of Saline and Alkali Soils - U.S. Dept. of Agriculture, Washington, D.C., Agric. Handbook, 60, 1-160, 1954.
21. ROUSSEAU, J. : La Validité des Noms de Quelques Hybrides Intergénériques des Graminées - Mémoires du Jardin Botanique de Montréal, Montréal, 29, 1-23, 1952.
22. TSURIEL, D. : *Agropyrum junceum* as Fodder and Pasture Plant - Hassadeh, Tel Aviv, 9, 613-614, 1954 (in Hebrew).
23. UNESCO : Utilization of Saline Water - Reviews of Research - UNESCO, Arid Zone Research, Paris, 2. Edition, 1-102, 1956.
 - a) GRILLOT, G.: The biological and agricultural problems presented by plants tolerant of saline or brackish water and the employment of such water for irrigation - (Bibliography contains 245 numbers), 9 - 35.
 - b) HAYWARD, H.E.: Plant Growth under Saline Conditions - (Bibliography contains 256 numbers), 37 - 71.
 - c) HOWE, E.D. : Utilizations of Sea Water - (Bibliography contains 307 numbers), 73 - 100.
24. UNESCO, ADVISORY COMMITTEE ON ARID ZONE RESEARCH: Survey of Institutions Engaged in Research on Saline Water Conversion - UNESCO/NS/AZ, Paris, 357, 1-40, 1958.
25. VAVILOV, N.I. : The Origin, Variation, Immunity and Breeding of Cultivated Plants - Chronica Botanica, Waltham, 13, 1/6, 1-366, 1951.
26. WALTER, H. : Grundlagen der Pflanzenverbreitung - Stuttgart, III, I: Standortstheorie, 525, 1951.
27. WERBER, A. : Is irrigation with Saline Water Possible ? - Hadar, Tel Aviv, IX, 9, 201-203, 1936.
28. WHITFIELD, CH.Y., and BROWN, R.L.: Grasses that fix Sand Dunes - in "Grass", the Yearbook of Agriculture, Washington, U.S. Dept. of Agric., 70-74, 1948.



Fig. 1: Organizing the planting of the desertgarden in Eilat by Dr. Elisabeth Boyko in November 1949. (Note the artificial reservoir above). All the plants to be seen are either already planted or prepared for planting. Water was brought by Command-car in barrels from Ber Orah, 18 km North of Eilat.



Fig. 2: First planting on the vegetationless gravel. Details of "methods" and of "soil" show the technical difficulties




Fig. 3: Views of the same Desert Garden at Elath in 1958. (For comparison with Fig. 1 and 2, taken 9 years earlier)





Fig. 4
H. and E. Boyko
Sea-water Irrigation, Rainfall and Temperatures




Fig 5
Distribution of Dry Periods in Coastal Areas



a




b

Fig. 6: SEAWATER IRRIGATION OF AGROPYRUM (Barrel No. 2)
(Agropyrum junceum sown on 9 Febr. 1958) a) left 20. May, 1958
b) right 20. Dec., 1958

The left half of the plants on photo 6b were cut for measurements of osmotic value. The plants were irrigated from 15. February, 1958 with 25% dilution of seawater, i.e. with water of 11,000 mg/l total salt content and 8,000 mg/l NaCl.

Fig. 7

Sea-water-Irrigation (of *Agropyrum junceum*)

Average Maximum of the 10 highest non fruiting stalks (Barrel No. 2)

Irrigation water { 11,000 mg/liter total
8,000 mg/liter Na Cl

Fig. 8


An Example of the Method of Measurement
Barrel No 1a (0%) on 2 March 1958

FIG. 8 gives an example of how the measurements were done. The two figures above the line indicate the height of the two highest leaves in this individual, the figure below the line indicates the number of green shoots on this day.

A single circle is the sign of an individual grown from a small seedling (3-4 cm on 31 July 1957); A double circle indicates an individual grown from a tall seedling (7-10 cm on 31 July 1957).

Fig.9

Sea-water-Irrigation. (Maximum Number of Shoots of *Juncus arabicus*)

Shoots





Fig.10

Sea-water-Irrigation. (Maximum Height of Leaves of *Juncus arabicus*)

mm



Irrigated with

- ▲ - 33,000 mg/liter total Saltcontent with 24,000 mg/liter NaCl
- △ - 22,000 mg/liter total Saltcontent with 16,000 mg/liter NaCl
- - 11,000 mg/liter total Saltcontent with 8,000 mg/liter NaCl
- - Freshwater (tapwater from Pardess Beth Dagon)



Fig. 11: SEAWATER IRRIGATION OF JUNCUS ARABICUS

Comparison of the plants above with those below shows the development of *Juncus arabicus*. Of the 4 barrels above (photos taken on 20. May 1958). 3 are to be seen below (photos taken on 25. Sept. 1958).

- No. 1a (second from the left above, first below)
Irrigated with fresh-water
- No. 2a (middle photo above, second from the left on the first photo below)
Irrigation with 25% seawater (11,000 mg/l total salt content, 8,000 mg/l NaCl)
- No. 3 (is not to be seen above, but shows below its healthy development)
Irrigation with 50% seawater (22,000 mg/l total salt content, 16,000 mg/l NaCl)
- No. 4 The plants were cut in order to measure their osmotic value.
Irrigation with 75% seawater (33,000 mg/l total salt content, 24,000 mg/l NaCl).
- No. 4a (first from the right above and below). The photo below is taken from a greater distance, but shows clearly the healthy development of the plants.
Irrigation with 75% seawater (33,000 mg/l total salt content, 24,000 mg/l NaCl).

PHYTOLOGICAL BIOCLIMATOLOGY

Section B1: General agricultural bioclimatology

SEAWATER IRRIGATION

A NEW LINE OF RESEARCH ON A BIOCLIMATIC PLANT- SOIL COMPLEX

by

Dr. Hugo and Elisabeth Boyko (Israel)

A B S T R A C T

Over a period of many years the authors have frequently observed the occurrence of non-halophytic plant species growing in islands in the midst of halophytic plant communities. Closer investigation has shown that these islands usually appear in places with high soil-permeability, or with good drainage.

In 1949, on the occasion of laying out and planting a desert-garden in Elath, on the Red Sea, this general principle was tested and used in a practical manner.

A garden was laid out and successfully irrigated with water of 2000-6000 mg/l (mainly sulphates) on completely barren slopes of broken rock debris with soil of good permeability. About 180 plant species, many of them of economic value, were introduced into this extreme desert area which has less than 25 mm average yearly rainfall. The garden is used today, as the town's recreation centre and the plants provide the basis for planting in several new settlements in the Wadi Araba.

On the basis of these results an experiment was started under extreme conditions in order to show that no accumulation takes place in the root layer in well-drained soil, even if irrigated regularly with very saline water, provided that the soil is deeper than the rootsystems of the plants grown therein. The experiments consisted of planting two species having extremely different ecological ranges on dune sand. One of the plants, *Agropyrum junceum* Beauv., is a xerophyte of shifting dunes - that is of the lightest soils. The other, *Juncus arabicus* Asch. et Buch., is a halophyte from salt swamps, and the heaviest soils. The economic value of the first lies in its high nutritive value and palatability as fodder; that of *Juncus* in its excellent qualities for papermaking (writing and printing paper).

The plants were irrigated with controlled amounts of various dilutions of East-Mediterranean seawater containing 43,600 mg/litre total salts and 31,600 mg/litre NaCl.

All soil-tests showed that even after 100 applications of seawater of the highest salt concentration (43,600 mg/litre total salt content), no accumulation occurred in the soil-layers at 10 cm, 30 cm and 60 cm depth.

These results therefore indicate that there are great possibilities for growing many plants of economic value with different ecological amplitudes, on sands and soils of similar permeability under irrigation with the large amounts of brackish waters available in deserts and, in the case of certain more salt tolerant species, with seawater. It is advocated that similar experiments should be tried out in other countries in order to coordinate and to develop their far reaching aspects.

Section C : Forest bioclimatology

Section D : Physiological phyto-bioclimateology

Section E : Pathological phyto-bioclimatology

PHYTOLOGICAL BIOCLIMATOLOGY

Section E: Pathological phyto-bioclimateology

METEOROLOGY AND THE TIMING OF FUNGICIDE APPLICATIONS AGAINST
POTATO BLIGHT

by

P.M. Austin Bourke (Eire) *)

1. INTRODUCTION

There has been a striking growth of interest in meteorology as applied to the control of plant disease since 1920, and particularly in the last ten years or so. This has, in large part, been due to the careful attention now being given to the economics of plant protection techniques and to a growing realisation that meteorological advice can contribute largely to bringing down the costs of protective measures, and at the same time increasing the efficiency of control of the disease.

Everyone is familiar with examples of the direct effect of weather on the host plant and on the life-cycles of fungus and insect pests. But even when there is no clear-cut environmental influence on the disease itself, meteorological factors may still play a decisive role through their effect on the population level and activity of vector insects. Thus in England in 1957, the mild preceding winter and dry spring were most favourable to the aphid which spreads the disease of sugar beet called virus yellows and as a result the crop suffered the worst outbreak since sugar beet became a principal crop in England.

In other cases, the connection between weather factors and a plant disease are even more complex. Bruzone disease of rice, for example, is caused by hydrogen sulphide developing in the soil. Weather affects the development of the disease in two ways (Reference 10). Firstly the gas is released from sulphides, reduced from sulphates, by the action of acids developing from the decomposition of organic matter. In the case of a prolonged winter and a cool spring, the decay of organic matter is hindered and the disease is correspondingly held in check. Secondly, the transport of the gas by irrigation water increases with cooler weather at the end of Summer because of the increase in the gas-absorbing capacity of the water with falling temperatures; hence the disease appears generally with decreasing temperatures at the end of Summer.

2. POTATO BLIGHT AND WEATHER

Amongst the plant diseases of major economic importance, potato blight (*Phytophthora infestans*) is undoubtedly the one whose appearance and spread is most completely under the direct control of environmental factors. In analysing the seasonal progress of other plant diseases, e.g. apple scab (*Venturia inaequalis*), weather data must be supplemented by observations on the initial level of inoculum, and on its ripening and dispersal. In most potato growing countries, however, the fungus responsible for potato blight is almost invariably present to a degree sufficient to give rise to an epidemic if only suitable weather occurs and persists. Again, contrary to what happens with many other crops, the host plants are susceptible to the disease at all stages of their development, so that no phenological observations, other than a knowledge that the plants are above ground, are required to supplement weather data. Finally, weather effects on the growth of the potato plant itself play no major role in blight epidemiology; it is permissible to concentrate entirely on environmental influences on the propagation and spread of the fungus.

In summary, then, weather data are a necessary and sufficient basis for forecasting the onset and spread of potato blight, provided the favourable conditions for the multiplication of the fungus can be defined.

*) Assistant Director, Irish Meteorological Service, Dublin. President of the Commission for Agricultural Meteorology, W.M.O.

3. DEFINITION OF "BLIGHT WEATHER"

The first necessary step, then, towards a practical Weather Warning Service against potato blight is to reach a SATISFACTORY DEFINITION IN TERMS OF ORDINARY METEOROLOGICAL OBSERVATIONS of the weather conditions favourable to the progress of the disease. A fair measure of success has been obtained in this objective (Reference 4).

Empirical rules were first developed in England and Holland. At the same time, laboratory experiments in the U.S.A. and elsewhere showed the necessity for a period of at least ten to twelve hours of saturated air within the crop cover, at temperatures of the order of 10° to 15° C, for the formation of blight spores. They also established that the spell of high humidity required to be followed immediately by a period of several hours during which free moisture was present on the foliage of the host plant in order to permit germination of the spores and reinfection of the host. These laboratory results were made the basis of definitions of "blight-weather" developed in the U.S.A., Germany, Holland and Ireland.

The author's system is used in the Blight Weather Warning Service which has been operated with success by the Irish Meteorological Service since 1952 (Ref. 1,2,3,6). It has also proved applicable in Chile (reference 5) and in Tasmania (Reference 11).

4. THE UTILITY OF FORECASTS OF POTATO BLIGHT

The possibility of identifying "blight-weather" had little practical value before the present century, when the practice of the regular spraying of the potato crops with copper or other fungicides was introduced in areas such as Ireland where the disease makes a regular appearance each year. The use of fungicides against potato blight does not eradicate the disease; its effect, rather, is to retard its spread by killing the spores which emerge during favourable weather, and thus, by lengthening the period of growth of green parts of the plant, to ensure a better crop of tubers. It soon became obvious that the timing of the applications of fungicide was all-important both from the point of view of effectiveness and economy. A spray applied after the disease had already taken a firm hold on the crop was too late to exercise any effective control on the subsequent damage. On the other hand, too early spraying lost most of its value because, by the time the disease was moved to activity by favourable weather, much of the protective layer of fungicide had been washed off the foliage and because there had been a substantial growth of new and unprotected leaves. Experience gradually dictated a fairly effective routine viz. first application of fungicide as soon as the earliest spots of the disease appeared in the crop, followed by other sprayings at fairly short intervals to protect the growing foliage.

However, the impact of disease varies very greatly from year to year with weather conditions, and it was clear that it should be possible to develop a more rational and elastic spraying programme based on meteorological information. Such a programme would indicate not only the best date for the initial spraying, but would also take into account the later probabilities of spread of the disease, enabling the number of fungicide applications to be restricted in years unfavourable to the epidemic development of blight and thus saving a substantial amount of money for the farmer.

5. MERE IDENTIFICATION OF "BLIGHT WEATHER" INADEQUATE

It was mentioned, in paragraph 3 above, that adequate rules for the identification of weather favourable to potato blight have been developed in several countries.

However, the satisfactory definition of "blight weather" in quantitative terms still leaves other problems to be solved before it can be applied to the epidemiology of the disease and in particular to the efficient timing of fungicide applications. It cannot be too strongly stressed that the simple use of identified blight weather periods for timing of anti-blight sprays has serious drawbacks, as illustrated schematically in Figure 1. In the three sub-diagrams X represents the date of initial visible appearance of blight in the crop, and XYZ the blight progress curve based on percentage of foliage diseased, using the valuable concept introduced by the English plant pathology school (Reference 9). P_1 and P_2 represent, respectively, periods of weather leading to the first sign of the disease and to significant subsequent progress. In several countries not operating a weather-warning system, a spraying advice (A) is issued to growers immediately following the visible appearance of the disease in the general crop. The possibility of identifying the favourable weather period, P_1 , as it occurs, enables the farmer to be warned (W) of the imminent appearance of the disease some time before it becomes visible on the foliage. This anticipation can be of considerable value if, as in subdiagram A, the next favourable weather period (P_2) follows in quick succession, since it will often give the farmer the extra

notice necessary to permit him to take protective measures before P_2 commences.

However the reverse applies in the case illustrated in sub-diagram B where a lengthy period elapses between periods P_1 and P_2 . Here, with the disease lying dormant for several weeks, spraying even at time A may be too early because of the effects of weathering and growth of new unprotected foliage before the next active period occurs. Spraying at the still earlier date W is even more premature in this case. Clearly for efficient spraying routine it is necessary that the identification of P_1 should be supplemented by means of foreseeing whether or not further favourable periods are likely to occur in the immediate future. Indeed, the ideal situation would be to issue a forecast warning (F.W. in subdiagram C) before the first of a series of favourable weather periods had begun.

6. A BASIS FOR FORECASTING "BLIGHT WEATHER"

Already it is possible, at least in some areas and in some years, to make reliable forecasts of the probable occurrence of weather favourable to potato blight, using the synoptic approach to plant disease forecasting. This involves relating the occurrence of blight-weather periods to the type of meteorological situation on synoptic weather maps which gives rise to them (Reference 6)

As regards NW Europe, the types of synoptic situation which are significant for blight epidemiology are comparatively few and easily identified. Two of the most important blight-fomenting situations in this area are:-

- (1) OPEN WAVES OF MARITIME TROPICAL AIR: The weather which accompanies this airmass, as it arrives over North-west Europe during the potato-growing season, is normally ideal for the development of potato blight, viz. overcast skies with rain or drizzle, saturated or near saturated air and temperatures of the order of 12-16°C.
- (2) STAGNANT OR SLOW-MOVING DEPRESSIONS GIVING LENGTHY PERIODS OF WET OVERCAST WEATHER: Ill-defined quasi-stationary fronts are frequently associated with this type of situation and widespread outbreaks of thunderstorms are not uncommon, particularly in the later stages.

One of the basic advantages of the synoptic approach is that it provides a sound basis for forecasting the probable trend of future weather as related to plant disease and thus overcoming the disadvantages discussed in connection with Figure 1. It also enables us to link potato blight epidemiology with the main body of research work on synoptic meteorology and long-term forecasting, and to take advantage of such concepts as "steering" (which gives a clue as to the probable track of future depressions) and "blocking" (which may check blight by shutting off the supply of maritime Tropical air for a considerable period).

It is true, of course, that despite the progress made, weather forecasting for more than a short period ahead is still fraught with difficulties. However some of the problems which beset general forecasting, e.g. the accurate timing of changes, do not apply in the case of plant disease epidemiology. In the latter case, it is sufficient to know that certain weather types are likely to predominate in the immediate future, irrespective of the exact dates on which they will occur. For this reason it is already possible on occasions to issue a "forecast warning" for potato blight purposes covering a period well in excess of that for which a detailed forecast could prudently be made.

7. SYNOPTIC BLIGHT FORECASTING IN PRACTICE

A few of the turning points in the 1958 blight forecasting system in Ireland may illustrate how the synoptic approach is applied in practice.

Early June showed abnormally high rainfall in most parts of Ireland. From June 13th to 18th, however, was generally dry, but it would have been premature to advise spraying unless it could confidently be said that the weather would soon revert to moist, warm conditions.

Figure 2 shows the surface weather chart of June 17th. A broad south-westerly current of maritime Tropical air covered much of the Western Atlantic, and it was clear that, although the anticyclonic ridge would give fine spraying weather for another day or two, an influx of warm moist air into Ireland was inevitable, and was likely to persist, or recur with short interruptions, probably to the end of the month.

Accordingly the first spraying warning for Ireland was broadcast on June 17th. The area which it was expected would be affected by the first wave of maritime Tropical air is illustrated in Figure 3(a), and what actually developed in Figure 3(b), in which the figures denote the effective duration of blight-weather at each observing station (Ref. 1,2,3,6). This synoptic weather system moved on into continental Europe, as did the immediately following depressions, including

a remarkably deep low on June 25th-26th. The period June 21st-30th gave rise to blight weather generally in potato-growing Europe, as far East as Poland.

Blight made a general appearance in Ireland at the end of June and beginning of July. The first spray warning was particularly well timed, in that it enabled the growers to take advantage of practically the only suitable spraying weather which occurred during the month.

A short lull in Ireland early in July enabled a second warning to be issued on July 7th, after which the weather reverted to conditions markedly favourable to the disease. Figure 4 illustrates the synoptic situation at the surface on July 21st. A ridge of high pressure moving in from the Atlantic promised better conditions for the first time in the 1958 Summer - but how long would they last? A glance at the marked East-West trough on the upper air chart (Figure 5), on which the analysis was confirmed by the upper wind reports shown, supplied a confident answer - barely long enough to enable spraying to be repeated before the bad weather systems appearing on the left of Figure 4 would again bring recurrent bad weather to Ireland.

Accordingly the third spraying warning, in the form of a general alarm issued by newspaper and radio, was circulated on July 21st. It stressed that the 1958 blight attack in Ireland was expected to be the worst for many years. By early August, this gloomy view was only too clearly justified, and poorly sprayed crops were destroyed in many areas by mid-August. In some parts of Ireland, the blight attack of 1958 was the worst during the present century.

8. THE FUTURE

The application of meteorology to the control of potato blight has made considerable progress in recent years. The synoptic approach, described above, has the merit of providing a possible means of considering simultaneously a whole group of plant diseases with similar general requirements for humidity, moisture and temperature. For example, Byrde (Reference 8) in a study of Brown Rot of Apple (*Sclerotinia fructigena*) has found that the weather conditions leading to outbreaks of sporulation in England were generally associated with either a slow-moving shallow low-pressure system or a moist south-westerly airstream of tropical or sub-tropical origin i.e. similar conditions to those favourable to the spread of potato-blight.

Progress in the case of other diseases is making rapid strides; a particularly promising feature is the application of upper air charts to problems of the dispersion of spores of cereal rusts and similar infections (Reference 7). In the case of soil borne diseases, the meteorologist is so far hampered by the limited observations available of soil temperatures and soil moisture.

The Commission for Agricultural Meteorology of the World Meteorological Organisation is alive to the importance of studies of meteorology as affecting the epidemiology of plant diseases. Apart from the report on potato blight already published (Reference 4), a Working Group to prepare a similar study on Apple Scab has recently been established.

In the general picture, the most important limiting factor is probably our still imperfect quantitative knowledge of the environmental factors of importance in the development of many plant diseases. As this is overcome, and with ever closer co-operation between plant pathologists and meteorologists there is a bright future for weather-disease studies, not only as applied to epidemiology, but also in the broader field of general plant pathology problems where meteorology can contribute to saving time and money by suggesting a rational approach in place of hit-or-miss empiricism.

R E F E R E N C E S

1. BOURKE, P.M.A. : Potato blight and the weather: a fresh approach - Irish Meteorological Service, Technical Note No. 12, 1953.
2. BOURKE, P.M.A. : The potato blight weather warning service in Ireland in 1952 - Irish Meteorological Service, Technical Note No. 13, 1953.
3. BOURKE, P.M.A. : Potato blight and the weather in Ireland in 1953 - Irish Meteorological Service, Technical Note No. 15, 1953.
4. BOURKE, P.M.A. : The forecasting from weather data of potato blight and other plant diseases and pests - World Meteorological Organisation : Technical Note No. 10, WMO-No. 42 TP. 16, 1955.
5. BOURKE, P.M.A. : Agricultural Meteorology - United Nations Technical Assistance Report No. TAA/CH1/2, 1956.
6. BOURKE, P.M.A. : The use of synoptic weather maps in potato blight epidemiology - Irish Meteorological Service, Technical Note No. 23, 1957.
7. BOURKE, P.M.A. : Meteorology and crop protection - WMO Bulletin, VII, 1, 14-7, 1958.
8. BYRDE, R.J.W. : Brown rot of apples - Transactions of the British Mycological Society, 39, 2, 280, 1956.
9. LARGE, E.C. : The interpretation of progress curves for potato blight and other plant diseases - Plant Pathology, 1, 4, 109-117.
10. VAMOS, R. : The connection between weather and the bruzone disease of rice - Időjárás, 58, 5, 273-7, 1954.
11. WADE, G.C. and WILLIAMS, P.G. : Late blight of potatoes : A progress Report of current research - Tasmanian Journal of Agriculture, 28, 11, 435-41.




FIG.1: RELATIONSHIP OF DIFFERENT CATEGORIES OF ADVICE AND WARNING ON THE DATE OF FIRST SPRAYING AGAINST POTATO BLIGHT.

P₁ = Period of blight-weather leading to first visible signs of disease.

P₂ = Subsequent period of blight-weather leading to intensification of disease.

X = Date of first visible signs of disease.

XYZ = Blight progress curve based on percentage of foliage attacked, as shown by vertical scale on the right.

A = Date of spraying advice based on first visible appearance of disease.

W = Date of spraying warning based on occurrence of significant period of blight-weather.

FW = Date of forecast spraying warning based on expectation of significant periods of blight-weather.




Fig. 2




Fig. 3




Fig. 4




Fig. 5

Section F : World literature

PHYTOLOGICAL BIOCLIMATOLOGY

Section F: World literature *

GENERAL PHYTOLOGICAL BIOCLIMATOLOGY

1. ALLARD, H.A. and EVANS, M.W.: Growth and flowering of some tame and wild grasses in response to different photoperiods - Jour. Agr. Res., 62, 193-228, 1941.
2. BALUT, S. : Der Einfluss der Tageslänge und der Temperatur auf den Verlauf des einjährigen Lebenszyklus der Samlinge FAGUS SILVATICA L. und ABIES ALBO Mill (Polish with German summary) - Ekol. Polska, Ser. A, 4, 225-292, 1956.
3. BLACK, J.N., BLACKMAN, G.E. and MARTIN, R.P.: Physiological and ecological studies in the analysis of plant environment. VIII. An inexpensive integrating recorder for the measurement of daylight - Ann. Bot. (Lond.) N.S., 17, 529-537, 1953.
4. BLACK, J.N. : Light intensity and plant growth - UNESCO, Arid Zone Research Proc. Canberra Symposium "Arid Zone Microclimatology" (in press).
5. BLACKMAN, G.E., BLACK, J.N. and KEMP, A.W.: Physiological and Ecological studies in the Analysis of Plant Environment X - Annals of Botany N.S., 19, 76, 1955.
6. BLACKMAN, G.E., and ROBERTSON-CUNINGHAME, R.C.: Interrelationships between light intensity and the physiological effects of 2:4-Dichlorophenoxyacetic Acid on the growth of Helianthus annuus - Journ. of Experimental Botany, 6, 17, 177-211, 1955.
7. BLACKMAN, G.E. : General Phytological Bioclimatology - See Areal report Dr. C.L. Cloudsley Thompson. Symposium Int. Soc. Bioclim. and Biomet. Paris 29- 31 August 1956.
8. BLEASDALE, J.K.A. : Smoke pollution and the growth of plants - Herbage Abstracts, 27, 3, 161-165, 1957.
9. BORTELS, H. : Azotobacter-Wachstum und Stickstoffbindung in Abhängigkeit vom Wetter - Zentralbl. f. Bakt. II, 94, 497-499, 1936.
10. BORTELS, H. : Entwicklung und Stickstoffbindung bestimmter Mikroorganismen in Abhängigkeit von Spurenelementen und vom Wetter - Ber. Dtsch. Bot. Ges., 56, 153-160, 1938.
11. BORTELS, H. : Ueber die Abhängigkeit bestimmter Bakterien von Spurenelementen und vom Wetter - Zentralbl. Bakt. 131, 494-496, 1938.
12. BORTELS, H. : Meteorobiologische Untersuchungen an Azobacter - Zentralbl. f. Bakt. II, 102, 129-153, 1940.
13. BORTELS, H. : Ueber die Abhängigkeit der Virulenz und anderer Eigenschaften pflanzenpathogener Bakterien sowie des Infektionserfolges vom Wetterverlauf - Phytopath. Zeitschr., 15, 376-393, 1948.
14. BORTELS, H. : Ueber mögliche Beziehungen zwischen Niederschlagsmenge und Wirkung von Molybdändüngungen auf Luzerne - Nachrichtenbl. BZA, 1, 140-142, 1949.
15. BORTELS, H. : Meteorobiologie vom Standpunkt des Mikrobiologen - Natur u. Volk, 82, 251-262, 1952.
16. BORTELS, H. : Das Gefrieren unterkühlten Wassers in Beziehung zu interdiurnen Luftdruckänderungen und zur Solaraktivität - Arch. f. Meteorolog. Geophys. u. Bioklim., 7, 269-276, 1956.
17. BORTELS, H. : Beziehungen zwischen Wetterwechsel, Atmung und Farbstoffbildung bei Mikroorganismen - Zentralbl. f. Bakt. 2, 109, 329-335, 1956.
18. BORTELS, H. : Ueber Beziehungen zwischen den Wirkungsgraden von Jute-Bandspinnmaschinen und der Farbstoffbildung durch Chromobacterium violaceum sowie interdiurnen Luftdruckänderung, der erdmagnetischen Aktivität und der Grenzfrequenz der sporadischen E. Schicht - Mellands Textilber. 1956.
19. BORTHWICK, H.A., HENDRICKS, S.B. and PARKER, M.W.: Photoperiodism - Radiation Biology 3, 479-517, 1956. (New York: McGraw-Hill Book Co., Inc.).

* References received from members by Secretariat of I.S.B.B. since 1957.

20. BOYKO, H. : Die Vegetationsverhältnisse im Seewinkel - Beihefte Bot. Zentralbl., 51, 600-747, 1934. (particularly p. 609-628). (In this paper Isoyetes are drawn on the basis of vegetation for the first time).
21. BOYKO, H. : Science and erosion problems in the middle East - J. of the Middle East Soc., 1/3, 1947.
22. BOYKO, H. : On the rôle of plants as quantitative climate indicators, and the Geo-ecological law of distribution - J. of Bot. Reh. Ser. J. of Ecol., 35/1, 138-157, 1947.
23. BOYKO, H. : On climatic extremes as decisive factors for plant distribution - Pal. J. of Bot. Reh. Ser., 7, 41-52, 1949.
24. BOYKO, H. : On regeneration problems of the vegetation in arid zones - Intern. Union of Biol. Sc. Ser. B., 9, 62-80, 1951.
25. BOYKO, H. : Climatic, Ecoclimatic and Hydrological influences on vegetation - Plant Ecology (Proc. of the Montpellier Symposium) Unesco series Arid Zone Research.
26. BOYKO, H. : Ecological Bioclimatology - See Areal report Prof. L. Emberger. Symposium Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956.
27. BROOKS, F.A. : Climatic environment: "A thermal system" - Syllabus, 1-153b, Aug. 1951
28. BURRICHTER, E. : Untersuchungen über die Massenentwicklung der Bodenbakterien im Laufe des Jahres - Ber. Dtsch. Botan. Ges., 71, 71-80, 1958.
29. CAILLEUX, A. : Richesse des flores, physiologie et évolution - Colloques Internat. du CNRS 41, 238-241, 1952.
30. CLAUSEN, J., KECK, D.D. and HIESEY, W.M.: Experimental studies on the nature of species. III. Environmental responses of climatic races of *Achillea* - Carnegie Inst. Wash. Publ., 581, 1948.
31. CLAUSEN, J. : Stages in the evolution of plant species - Ithaca, New York: Cornell University Press, 1951.
32. DAHL, E. : On the relation between summer temperature and the distribution of alpine vascular plants in the lowlands of Fennoscandia - *Oikos*, 1, 22-52, 1951.
33. DAIGO, Y. : A study on the Agroclimate of Japan - *Geophysical Mag.*, 28, 2, 149-236, 1957.
34. DUIN, R.H.A. van : De waterbehoefte der gewassen - *Landbouwkundig Tijdschrift*, 66, 4, 240-243, 1954.
35. DUIN, R.H.A. van : Influence of tilth on soil- and airtemperature - *Netherlands Journal of Agricultural Science*, 2, 4, 229-241, 1954.
36. DUIN, R.H.A. van and VRIES, D.A. de: A recording apparatus for measuring thermal conductivity, and some results obtained with in soil - *Neth. J. of Agr. Sc.*, 2, 3, 168-175, 1954.
37. DUIN, R.H.A. van : De betekenis van de grondbewerking voor vocht- en warmtetransport in de bodem - Presented at "Natuur- en Geneesk. Congres", Wageningen, 1955.
38. DUIN, R.H.A. van : Tillage in relation to rainfall intensity and infiltration capacity of soils - *Neth. J. of Agr. Sc.*, 3, 3, 182-191, 1955.
39. DUIN, R.H.A. van : On the influence of tillage on conduction of heat, diffusion of air and infiltration of water in soil - *Verslagen van Landbouwkundige Onderzoekingen*, Den Haag, 62, 7, 1956.
40. DUVDEVANI, S. : Effects of dew on plant growth under arid conditions - *American Jnl. of Botany*.
41. DUVDEVANI, S. : Dew in relation to plants in Palestine - *The Palestine Botanical Society Rehovoth*, 1939.
42. DUVDEVANI, S. and USHEROV: Analytical key for the identification of wild plants from their leaves, stems and roots - Tel Aviv, 1946 (book, 350 pp.)
43. DUVDEVANI, S. : An optical method of dew estimation - *Royal Meteorol. Soc.*, London 1947
44. DUVDEVANI, S. : Dew investigations under arid conditions - M.I.T. and Harvard Univ. Seminar, Cambridge, Mass., 1948.
45. DUVDEVANI, S. : Dew measurements - National meeting of American Meteorological Soc. New York, 1949.
46. DUVDEVANI, S. : Dew and its significance - Un. Nations Scientific Conference on the Conservation and Utilization of Resources, Lake Success, N.Y., 1949.
47. DUVDEVANI, S. : Dew gradients and the topography, climate and soil in arid areas - Int. Symposium on desert research, Jerusalem, 1952.
48. ERMICH, K. : Microclimatic investigations in plant communities of the Puszcza Niepolomicka - *Acta Societatis Botanicorum Poloniae*, 22, 483 - 559, 1953.
49. ERMICH, K. and ZURZYCKA, A.: The horizontal precipitation and plant water economy - *Wszeczeńświat*, 165-168, 1957.

50. FAIRBAIRN, W.A. : Ecological succession due to biotic factors, Northern Nigeria - Imperial Forestry Institute (Oxford), Paper no. 22, 1939.
51. GARNER, W.W. and ALLARD, H.A. : Further studies in photoperiodism, the response of the plant to relative length of day and night - Jour. Agr. Res., 23, 871-885, 1923.
52. GAUSSEN, H. : Végétation de la moitié orientale des Pyrénées - Climat, sol, végétation, Paris, 1926.
53. GAUSSEN, H. : Climat et végétation des Pyrénées - Toulouse, 1928.
54. GAUSSEN, H. : Le climat et la végétation de l'Ariège - Bordeaux, 1930.
55. GAUSSEN, H. : Sol, Climat, Végétation des Pyrénées Espagnoles - Zaragoza, 1935.
56. GAUSSEN, H. : Climat et flore dans le bassin de Paris - Paris, 1936.
57. GAUSSEN, H. : Projets pour diverses cartes du monde au 1.000.000^e. La carte écologique du tapis végétal - Congrès international de géographie de Lisbonne, 1949.
58. GAUSSEN, H. : A proposed ecological vegetation map - Surveying and Mapping, 13, 1953.
59. GAUSSEN, H. and BAGNOULS, F. : Période de sécheresse et végétation - Compte-rendu à l'Académie des Sciences, 1953.
60. GAUSSEN, H. : Le végétation et l'enneigement au Pyrénées - Paris, 1954.
61. GIACOBBE, A. : Schema di una teoria ecologica per la classificazione della vegetazione italiana - Nuovo Giornale Botanico Italiano, Firenze, 2, 1938.
62. GIACOBBE, A. : Le basi concrete per una classificazione ecologica della vegetazione Italiana - Archivio Botanico, Forlì, 1949.
63. GIACOBBE, A. : Ricerche ecologiche sull'aridità nei paesi del mediterraneo occidentale - da Webbia, 14, 1, 1958.
64. GODARD, M. and GESLIN, H. : Les gelées de décembre 1938 - C.R. Ac. Ag. de France, 122 - 130, 1939.
65. GODARD, M. and GESLIN, H. : L'actinométrie à distillation de Bellani et l'enregistrement de la radiation solaire - Le Météorologie, 151-156, 1940.
66. GODARD, M. : Le climat de la plante - Annales Agronomiques, 220-214, 1944.
67. GODARD, M. and HENIN, S. : Relation entre l'évaporation, la température et l'indice d'aridité - C.R. Ac. des Sc., 219, 559-560, 1944.
68. GODARD, M. : Radiation solaire et échauffement de la surface du sol - C.R. Ac. des Sc., 219, 624-626, 1944.
69. GODARD, M. : Observation sur le climat du sol - Ac. Agr. de France, 515, 1944.
70. GODARD, M. : Microclimats et Mésoclimats du point de vue agronomique - Annales Agronomiques, 578-604, 1949.
71. GODARD, M. : La transpiration végétale et le complexe chaleur-évaporation dans la France méditerranéenne - VIII^e Congrès International de Botanique, 1954.
72. GODARD, M. and NIGOND, J. : Les gelées de printemps à l'Ecole Nationale d'Agriculture de Montpellier - La Météorologie, 319-344, 1954.
73. HAARLØV, N. : Phytological Bioclimatology - See Areal report Prof. Dr. K. Knoch. Symposium Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956.
74. HICKS, W.W. : A series of experiments on trees and plants in electrostatic fields - Jnl. of the Franklin Inst., 264, 1, 1-5, 1957.
75. HUDSON, J.P. : General Phytological Bioclimatology - See Areal report Dr. J.L. Cloudsley-Thompson. Symposium Int. Soc. Bioclim. and Biomet., Paris 29 - 31 August 1956.
76. HUECK, K.C.A. : Urlandschaft, Raublandschaft und Kulturlandschaft in der Provinz Tucuman in N.W. Argentinien - Bonner Geogr. Abhandl. 10, Bonn, 1933.
77. HUECK, K.C.A. : Distribuicao de Pinheiro de Brasil e o problema dos campos no Brasil meridional - Ciencia e Cultura, Sao Paulo, 1952.
78. HUECK, K.C.A., RAWITSCHER and MORELLO : Algumas observacoes sobre a ecologia da vegetacao das Caatingas - Anais Academia Brasileira Ciencias, Rio, 1952.
79. HUECK, K.C.A. : Vegetationskarten aus Argentinien - Die Erde, Berlin, 1953.
80. HUSTICH, I. : On the correlation between growth and the recent climatic fluctuation - Geografiska Annaler, Stockholm, 1-20, 1949.
81. HUSTICH, I. : The recent climatic fluctuation in Finland and its consequences (Ed.) - Fennia, 75, 1-128, 1952.
82. JEFFREE, E.P. : Temperature ranges and distributions of some vascular plants - Nature, 176, 703-704, 1955.
83. JUNGES, W. : Die Wirkung von Tageslänge und Lichtintensität in der Präthermophase biennier Pflanzen - Ber. der Dt. Bot. Ges. 71, 5, 197-204, 1958.
84. KIEPENHEUER, K.O. : Ueber die Wirkung von Meterwellen auf das Teilungswachstum der Pflanzen - Die Naturwissenschaften, 36, 27, 1949.
85. KIRA, T. : Metabolic phytometer; a preliminary approach - Bull. Soc. Plant Ecol., 1, 126-129, 1951.

86. KNAPP, R. : Ueber den Einfluss der Höhenlagen und des Klimas auf die Artenzusammensetzung von Wiesen im mittleren Deutschland - Ltb. Samml., 6, 9-16, 1951.
87. KNAPP, R., LINSKENS, H.F., LIETH, H. and WOLF, F.: Untersuchungen über die Bodenfeuchtigkeit in verschiedenen Pflanzengesellschaften nach neueren Methoden - Ber. d. deutschen Botan. Ges., 65, 113-132, 1952.
88. KNAPP, R. : Studien zur Vegetation und Pflanzengeographischen Gliederung Nordwest-Italiens und der Süd-Schweiz - Kölner Geogr. Arbeiten, 4, 1-59, 1953.
89. KNAPP, R. and LINSKENS, H.F.: Beobachtungen über den Einfluss einiger mediterraner Pflanzengesellschaften auf Mikroklima und Bodenfeuchtigkeit - Angew. Botanik, 27, 48-69, 1953.
90. KNAPP, R. : Experimentelle Soziologie der höheren Pflanzen - Stuttgart-Ludwigsburg, 1954 (Book, 202 pp)
91. KNAPP, R. : Ueber den Verlauf der Keimung von einigen Gräsern, Leguminosen und Iberis amara bei unterschiedlichen Temperaturen - Angew. Botanik, 29, 125-138, 1955.
92. KNAPP, R. : Effekte wechselnder hoher und tiefer Temperaturen bei der Keimung von Sochus arvensis L. - Naturwissenschaften, 43, 41-42, 1956.
93. KNAPP, R. : Ueber den Einfluss der Temperatur auf die Entwicklung der nächsten Generation bei Agrostemma githago L. - Naturwissenschaften 43, 115, 1956.
94. KNAPP, R. : Untersuchungen über die Wirkungen täglicher Temperaturschwankungen auf Wachstum, Blütenentwicklung und Fertilität - Ber. d. Dtsch. Bot. Ges., 49, 399-412, 1956.
95. KNAPP, R. : Ueber die Wirkung von Giberellin auf Wachstum und Blütenbildung bei verschiedenen Temperatur- und Lichtverhältnissen - Zeitschr. f. Naturforschung, 11b, 698-704, 1956.
96. KRIVSKÝ, L. : Bestimmung der vorherrschenden Windrichtung aus Windrichtung aus Windfahnenbäumen - Met. Rundschau, 11, 3, 86-90, 1958.
97. LARSEN, E.C. : Photoperiodic responses of geographical strains of ANDROPOGON SCOPARIUS - Bot. Gaz., 109, 132-149, 1947.
98. LEMEE, G. : Recherches écologiques sur la végétation du Perche - Th.Sci., Paris, 1937 (388 pp).
99. LEMEE, G. : La méthode de l'analyse pollinique et ses apports à la connaissance des temps quaternaires - L'Année biol., 24, 2, 49-75, 1948.
100. LYSGAARD, L. : On the present climatic variation - Roy. Meteor. Soc. Cent.Proc., 206-211, 1950.
101. MÄDE, A. and ULLRICH, H.: Studien über die Ursachen der Frostresistenz. I. Untersuchungen des Temperaturtauses an Rizinusblättern - Planta, Arch. für Wiss. Botanik, Berlin, 28, 344-351, 1938.
102. MÄDE, A. : Das Fadenwiderstandsthermometer als Messgerät zur Bestimmung der Oberflächentemperaturen von Blättern - Biokl. Beibl. 6, 11-13, 1939.
103. MÄDE, A. : Ueber den Temperaturgang in Gewächshäusern, Dunkelkästen und Mistbeetanlagen (Ein Beitrag zum Mikroklima gärtnerischer Zuchtenlagen) - D. Gartenbauwissensch., 14, 626-641, 1940.
104. MÄDE, A. : Ueber den Temperaturverlauf in Beständen - D. Gartenbauwissensch. 15, 312-333, 1940.
105. MÄDE, A. und ULLRICH, H.: Studien über die Ursachen der Frostresistenz. II. Untersuchungen über den Temperaturverlauf beim Gefrieren von Blättern und Vergleichsobjekten - Planta, Archiv f.wiss.Botanik, Berlin, 31, 251-262, 1940.
106. MÄDE, A. und ULLRICH, H.: Temperaturuntersuchungen an erfrierenden Blättern - Umschau, 45, 43 - 45, 1941.
107. MÄDE, A. : Die Agrarmeteorologie in der Pflanzenzüchtung - Ein Bericht über Arbeiten der Agrarmeteorologischen Forschungsstelle Müncheberg/Mark des RfW - Wiss. Abh.d. RfW., 9, 6, 48, 1942.
108. MÄDE, A. : Agrarmeteorologische Forschung in Sachsen-Anhalt - Ztschr. f. Meteor., 2, 4/5, 115-116, 1948.
109. MÄDE, A. : Zum Mietenklima - Met. Rundschau, 3, 135-137, 1950.
110. MÄDE, A. : Meteorologie und Gartenbau - Dt. Agrartechnik, 1, 248-251, 1951.
111. MÄDE, A. : Temperaturbeobachtungen an vereinfachten Klimastationen - Angew. Met., 1, 53-56, 1951.
112. MÄDE, A. : Zur Methodik phänologischer Kartenwürfe - Angew. Met., 1, 139 - 142, 1952.
113. MÄDE, A. : Zur Methodik der Bodenfeuchtheitsmessungen - Ber. d. Dtsch. Wetterdienstes in der US-Zone, Bad Kissingen, 38, 1952.
114. MÄDE, A. : Bemerkungen zur Vereinfachung des phänologischen Meldeprogramms - Ber. Dt. Wetterd. US-Zone, Bad Kissingen, 42 (Knoch-Heft), 110-114, 1952.

115. MÄDE, A. : Agrarmeteorologie und Pflanzenschutz - Sonderdruck aus der "Pflanzenschutztagung"; Biologische Zentralanstalt Berlin DAL, Deutschen Bauernverlag, Berlin, 3, 7-12, 12-14, 1952.
116. MÄDE, A. : Ueber die Höhe des nächtlichen Taufalles im mitteldeutschen Trocken- gebiet - Wiss. Z. Univ. Halle, Math.-Nat., 4, 185-189, 1954.
117. MASSON, H. : L'eau de condensation et la végétation - Extrait du Bulletin de l' IFAN, 130-181, 1948. (2 clichés, 16 graphs).
118. MCCORMICK, J. : Phytological Bioclimatology - See Areal report Dr. D.H.K. Lee. Sym- posium Int. Soc. Bioclim. and Biomet. Paris 29-31 August 1956.
119. McMINN, R.G. : Role of soil drought in the distribution of vegetation in the nor- thern Rocky Mountains - Ecology 33.
120. MILLER, A.A. : Climatic requirements of some major vegetational formations - Ad- vancement of Science, 7, 25, 1950.
121. MILTHORPE, F.L. : Plant growth and the weather - Inaugural Lecture of the Univ. of Not- tingham, 1955.
122. MORELLO, J. : Wurzelsysteme der Vegetation von Julipao. Valle de Yocavil, Provinz Tucumén - Die Erde, 2, 195, 1950/51.
123. MORELLO, J. : Eficacia de método de pesadas rapidas no estudo da transpiracao de plantas. (en coautoría con F. Rawitscher) - Ciencia e Cultura, 1951.
124. MORELLO, J. : El Bosque de Algaerobo y la estepa de jarilla en el valle de Santa Mariá (Provincia de Tucumán) - Darwiniana, 9, 3, 4, 515-547, 1951.
125. MORELLO, J. : Eficacia des método de pesadas rápidas en el estudio de la transpira- ción de plantas (en coautoría con F. Rawitscher) : Ciencia e Inves- tigación, 8, 4, 183-185 and 6, 275-285, 1952.
126. MORELLO, J., RAWITSCHER, F., PAFFEN, K. and HUECK, K.: Algunas observacoes sobre a ecologia da vegetacao das Caatingas - An-acad. Bras. Cien., 24, 3, 287 - 301, 1952.
127. MORELLO, J. : Cuanta agua transpiran las plantas ? - Ciencia e Investigación, 9, 51-60, 1953.
128. MORELLO, J. : Transpiración y balance de agua de la bananera en las condiciones de la ciudad de Sao Paulo - Bol.Fac.Fil. Cien. Letr.Univ. Sao Paulo Bot., 10, 27-97, 1953.
129. MORELLO, J. : Ecología de una planta reviviscente de la Caatinga - Rev.Brasil.Bio- logía, 14, 1, 83-108, 1954.
130. MORELLO, J. : Estudios botánicos en las regiones áridas de la Argentina. III. Am- bientes, morfología y anatomía de 4 arbustos resinosos de follaje permanente - Rev. Agr. Noroeste, Arg. en prensa, 1955.
131. MORELLO, J. : Estudios botánicos en las regiones áridas de la Argentina. El Monte, características y límites - Lilloa en prensa, 1955.
132. MORELLO, J. : Regiones áridas de la Argentina. I Vegetación y monimientos del suelo en Neuquén extra andino - Lilloa en prensa, 1955.
133. MOSHKOV, B.S.: : Photoperiodismus und Frosthärte ausdauernder Gewächse - Planta, 23, 774-803, 1935.
134. NOVAK, V. : Begriff und Probleme des Bodenklimas (Czechosl. language) - Soil Re- search, 5, 1936.
135. NOVAK, V. : Taubildung im südmarischen Klima (Czechosl. language) - Annals of Cz. Academy of Agriculture, 1938.
136. NUTTONSON, M.Y. : Some preliminary observations of phenological data as a tool in the study of photoperiodic and thermal requirements of various plant ma- terial - Vernalization and Photoperiodism. A Symposium, 1, 1948.
137. OLMSTED, C.E. : Growth and development in range grasses. IV. Photoperiodic response in twelve strains of side-oats grama - Bot. Gaz., 106, 46-74, 1944.
138. OLSON, J.S. and NIENSTAEDT, H.: Photoperiod and chilling control growth of hemlock - Science, 125, 492-494, 1957.
139. OVINGTON, J.D. : General Phytological Bioclimatology - See Areal report Dr. J.L. Clouds- ley Thompson. Symposium Int. Soc. Bioclim. and Biomet. Paris, 29 - 31 August 1956.
140. PEREIRA, H.C. and HURSH, C.R.: Field moisture Balance in the Shimba Hills, Kenya - E. Afr. Agric. J., 18, 139-145, 1953.
141. PEREIRA, H.C. : Moisture measurements in East African soils - Proc. 2nd. Inter- Afr. Soils Conference, Leopoldville, 425-430, 1954.
142. PEREIRA, H.C. : Soil water storage under catchment area vegetation - Proc. 5th Int. Soil Sci. Congress, Leopoldville, 1954.
143. PIERCE, L.T. : Estimating Seasonal and Short-Term fluctuations in Evapo- Transpira- tion from Meadow Crops - Bull. of the Am.Met.Soc., 39, 2, 73-78, 1958.
144. PLATT, R.B. : An ecological study of the mid-Appalachian shale barrens and of the plants endemic to them - Ecological Monographs, 21, 269-300, 1951.

145. PLATT, R.B. : Broader horizons for the biologist in microclimatic research - Jnl. of the Tennessee Academy of Science, 27, 199, 1952.
146. PLATT, R.B. and AMSLER, F.P.: Growth rate determinations for tree lichens - Jl. of the Tennessee Academy of Science, 27, 203, 1952.
147. PLATT, R.B. and WOLF, J.N.: A method for controlling natural soil temperature gradients - Ecology, 33, 2, 1952.
148. PLATT, R.B. : Research status and needs for the biologist in microenvironment research - Bull. of the Ecol. Soc. of America, 34, 75, 1953.
149. PLATT, R.B. : Autenvironment and plant ecology - an assessment - Rapports et Communications, VIII Congrès International de Botanique, Sect. 7, 225-227, 1954.
150. PLATT, R.B. : Microenvironmental research at the Emory University Field Station - Bull. of the Georgia Academy of Science, 13, 1-6, 1955.
151. PLATT, R.B. and AMSLER, F.P.: A basic method for the immediate study of lichen growth rates and succession - Jl. of the Tennessee Academy of Science, 30, 177-183, 1955.
152. PRESCOTT, J.A. : The soils of Australia in relation to vegetation and climate - C.S. I.R. Bull., 52, 82, 1931.
153. PRESCOTT, J.A. : Evaporation from a water surface in relation to solar radiation - Trans. Roy. Soc. of S. Aust., 64, 114-118, 1940.
154. PRESCOTT, J.A. : A climatic index for the leaching factor in soil formation - Journal Soil.Sci., 1, 9-19, 1949.
155. PRESCOTT, J.A. and TOMAS, J.A.: The length of the growing season in Australia as determined by the effectiveness of the rainfall. A revision. - Proc. Roy. Geogr. Soc. S. Aust., 50, 42-46, 1949.
156. PRESCOTT, J.A. and COLLINS, J.A.: The lag of temperature behind solar radiation - Quart. Jl. Roy. Met. Soc., 57, 121-126, 1951.
157. PRESCOTT, J.A. and THOMAS, J.A.: Climatic expressions and generalised climatic zones in relation to soils and vegetation - Specialist Conf. in Agric. Australia 1949, Proc. 27-33, 1951.
158. PRESCOTT, J.A., BONYTHON, C.W. and COLLINS, J.A.: Evaporation in Australia in January and July - Trans. Roy. Soc. of S. Aust., 1954.
159. PRIMAULT, B. : Die Auswirkung von Gewitterniederschlägen auf die Temperatur des Erdbodens - Der Landfreund, 1951. (24 August)
160. PRIMAULT, B. : Quelques considérations sur le gel en Valais et les moyens disponibles pour en diminuer les effets - Le Confédéré, Martigny, 1951 (23/4)
161. PRIMAULT, B. : Le gel et les moyens d'en diminuer les dégâts - Le Sillon romand, 1951 (27/4).
162. PRIMAULT, B. : Der Frost und die heutigen Mittel um seine Schaden zu bekämpfen - Der Landfreund, 1951 (11/5).
163. PRIMAULT, B. : Les températures dans le sol, à Zürich, au cours de l'année 1952 - Annalen der Schweizerischen Meteorologischen Zentralanstalt, 1 -12, 1953.
164. PRIMAULT, B. : Leçon à tirer d'une nuit de gel - Revue romande d'Agriculture, de Viticulture et d'Arboriculture, 9ème année, 10, 1953.
165. PRIMAULT, B. : La température d'un pré - Annalen der Schw. Meteorologischen Zentralanstalt, 1954.
166. PRIMAULT, B. : De l'évapotranspiration en Suisse romande - Le Sillon Romand. 59ème année, 29 and 30, 1956 (20 et 27/7).
167. PRIMAULT, B. : Phytological Bioclimatology - See Areal report Prof. Dr. K. Knoch. Symp. Int. Soc. Bioclim. and Biomet. Paris 29-31 August 1956.
168. PUJOS, A. : Sur un sous-étage de végétation méditerranéen aride froid à Chénopodiacees et la présence de l'étage méditerranéen saharien dans le Bassin de la Moulouya (Maroc) - Comptes rendus des séances de l'Académie des Sciences, 240, 1010-1012, 1955.
169. RAMDAS, L.A. : Meteorology of air layers near the ground. I. - Technical Bull. nr. 3. of Ind. Met. Dept.
170. RAMDAS, L.A. : Meteorology of air layers near the ground. II. - Technical Bull. nr. 9. of Ind. Met. Dept.
171. RAMDAS, L.A. : The climates of the air layers near the ground at Poona. III. Technical Note nr. 21, Ind. Met. Dept.
172. RAMDAS, L.A. and ATMANATHAN, S.: The vertical distribution of air temperature near the ground during night - Gerlands Beitrage Zur Geophysik, 37, 116, 1932.
173. RAMDAS, L.A. : Agricultural Meteorology - Cur. Sci., 1, 7, 191, 1933.
174. RAMDAS, L.A. : Microclimatology - Cur. Sci., 2, 11, 445, 1934.
175. RAMDAS, L.A. : Annual report of the Agricultural Meteorology Section, India Meteorological Dept., for the year ending 21st August 1933 - Special publications of the Indian Council of Agr. Research.

176. RAMDAS, L.A. and DRAVID, R.K.: Soil temperatures - Current Science, 3, 6, 226, 1934.
177. RAMDAS, L.A., KALAMKAR, R.J. and GADRE, K.M.: Agricultural Meteorology: Studies in Micro-climatology. I. - Ind. Jour. of Agric. Sci., 4, 3, 1934.
178. RAMDAS, L.A. and KATTI, M.S.: The variation of moisture in the surface layer of the soil in relation to the diurnal variation of meteorological factors - Current Science, 3, 1, 24, 1934.
179. RAMDAS, L.A. and KATTI, M.S.: Agricultural Meteorology: Preliminary studies on soil moisture in the surface layer of the atmosphere during the clear season at Poona - Ind. Jour. of Agric. Sci., 4, 6, 1934.
180. RAMDAS, L.A. : Measurement of evaporation with special reference to (a) water supply and (b) Agriculture - Report of the Conference of Empire Meteorologists, London, 180, 1935.
181. RAMDAS, L.A. : Correlation of general crop observations with meteorological data - Report of the Conference of Empire Meteorologists, London, 173, 1935.
182. RAMDAS, L.A. : Note on soil temperature - Report of the Conference of Empire Meteorologists, London, 160, 1935.
183. RAMDAS, L.A. : Report on the progress of Agricultural Meteorology in India during the period 1932-34 - Kommission für Landwirtschaftliche Meteorologie, Protokolle der Tagung in Danzig, 1935 (no. 24).
184. RAMDAS, L.A., KALAMKAR, R.J. and GADRE, K.M.: Agricultural Meteorology: Studies in micro-climatology, II. - Ind. Jour. of Agric. Sci., 5, 1, 1935.
185. RAMDAS, L.A. : Frost hazard in India - Current Science, 3, 3, 325, 1935.
186. RAMDAS, L.A. and RAMANATHAN, K.R.: Derivation of Angstrom's formula for atmospheric radiation and some general considerations regarding nocturnal cooling of air layers near the ground - Proc. Ind. Acad. of Sci., 1, 11, 1935.
187. RAMDAS, L.A. and KATTI, M.S.: The diurnal variation of moisture in the soil during the clear season - Current Science, 3, 12, 1935.
188. RAMDAS, L.A. and DRAVID, R.K.: Soil temperature in relation to other factors controlling the disposal of solar radiation at the earth's surface - Proc. of the Nat. Inst. of Sci., 3, 3, 1936.
189. RAMDAS, L.A. and KATTI, M.S.: Agricultural Meteorology: Studies on soil moisture in relation to moisture in the surface layers of the atmosphere during the clear season at Poona - Ind. Jour. of Agr. Sci., 6, 6, 1936.
190. RAMDAS, L.A. : Report on simple methods of measurement in Agricultural Meteorology - Sent to the Intern. Commission on Agr. Met., Sept. 1937.
191. RAMDAS, L.A. : Report on visible condensation - Sent to the Intern. Commission on Agr. Met., Sept. 1937.
192. RAMDAS, L.A. and NARASIMHAN, M.: Agricultural Meteorology: The prediction of the minimum temperature on clear day at selected stations in India - Ind. Jour. of Agric. Sci., 7, 5, 1937.
193. RAMDAS, L.A., SREENIVASIAH, B.N. and RAMAN, P.K.: Variation in the nocturnal radiation from sky with zenith distance and with time during the night - Proc. Ind. Acad. of Sci., 5, 1, 1937.
194. RAMDAS, L.A. : The variation with height of the water vapour content of the air layers near the ground at Poona - Bioklimatische Beiblätter, I, 1938.
195. RAMDAS, L.A. and MALLIK, A.K.: The exchange of moisture between air layers near the ground and substances like soils, plant leaves, seeds, etc. exposed to the atmosphere - Current Science, 6, 9, 1938.
196. RAMDAS, L.A. and MALLIK, A.K.: Loss of water by evaporation from the upper surface of soil columns resting on a water table - Current Science, 8, 6, 1939.
197. RAMDAS, L.A. and MALLIK, A.K.: The thickness of the surface layer of the soil exchanging moisture with the adjacent air layers during the clear season at Poona - Current Science, 8, 4, 1939.
198. RAMDAS, L.A., RAMAN, P.K. and SREENIVASIAH, B.N.: Variation in the nocturnal radiation from the sky with zenith distance and with time during the night - Proc. of Ind. Acad. Sci., 9, 5, 1939.
199. RAMDAS, L.A. : Cold waves and frosts - Indian Farming, 1, 11, 1940.
200. RAMDAS, L.A. and JOGLEKAR, S.Y.: Proc. of Ind. Acad. of Sci., 13, 1941.
201. RAMDAS, L.A. : The micro-climates of plant communities - The Indian Ecologist, 1, 1, 1946.
202. RAMDAS, L.A. : Rainfall of India. A brief review - Empire Jl. of Exp. Agric., 14, 54, 1946.
203. RAMDAS, L.A. and RAMAN, P.K.: A method of estimation on the thickness of the "laminar" layer above an evaporating water surface - Proc. of Ind. Acad. Science, 23, 1946.
204. RAMDAS, L.A. and MALLIK, A.K.: The effect of sodium chloride in improving the permeability of alkali soils - Current Science, June 1947.

205. RAMDAS, L.A. : Soil Moisture and Evaporation investigations - Central Board of Irr. Jour., 5, 3, 1948.
206. RAMDAS, L.A. : Report on evaporation from bare soil in relation to the depth of the water table or the zone of saturation - Communicated to the meeting of the Intern. Union of Geodesy & Geophys., Oslo, August 1948.
207. RAMDAS, L.A. : The physics of the bottom layers of the atmosphere - Presidential Address to the Physics section of the Indian Science Congress Patna 1948.
208. RAMDAS, L.A. : Effect of salts on soil permeability and rectification of alkali ridden soils - Jour. of C.S.I.R., 3, 1, 1949.
209. RAMDAS, L.A. : Desert Hydrology: Condensation and evaporation phenomena in the Rajasthan Desert - Bull. of the Nat. Inst. of Sci. of India, 1, 1952.
210. RAMDAS, L.A. : Convective phenomena near a heated surface - Proc. of Ind. Acad. of Sci., 37, 2, 1953.
211. RAPPE, G. : Den växtlösliga näringshaltens i jorden växlingar med årstiderna - Nordisk Jordbruksforskning, 3, 4, 1933. (German summary)
212. RAPPE, G. : Våra stråsådessorter deras avkastning och utbredning jämte en inledande översikt över Stråsådesförädlingen - Lund, Gleerupska Univ. Bokhandeln, 1933.
213. RAPPE, G. : Prövning av en ny metod för utförande av enkla fältförsök - Vänersborg, C.W. Carlsson Eftr:s Boktryckeri, 1934.
214. RAPPE, G. and JOHNSON, H.: Qualitätsvergleiche zwischen Mineralboden- und Torfbodenheu in Süd- und Nordschweden (Swedish) - Svenska Mosskulturföreningens Tidskrift, 52, 2, 1938.
215. RAPPE, G. : Skördetidens och salpetersgödslingens inflytande på ängsgrödhöets kvalitet - Svenska Vall- och mosskulturföreningens meddelanden, 1, 1939.
216. RAPPE, G. : Hökvalitet, skördetid och kvävegödsling - Svenska Vall- och Mosskulturföreningens Kvartalskrift, 1, 1939.
217. RAPPE, G. : Något om betes- och vallproduktionens anpassning till föreliggande marktillgångar - Svenska Vall- och Mosskulturföreningens kvartalskrift, 3, 1942.
218. RAPPE, G. : Jahresbericht über die Stüdienhofs-Tätigkeit der "Svenska Vall- och Mosskulturföreningen" - Norrtälje, Norrtälje Tidnings Boktryckeri A. B., 1944.
219. RAPPE, G. : Die Jahreszeitsvariationen der Egner'schen Kali- und Phosphorsäurezahlen für besäten und unbesäten Mineralboden, Niedermoorboden, und Hochmoorboden (Swedish) - Svenska Vall- och Mosskulturföreningens Tidskrift, 4, 1944.
220. RAPPE, G. : Seasonal variations observed in the nutritive substances of a cultivated sphagnum peat soil - Transactions of the International Congress of Soil Science, Amsterdam, 1, 1950.
221. RAPPE, G. : Seasonal variations in some properties of Arable soil - Plant and Soil, 4, 3, 1953.
222. RAPPE, G. : Observations on Season, Grassgrowth and Soil Conditions - Societas pro Studio- Rhythmi Biologici. Fifth Conference 1955. Resumé.
223. ROUSSEAU, J. : Notes sur les mouvements périodiques de quelques plantes - Nat.can., 57, 3, 1, 109-112, 1930, and Devoir, 25, 127, 10, 1934.
224. ROUSSEAU, J. : Le climat, la plante et la civilisation - La Revue Populaire, 30, 6-7, 1937.
225. ROUSSEAU, J. and ROUSSEAU, B.: Etude comparative de la température de l'estuaire du Saint-Laurent et des habitants voisins - Nat.can., 64, 127-131, 1937, and Contrib. Lab. Bot. Univ. Montreal, 29, 65-69, 1937.
226. ROUSSEAU, J. and MARIE-VICTORIN, F.: Nouvelles entités de la flore phanérogamique du Canada oriental - Contrib. Inst. bot. Univ. Montreal, 36, 74, 1940.
227. ROUSSEAU, J. : The vegetation and life zones of George River, Eastern Ungava, and the welfare of the natives - Arctic, 1, 2, 93-96, 1948.
228. ROUSSEAU, J. : Modification de la surface de la toundra sous l'action d'agents climatiques - Revue can. de géographie, 3, 1/4, 43-51, 1949.
229. ROUSSEAU, J. : Les zones biologiques de la péninsule Québec-Labrador et l'hémiarctique - Can Journ. of Botany, 30, 4, 436-474. 1952 and Mémoire du Jardin botanique de Montreal no. 27.
230. ROUSSEAU, J. and RAYMOND, M.: Le Gentiana tenella Rottb. dans le Québec arctique - Le Naturaliste canadien, 79, 2, 76-80, 1952.
231. SAUBERER, F. : Ueber die Strahlungseigenschaften der Pflanzen im Infrarot (Vorläufige Mitteilung) - Wetter und Leben, 1, 231, 1949.
232. SAUBERER, F. : Niederschlagsmessungen am Hordhang des Kleinen Hetzkogels - Wetter und Leben, 1, 297, 1949.

233. SAVAGE, D.A. : Drought survival of native grass species in the central and southern Great Plains - U.S. Dept. Agr. Tech. Bull., 549, 1937.
234. SMATOK, I.D. : The effect of day length under polar conditions on the carbohydrate and mineral content of *CARAGANA ARBORESCENS* and *SYRINGA VULGARIS* - Akad. Nauk. S.S.R.R. Dok., 111, 213-216, 1956.
235. SMILEY, T.L. : Geochronology as an aid to research in the study of arid lands - The Future of Arid Lands, edited by Gilbert F. White. Proc. of the Intern. Arid Lands Symp. and Conf., April-May 1955. A.A.A.S. Symposium Publication.
236. SMOLIK, L. : To the evaporation in the air near the soil - Sbornik CAZ, 1938.
237. SMOLIK, L. : The second maximum of the air temperature near the soil - Sbornik, CAZ, 1939.
238. SMOLIK, L. : Influence of the wind break upon the evaporation - Vestnik CAZ, 1946.
239. SMOLIK, L. : The depth of the freezing of our soils 1939/40 - Vestnik CAZ, 1946.
240. SMOLIK, L. : Humidity of the air as a source of soil moisture - N.O. Zem.vyd, 1946.
241. SMOLIK, L. : Condensation of the air water vapour and soil moisture - Scientific conference of the Polytechnical University, 1957.
242. SOLOMON, M.E. : Control of humidity with potassium hydroxide, sulphuric acid or other solutions - Bull.ent.Res., 42, 3, 543-554, 1951.
243. STANHILL, G. : Evaporation of water from soil under field conditions - Nature, 176, 82, 1955.
244. STEPHENS, C.G. : The soils of Tasmania - C.S.I.R. Bull., 139, 1941.
245. STRUZKA, V. and UHLÍR : Measuring in Wind Screens in Bulgaria (Czechosl.) - Meteorologické zprávy, 3, 1952.
246. SUBRASMANIAN, V.K. and MISRA, R.K. : Sub-soil irrigation in relation to plant growth, II. - Journal of the Central Board of Irrigation, 5, 4, 1948.
247. TAKASU, K. : Microclimatic Study.V.: Leaf temperatures under natural environments - Mem. Coll.Sci., Kyoto Univ., Ser. B., 20, 3, 8, 179-187, 1953.
248. TIMMERS, H.J., THIJSSSEN, H.A.C., WIT, C.T. de, VOLLENHOVEN, E.v. and ADMIRAAL, L.: New instruments for agricultural research - Neth.J. Agr. Sci., 2, 209-214, 1954.
249. TIMMERS, H.J. : Determination of soil permeability in situ - Neth. J. of Agr. Sci., 3, 2, 119-126, 1955.
250. TRANQUILLINI, W. : Der Ultrarotabsorptionsschreiber im Dienste ökologischer Messungen des pflanzlichen CO_2 - Umsatzes - Ber. dtsh.bot. Ges., 65, 102, 1952.
251. TRANQUILLINI, W. : Ueber den Einfluss von Uebertemperaturen der Blätter bei Dauereinschluss in Küvetten auf die ökologische CO_2 - Assimilationsmessung - Berichten der deutschen bot. Gesellschaft, 67, 191, 1954.
252. TRUMBLE, H.C. : The relation of pasture development to environmental factors in South Australia - Jour. Agric. of S. Aust., 38, 1460-1487, 1935.
253. TRUMBLE, H.C. and CORNISH, E.A.: The influence of rainfall on the yield of a natural pasture - C.S.I.R. Journal, 9, 19-28, 1936.
254. TRUMBLE, H.C. : Some factors affecting the germination and growth of herbage plants in South Australia - Jour. Agric. of S. Aust., 40, 779-786, 1937.
255. TRUMBLE, H.C. : Agricultural Climatology in Australia - Journal Aust.Inst. of Agric. Sci., 2, 3, 115-119, 1945.
256. TRUMBLE, H.C. : Rainfall, evaporation and drought frequency in South Australia - J. Dept. Agr.Sth.Aust., 52, 55-64, 1948.
257. TRUMBLE, H.C. : Climatic and soil factors affecting future land use in South Australia - Proc. Roy. Geogr. Soc. Sth.Aust., 49, 18-32, 1948.
258. TRUMBLE, H.C. : The ecological relations of pastures in South Australia - J. British Grassland Soc., 4, 135-160, 1949.
259. TRUMBLE, H.C. : Grassland agronomy in Australia - Advances in Agronomy, 4, 1-65, 1952.
260. TURESSON, G. : The selective effect of climate upon the plant species - Hereditas, 14, 99-152, 1930.
261. TURNER, H. : Ueber das Licht- und Strahlungsklima einer Hanglage der Oetztaaler Alpen bei Obergurgl und seine Auswirkung auf das Mikroklima und auf die Vegetation - Arch.Met.Geoph. Biokl., B, 8, 3/4, 273-325, 1958.
262. TURRELL, F.M. : A list of early flowering herbaceous plants near Sorento, Illinois - Private publication, 1932.
263. TURRELL, F.M. : The relation between chlorophyll concentration and the internal surface of mesomorphic and xeromorphic leaves grown under artificial light - Proc. Iowa Acad. Sci., 46, 107-117, 1939.
264. TURRELL, F.M. and KERSTEN, H.: A qualitative study of the effect of soft X-rays on the activity of B-indolyl acetic acid as a growth substance - Trans. Iowa Acad. Sci., 49, 139-143, 1942.
265. TURRELL, F.M. and TURRELL, M.: The effect of the great drought of 1934 on the leaf structure of certain Iowa plants - Proc. Iowa Acad. Sci., Leaf Structure, 50, 185-194, 1943.

266. TURRELL, F.M. : Correlation between internal surface and transpiration rate in mesomorphic and xeromorphic leaves grown under artificial light - The Botanical Gazette, 105, 413-425, 1944.
267. TURRELL, F.M. : Maybe this is the cause (Scientist investigates effect of sulphur gases in industrial smoke on vegetation - Citrus Leaves, 25, 11, 6 - 7, 1945.
268. TURRELL, F.M. : Effect of sulphur gases in industrial smoke on vegetation - Calif. Citrograph, 31, 2, 40-41, 1945.
269. TURRELL, F.M. : The reaction of sulphur dust with plant constituents - Amer. Soc. Plant Physiol. Abstract and Titles of Papers, 20th Annual Meeting, 3, 1946.
270. TURRELL, F.M. : Tables of corresponding pH and (H⁺) Values - Univ. of California Citrus Experiment Station Riverside, California, 1946.
271. UNGER, H. : Bodenfeuchtigkeit und Asternwelke - Angew. Met., 1, 9, 278-280, 1953.
272. UNGER, K. : Zur Abschätzung von Verdunstungsunterschieden verschiedener Pflanzenbestände am natürlichen Standort - Angew. Met., 2, 1, 1-14, 1954.
273. VAARTAJA, O. : General Phytological Bioclimatology - See Areal report Dr. W.G. Wellington. Symposium Int. Soc. Bioclim. and Biomet., Paris 29 - 31 August 1956.
274. VEGIS, A. : Ueber den Einfluss der Temperatur und der täglichen Licht-Dunkel-Periode auf die Bildung der Ruheknochen; zugleich ein Beitrag zur Entstehung des Ruhezustandes - Symbolae Bot. Upsaliensis, 14, 1 - 175, 1955.
275. VRIES, D.A. de : Natuurkundige beschouwingen omtrent de diffusieweerstand en het warmtegeleidingsvermogen van gronden - Landbouwkundig Tijdschrift, 65, 11, 676-683, 1953.
276. VRIES, D.A. de and DUIN, R.H.A. van: Some considerations on the diurnal variation of transpiration - Neth. J. Agr. Sci., 1, 27-34, 1953.
277. VRIES, D.A. de and VENEMA, H.J.: Some physical considerations on the behaviour of the Piche evaporimeter - Vegetatio, Acta geobotanica, 5/6, 1, 8, 225-234, 1954.
278. VRIES, D.A. de and WIT, C.T. de: Die thermische Eigenschaften der Moorböden und die Beeinflussung der Nachtfrostgefahr dieser Böden durch eine Sanddecke - Meteorologische Rundschau, 7, 3/4, 41-45, 1954.
279. VRIES, D.A. de : On the evaluation of the eddy diffusivity in the lower air layers from the diurnal temperature wave - Quarterly J. of Roy. Met. Soc., 1955.
280. VRIES, D.A. de : Solar radiation at Wageningen - Mededelingen L.H.S. (Commun. Agric. Univ., Wageningen) 55, 1956.
281. VRIES, D.A. de : On the evaluation of the thermal (or eddy) diffusivity from the diurnal temperature wave - Netherlands Journal of Agr. Sci., 4, 1, 136-139, 1956.
282. WATERHOUSE, F.L. : Microclimatological profiles in grass cover in relation to biological problems - Quarterly J. of the Royal Met. Soc., 81, 347, 1955.
283. WEAVER, J.E. and ALBERTSON, F.W.: Major changes in grassland as a result of continued drought - Bot. Gaz., 100, 576-591, 1939.
284. WENT, F.W. : The response of plants to climate - Science, 112, 1950.
285. WENT, F.W. : Phytological Bioclimatology - See Areal report Dr. D.H.K. Lee. Symposium Int. Soc. Bioclim. and Biomet., Paris 29 - 31 August 1956.
286. WILLIAMS, C.B. : A short bioclimatic study in the Egyptian Desert - Min. Agr. Egypt. Tech. & Scientif. Service Bull., 29, 1-20, 1923.
287. WILLIAMS, C.B. : Bioclimatic observations in the Egyptian Desert in March 1923 - Min. Agr. Egypt. Tech. & Scien. Service Bull., 37, 1-18, 1924.
288. WILLIAMS, C.B. and MCKENZIE-TAYLOR, E.: A comparison of sand and soil temperatures in Egypt - Min. Agr. Egypt. Tech. & Scient. Service Bull., 40, 1-24, 1924.
289. WILLIAMS, C.B. : A third bioclimatic study in the Egyptian Desert - Min. Agr. Egypt. Tech. & Scient. Serv. Bull., 50, 1-32, 1925.
290. WILLIAMS, C.B. : Some bioclimatic observations in the Egyptian Desert. From " Biology of Deserts " - Proc. of Symposium on Biology of hot and cold deserts in 1952, Inst. Biology London, 18-27, 1954.
291. WIJK, W.R. van, VRIES, D.A. de and DUIN, R.H.A. van: Potential evapotranspiration - Neth. J. Agr. Sci., 1, 35-39, 1953.
292. WIJK, W.R. van : Laboratorium für Physik und Wetterkunde der Landwirtschaftlichen Hochschule, Wageningen, Niederlande - Achema-Jahrbuch, Sect. II, 125-126, 1953/55.
293. WIJK, W.R. van : Het klimaat nabij de grond - Elseviers Weekblad, 20 May 1954.
294. WIJK, W.R. van and VRIES, D.A. de: Evapotranspiration - Neth. J. Agr. Sci., 2, 105-119, 1954.
295. WIJK, W.R. van and WESSELING, J.: Optimal depth of drainage - Neth. J. of Agr. Science, 3, 2, 106-118, 1955.

296. WIJK, W.R. van and HIDDING, A.P.: Onderzoek over de verandering van het klimaat achter windschermen - Landb. Tijdschrift 67, 707-712, 1955.
297. WYNNE-EDWARDS, V.C. : Isolated arctic-alpine floras in eastern North America. A discussion of their glacial and recent history - Tr. Roy.Soc. Canada, 5, 3, 31, 1-26, Walker prize (Boston Soc. Nat. Hist.), 1937.
298. WYNNE-EDWARDS, V.C. : Some factors in the isolation of rare alpine plants - Trans. Roy.Soc. Can., 5, 3, 33, 35-42, 1939. (pl. I-III).

GENERAL AGRICULTURAL BIOCLIMATOLOGY

1. BEALL, J.M. : A proposal for a revitalized agricultural meteorological service - AMS Bulletin, 35, 9, 1954.
2. BEALL, J.M. : New horizons in agricultural meteorology - Florida State Horticultural Society, 1954.
3. BEALL, J.M. : How greater utilization of weather information can pay off to farmers and canners - Information Letter, National Canners Association, 1526, 1955.
4. BEALL, J.M. : Crop Dusting - Weekly Weather and Crop Bulletin, 43, 4, 1956.
5. BEALL, J.M. : Weather and agricultural research - Chemurgic Digest, 15, 2, 1956.
6. BEALL, J.M. : Weather service for the farmer. A review and forecast - Presented at Fifty-Second Meeting Washington State Horticultural Assoc., Yakima, Washington, 1956.
7. BEALL, J.M. : Weather and the production of Burley tobacco - AMS Bulletin, 38, 2, 1957.
8. BIDER, M. : Phänologische Beobachtungen in den Kantonen Baselland, Basel-Stadt, Uri und Graubünden - Tätigkeitsbericht der Nat. Ges. Baselland, 57-90, 1936/38.
9. BIDER, M. : Korrelationen zwischen Witterung und Ernteerträgen - Verhandl. der Schweiz. Naturforsch. Ges., 135, 1945.
10. BIDER, M. and MEYER, A.: Lässt sich der Zeitpunkt der Kirschenernte in der Nordwestschweiz vorausbestimmen? Versuch einer frühzeitigen Prognose des Kirschen-ertrages - Schw. Zeitschr. Obst.u. Weinbau, Wädenswil, 1946.
11. BIDER, M. : Aufblühdaten eines Kirschbaumes während 6 Dezennien. Quotienten der Niederschlagsmenge: Sättigungsdefizit. Mit A. Meijer: 10 Jahre Kirschen-ernte Prognose in der Nordwestschweiz - Wiss. Ges. für Landw. und Forstkunde, Freiburg i/Breisgau. Intern. Kolloquium, 1953.
12. BLACK, J.N. : The influence of varying light intensity on plant growth with particular reference to graminaceous and leguminous species - Thesis, Univ. of Oxford, 1952.
13. BLACK, J.N. : The interaction of light and temperature in determining the growth rate of subterranean clover - Aust. Jl. Biol.Sci., 8, 330-343, 1955.
14. BLACK, J.N., BLACKMAN, G.E. and KEMP, A.W.: An analysis of the effects of seasonal variation in daylight and temperature on the growth of *HELIANTHUS ANNUUS* in the vegetative phase - Ann. Bot. (Lond.) N.S., 19, 527-548, 1955.
15. BLACK, J.N. : The influence of varying light intensity on the growth of herbage plants; a review - Herbage Abstracts, June 1957.
16. BORTELS, H. : Fortgeführte Untersuchungen über die Beziehung zwischen Infektionserfolg und Wetterverlauf bei der Kartoffelnassfäule - Phytopatholog. Zeitschr., 18, 360-362, 1951.
17. BORTELS, H. : Ueber Beziehungen zwischen der durch *Pseudomonas tabaci* (Wolf und Foster) Stevens hervorgerufenen "Wildfeuer"-Erkrankung des Tabaks, Luftdruckänderungen und Solaraktivität - Phytopathologische Zschr., 33, 4, 403-425, 1958.
18. BOURKE, P.M.A. : Potato blight and the weather: a fresh approach - Technical Note no. 12 of the Irish Meteorological Service, 1953.
19. BOURKE, P.M.A. : The potato blight weather warning service in Ireland in 1952 - Technical Note no. 13 of the I.M.S., 1953.
20. BOURKE, P.M.A. : Potato blight and the weather in Ireland in 1953 - Technical Note No. 15 of the I.M.S., 1953.
21. BOURKE, P.M.A. : The forecasting from weather data of potato blight and other plant diseases and pests - World Met. Organization: Technical Note no. 10-WMO No. 42 TP 16, 1955
22. BOURKE, P.M.A. : Agricultural Meteorology - United Nations Technical Assistance Programme Report No. TAA/CHI 2, 1956.

23. BOURKE, P.M.A. : The use of synoptic weather maps in potato blight epidemiology - Technical Note no. 23 of the I.M.S., 1957.
24. BROOKS, F.A., SCHOONOVER, W.B. and WALKER, H.B.: Protection of Orchards against frost - Agric. Ext. Service Univ. of California, Circular 111, 1939.
25. BROOKS, F.A. and Deering, R.B.: Landscaping for Summer Shade - Cal. Agric., 7, 5, 11-16, 1953.
26. BROOKS, F.A., RHOADES, D.G. and LEONARD, A.S.: Wind machines tests in Citrus - Cal. Agric. 8, 8, 8-10, 1954.
27. BROOKS, F.A., RHOADES, D.G., LEONARD, A.S. and SCHULTZ, H.B.: Frost protection in Almonds - Cal. Agric., 9, 8, 3-5, 14, 1955.
28. COLLINGWOOD, C.A. : Fruit tree spraying trials in the West Midlands - Herefords. Agric. Journ., 12, 275-277, 1951.
29. COLLINGWOOD, C.A. : Pests of soft fruit - Worcs. Agric. Chron., 21, 243-245, 1953.
30. DUIN, R.H.A. van and SCHOLTE UBING, D.W.: De invloed van het weer op de opbrengst van de aardappel - Landbk. Tijdschrift, 67, 795-802, 1955.
31. DUVDEVANI, S, REICHERT and PALTÍ: The development of downy and powdery mildews in cucumbers as related to environmental factors - Palestine Journ. of Bot., 1946.
32. DUVDEVANI, S. : Agrobotanical atlas for students - Pardess Hanna agric. School, 1949.
33. GAUSSEN, H. : Le climat de la région de Toulouse et son influence sur l'Agriculture - Toulouse, 1945.
34. GAUSSEN, H. : L'Olivier et le climat insubrien - Paris, 1951.
35. GODARD, M. : Influence des facteurs climatiques sur la croissance de la betterave sucrière - Annales Agronomiques, Paris, 697-727, 1937.
36. GODARD, M., BALACHOWSKY, A. and GESLIN, H.: Recherches sur l'utilisation du froid dans la lutte contre le ver des cerises (*Rhagoletis cerasi* L.) - Annales des Epiphyties, 137-140, 1937.
37. GODARD, M. : La température du sol et le développement automnal de la betterave sucrière - An/Ag. de France, 357-364, 1938.
38. GODARD, M. : Le climat solaire et la croissance de la betterave sucrière - C.R. Ac. des Sc., Paris, 208, 929, 1939.
39. GODARD, M. and LIMACET, P.: Nouvelles recherches sur le *Phytophthora infestans* (Mont.) de Bary - Annales des Epiphyties, 145-156, 1940.
40. GODARD, M. : La maturation de l'oeillette et la radiation solaire - Ac. Agr. de France, 47, 1945.
41. GODARD, M. : Facteurs physiques du milieu et transpiration du Maïs: Action de la température - C.R. Ac. des Sc., Paris, 229, 850-852, 1949.
42. GODARD, M. : Facteurs physiques et transpiration du Maïs. Action du pouvoir évaporant de l'air: Influence du vent - C.R. Ac. des Sc., Paris, 229, 900-901, 1949.
43. GODARD, M. : Action de la température sur la transpiration du Maïs: influence varietale - C.R. Ac. des Sc., Paris, 238, 2343-2345, 1954.
44. GOODALL, D.W. : Seasonal and diurnal changes in the water content of the tomato seedling - Rep. exp. Res. Sta. Cheshunt., 92-96, 1936.
45. GOODALL, D.W. : Some preliminary observations on the position of the first inflorescence in the tomato plant - Rep. exp. Res. Sta. Cheshunt., 87 - 92, 1936.
46. GOODALL, D.W. : Further observations on factors affecting the position of the first inflorescence in the tomato - Rep. exp. Res. Sta. Cheshunt., 73-78, 1937.
47. GOODALL, D.W. : The distribution of weight change in the young tomato plant. I. Dry-weight changes of the various organs - Ann. Bot. London, N.S. 9, 101-139, 1945.
48. GOODALL, D.W. : The distribution of weight change in the young tomato plant. II. Changes in dry weight of separated organs and translocation rates - Ann. Bot., London, N.S. 10, 305-338, 1946.
49. GOODALL, D.W. : Growth of cocoa seedlings as affected by illumination - Paper read to International Horticultural Congress, Scheveningen, Neth., 1955.
50. GRAINGER, J. : Potato blight forecasting and its mechanization - Nature, 171, 1012, 1953.
51. GRAINGER, J. : Climate, host and parasite in crop disease - Quarterly Jl. of the Roy. Meteorological Soc., 81, 347, 1955.
52. GRAINGER, J. : The "auchincruive" Potato blight forecast recorder - Weather, 10, 7, 1955.
53. GRAINGER, J., SNEDDON, J.L., CHISHOLM, C. and HASTIE, A.: Climate and the yield of cereal crops - Quarterly Jl. of the Royal Meteorological Soc., 81, 347, 1955.
54. HESSE, W. : Agricultural Bioclimatology - Areal report: Symposium Int. Soc. Bioclim. and Biomet., Paris, 29-31 August 1956.
55. HUSTICH, I. : Yields of cereals in Finland and the recent climatic fluctuation - Fennia, 73, 3, 1-48, 1950.

56. KNOCH, K. : Die Vorhersage der Reisernte in Nordjapan - Die Naturwissenschaften, 993-995, 1922.
57. KNOCH, K. : Die Möglichkeiten der Abschätzung des Ernteertrages auf Grund meteorologischer Angaben. Ein wichtiges Kapitel aus dem Gebiete der landwirtschaftlichen Meteorologie - Die Naturwissenschaften, 769-776, 1923.
58. KNOCH, K., GEIGER, R. und HOLDEFLEISS, P.: Bericht über die seit 1928 in Deutschland auf dem Gebiete der landwirtschaftlichen Meteorologie durchgeführten Untersuchungen - Commission de Météorologie Agricole. Proces-Verbaux des Séances de Munich 1932, Utrecht, 40-57, 1933.
59. KNOCH, K., GEIGER, R. und HOLDEFLEISS, P.: Bericht über die seit 1932 in Deutschland auf dem Gebiete der landwirtschaftlichen Meteorologie durchgeführten Untersuchungen - Secrétariat de l'Organisation Met. Intern. Nr. 24 Kommission für Landwirtschaftliche Met., Danzig, 133-156, 1935.
60. KNOCH, K. : Hagelverbreitung und Hagelschäden in Deutschland - Organisation Météorologique Intern., Leyden, 36, 67-70, 1938.
61. KNOCH, K. : Bericht über agrarmeteorologische Forschungen des Reichsamtes für Wetterdienst und seiner Aussenstellen - Organisation Météorologique Intern., Leyden, 36, 72-77, 1938.
62. KNOCH, K. : Angewandte Klimatologie als Förderer wirtschaftlicher Probleme - Zeitschrift für angew. Met., 55, 39-48, 1938.
63. KOSHIMIZU, T. : Carpobiological studies of *Crinum asiaticum* var. *japonicum* Bak. - Mem. col. Sci. Kyoto Imp. Univ. B, 5, 2, 1930.
64. KOSHIMIZU, T. : Phytogeographical distribution of *Crinum* throughout the World - Bot. Mag., 52, 613, 1938.
65. KOSHIMIZU, T. : On the *Crinum* line in the flora of Japan - Bot. Mag., 52, 615, 1938.
66. KOSHIMIZU, T. : The giant *Crinum* growing in Java is nothing but a ecotype of *Crinum asiaticum* var. *japonicum* - Jour. Japanese Bot., 27, 6, 1952.
67. LAKE, J.V. : The effect of microclimatic temperature on the date of flowering of the apple - J. hort. Sci., 31, 244-257, 1956.
68. LANDSBERG, H.E. and MÜLLER, H.: Frostbekämpfungsversuche in Rheingau - Wein und Rebe, 1935.
69. LANDSBERG, H.E. : Agricultural Bioclimatology - See Areal report Dr. D.H.K. Lee, Symposium Int. Soc. Bioclim. and Biomet. Paris, 29-31 August 1956.
70. LEE, D.H.K. : Factors affecting the vitamin C content of fruit and vegetables - Med. J. Aust., 1, 786, 1940.
71. LORCH, J. : Analysis of Windbreak Effects - The Bull. of the Research Council of Israel, 6D, 4, 211-220, 1958.
72. MADE, A. : Ein Beitrag zum Mikroklima eines Obstbaumes (Mikroklimatische und blütenphänologische Beobachtungen) - Die Gartenbauwissenschaft, 12, 127-137, 1938.
73. MADE, A. und RUDORF, W.: Zweck und Aufbau moderner bewetterter Gewächshäuser und ihr Temperaturgang im Vergleich mit dem des Freilandes - Biokl. Beibl. 5, 145.153, 1938.
74. MADE, A. : Temperaturuntersuchungen an Obstbäumen (Zur Frage der Entstehung von Frostschäden an Obstbäumen im Klagenfurter Becken) - Gerlands Beitr. z. Geophysik, 59, 201-213, 1942.
75. MADE, A. : Methodische Voruntersuchungen für eine Agrarklimatologie der DDR - Habilitationsschrift 1957. Manuscript.
76. MORRIS, L.L. : Field and transit chilling of fall-grown tomatoes - Proc. Conf. on Transportation of Perishables. Univ. of Cal. Davis, April 1954.
77. MÜLLER, K. : Die Entwicklungsgeschwindigkeit des Winterroggens zwischen Blüte und Vollreife in Abhängigkeit von den Witterungselementen - Diss. Köln, 1950 (55 pp and figures).
78. NÄGELI, W. : Ueber die Bedeutung von Windschutzstreifen zum Schutze landwirtschaftlicher Kulturen - Schweizerische Zeitschrift für Forstwesen, 11, 1941.
79. NISHIUCHI, H. : The accumulative temperature in frostless season as an index of temperature climate of warm season at the stand of crops (In Japanese) - Proc. of the Crop Sci. Soc. of Japan, 15, 179-184, 1944.
80. NISHIUCHI, H. : On the index of temperature climate at the limit of water rice crop culture in the world - Journ. of Agric. Met., 8, 14-16, 1953.
81. NISHIUCHI, H. : Relation between the isothermal lines of the mean air temperatures of the coldest and warmest months and the limit of the distributions of crops in the world - Jl. of Agric. Met., 10, 61-62, 1954.
82. NOVAK, V. : Ein Beitrag zur Lösung des Gartenmikroklimas (Czechosl. language) - Annals of Cz. Academy of Agriculture, 1938.
83. NUTTONSON, M.Y. : Ecological Plant Geography of Albania, its agricultural crops and some North American Climatic Analogues - American Institute of Crop Ecology, Washington, D.C.

84. NUTTONSON, M.Y. : Agricultural Climatology of Poland and its agro-climatic analogues in North America - Am. Inst. of Crop Ecology, Washington, D.C.
85. NUTTONSON, M.Y. : Agricultural Climatology of Yugoslavia and its agro-climatic analogues in North America - Am. Inst. of Crop Ecology, Washington, D.C.
86. NUTTONSON, M.Y. : Agricultural Climatology of Czechoslovakia and its agro-climatic analogues in North America - Am. Inst. of Crop Ecology, Washington, D.C.
87. NUTTONSON, M.Y. : Ecological crop geography of the Ukraine and the ukrainian agro-climatic analogues in North America - Intern. Agro-climatological series, Study no. 1, 1947.
88. NUTTONSON, M.Y. : Ecological crop geography of China and its agro-climatic analogues in North America - Am. Inst. of crop Ecology. Int. agroclimatological series, Study no. 7, 1947.
89. NUTTONSON, M.Y. : Ecological crop geography of Greece and its agro-climatic analogues in North America - Int. agro-climatological series, Study no.5, 1947.
90. NUTTONSON, M.Y. : Agroclimatology and crop ecology of Palestine and Transjordan and climatic analogues in the United States - The Geographical Review, 37, 3, 436-456, 1947.
91. NUTTONSON, M.Y. : Ecological crop geography of Germany and its agroclimatic analogues in North America - Am. Inst. of Crop Ecology, Washington, D.C., 1949.
92. NUTTONSON, M.Y. : Ecological crop geography of Finland and its agroclimatic analogues in North America - Int. Agro-climatological series, Study no.10, 1950.
93. NUTTONSON, M.Y. : Agricultural climatology of Sweden and its agro-climatic analogues in North America - Am. Inst. of crop-ecology, Int. agroclimatological series. Study no. 11, 1950.
94. NUTTONSON, M.Y. : Ecological crop geography of Norway and its agro-climatic analogues in North America - Int. agro-climatological series. Study no.12, 1950.
95. NUTTONSON, M.Y. : Agricultural climatology of Siberia, its natural Belts and agro-climatic analogues in North America - Am. Inst. of Crop Ecology, 1950.
96. NUTTONSON, M.Y. : Agricultural Climatology of Japan and its agro-climatic analogues in North America - Am. Inst. of Crop Ecology, Washington, D.C.
97. NUTTONSON, M.Y. : Ecological crop geography and field practices of Japan, Japan's natural vegetation, and agro-climatic analogues in North America - Am. Inst. of Crop Ecology, Washington, D.C., 1951.
98. NUTTONSON, M.Y. : Ecological crop geography and field practices of the Ryukyu islands, natural vegetation of the Ryukyu's and agro-climatic analogues in the Northern Hemisphere - Am. Inst. of Crop Ecology, Washington D.C., 1952.
99. NUTTONSON, M.Y. : Phenology and thermal environment as a means for a physiological classification of wheat varieties and for predicting maturity dates of wheat - Am. Inst. of Crop Ecology, Washington D.C., 1953.
100. NUTTONSON, M.Y. : Wheat-climate relationships and the use of Phenology in ascertaining the thermal and photo-thermal requirements of wheat - Am. Inst. of Crop Ecology. A research organisation devoted to problems of plant adaption and introduction, 1955.
101. PASCALE, A.J. : Comportamiento bioclimático de tres variedades de trigo en un ensayo de siembras continuadas - Tesis. Fac. de Agr.y.Vet., Bs.Aires, 1950.
102. PASCALE, A.J. : Reacción de los trigos argentinos a la vernalización - Servicio Meteorológico Nacional, Bs. Aires, 1951 (not published)
103. PASCALE, A.J. : Comportamiento fotoperiódico de algunos trigos argentinos - METEOROS 33, 1, 1953.
104. PASCALE, A.J. : El Índice Heliotérmico aplicado a los trigos argentinos (en colaboración) - METEOROS, 4, 3, 1954.
105. PASCALE, A.J. : Método para determinar las características bioclimáticas de una variedad de trigo - METEOROS, 5, 1-2, 1955.
106. PASCALE, A.J. : Comportamiento bioclimático de una colección de ciruelos (en colaboración) - Servicio Meteorológico Nacional, Bs. Aires, 1955 (not published).
107. PRESCOTT, J.A. : The nitrogen problem in the Australian wheat belt - Pan-Pacific Sci. Congress (Fifth), Vancouver 1933.
108. PRESCOTT, J.A., TRUMBLE, H.C. and DAVIDSON, J.: Climatology in relation to biology and agriculture - Journ. Aust.Inst. Agric. Sci., 3, 77-79, 1937.
109. PRESCOTT, J.A. : Indices in agricultural climatology - Journ.Aust.Inst.Agric. 4, 33-40, 1938.
110. PRESCOTT, J.A. : The climate of Tropical Australia in relation to possible agricultural occupation - Trans. Roy. Soc. of S.Aust., 62, 229-240, 1938.
111. PRESCOTT, J.A. : The response of the growth of pastures to temperature - Aust.Jl.Sci., 11, 24-25, 1948.
112. PRESCOTT, J.A. and BUTLER, P.F.: Evapotranspiration from wheat and pasture in relation to available moisture - Aust.Jr.Agr.Sci., 6, 52-61, 1955.

113. PRIMAULT, B. : Chronique météorologique. Suite d'aperçus de l'évolution du temps dans le Canton de Vaud et de ses répercussions sur les cultures - La Terre Vaudoise, Lausanne.
114. PRIMAULT, B. : Les coiffes anti-gel et leurs répercussions sur le développement des ceps - Geofisica pura e applicata, Milan, 20, 1951.
115. PRIMAULT, B. : Temperaturuntersuchungen in der Kambiumzone von Obstbäumen auf der Südwest- und Nordostseite - Wissenschaft. Ges. für Land- und Forstwirtschaft, Freiburg i.B., 1953.
116. PRIMAULT, B. : Die Verwendbarkeit von Strohschirmen als Frostschutz für Reben im Lichte der Maifröste 1953 - Schw. Zschr. für Obst- und Weinbau, Nr.8, 1954.
117. PRIMAULT, B. : L'influence de l'insolation sur la température du cambium des arbres fruitiers - Rev. Romande d'Agricult., de Viticult. et d'Arboricult., 10, 3, 1954.
118. PRIMAULT, B. : Die agrarmeteorologische Dreiländertagung am 25.Mai in Hallau - Landwirtschaftliche Monatshefte, July 1955.
119. PRIMAULT, B. : Arboriculteurs, méfiez-vous du soleil du printemps - La Terre vaudoise, 26. May 1955.
120. PRIMAULT, B. : Un moyen pour améliorer et régulariser la production des vergers - La Terre vaudoise, 7 May 1955.
121. PRIMAULT, B. : Utilisation de données climatologiques en agriculture - Conf. diffusée par Sottens le 15 juillet 1956 dans le cadre de "Problèmes de la vie rurale".
122. RAINY, R.C. : Climatic characteristics of the South African cotton-growing areas - Reprints from Experiment Station, London 1948-'49, 26-30, 1950.
123. RAMDAS, L.A. : Crop Protection from the effects of weather - Proc. of the crop and soil wing of the Board of Agr. and Animal Husbandry, Dec. 1937.
124. RAMDAS, L.A. and KALEMKAR, R.J. : Statistical investigations on crop-weather relationships in India - Sankhya, May 1938.
125. RAMDAS, L.A. : Annual Report of the Agricultural Meteorology Section, India Meteorological Dept., for the year 1941-42 - Special publication of the Indian Council of Agricultural Research (Roneoed).
126. RAMDAS, L.A. : Climatic factors in Agriculture - Ind. Farming, 3, 5, 1942.
127. RAMDAS, L.A. : The effect of climatic factors on plant growth, crop yield and the incidence of pests and diseases - Proc. of the 5th meeting of the Crops and Soils Wing of the Board of Agriculture and Animal Husbandry, 1942.
128. RAMDAS, L.A. and MALLIK, A.K. : Studies on soils. Part I: The upward movement of water and salt solutions in black cotton soil - Proc. of Ind. Acad. of Sci., 16, 1, 1942.
129. RAMDAS, L.A. and MALLIK, A.K. : Studies on soils. Part II: A microscopic study of the behaviour of the black cotton soil in salt solutions - Proc. of Ind. Acad. of Sci., 16, 1, 1942.
130. RAMDAS, L.A., MALLIK, A.K. and PANDIT, U.P. : The effect of concentration on the capillary movement of some salt solutions through the black cotton soil - Current Science, July 1942.
131. RAMDAS, L.A. : Annual report of the Agricultural Meteorolog. Section, Indian Meteorological Dept. for the year 1942-43 - Special Publications of the Indian Council of Agr. Research (Roneoed).
132. RAMDAS, L.A. : Annual report of the agricultural Meteorological Section, Indian Meteorological Dept., for the year 1943-44 - Special Publications of the Indian Council of Agricultural Research (Roneoed).
133. RAMDAS, L.A. and MALLIK, A.K. : On the effect of concentration of sodium carbonate in aqueous solutions on (A) the formation of cracks (B) swelling and dispersion and (C) Capillary ascent in the black cotton soil - Current Science, February 1944.
134. RAMDAS, L.A. : Annual report of the Agricultural Meteorological Sect., Indian Meteorological Dept., for the year 1944-45 - Special publications of the Indian Council of Agricultural Research (Roneoed).
135. RAMDAS, L.A. : Annual report of the Agricultural Meteorological Sect., Indian Meteorological Dept., for the year 1945-46 - Special publications of the Indian Council of Agricultural Research (Roneoed).
136. RAMDAS, L.A. : Report on recent progress in Agricultural Meteorology in India (with five appendices) - Communicated to the International Commission on Agr.-Met. held at Toronto, August 1947.
137. RAMDAS, L.A. : Report on "The Coordinated Crop weather Scheme-India" - Communicated to the International Commission on Agr. Met. held at Toronto, August 1947.

138. RAMDAS, L.A. and MALLIK, A.K.: Phenology of the Wheat Crop in India - Communicated to the International Commission on Agr. Met., held at Toronto, August 1947.
139. RAMDAS, L.A. and MALLIK, A.K.: Studies on soils. Part III: On the effect of concentration of some salt in aqueous solution on capillary ascent, dispersion and associated phenomena in the black cotton soil - Proc. Ind. Acad. Sci., 26, 1947.
140. RAMDAS, L.A. and MALLIK, A.K.: Weather Service for agriculture: Crop-weather calendars, Farmer's Weather Bulletins, warnings for adverse weather (with three appendices) - Communicated to the International Commission on Agr. Met. held at Toronto, August 1947.
141. RAMDAS, L.A. : Some new instruments and experimental techniques developed in the agricultural Meteorology Section at Poona - Jour. of Sci. & Industrial Research, 7, 16-29, 1948.
142. RAMDAS, L.A. and MALLIK, A.K.: The objective assessment of the crop outlook week by week on the basis of the progress of weather week by week during the growing season - Agricultural Situation in India, September 1948.
143. RAMDAS, L.A. : Rainfall and Agriculture - Ind. Jour. of Met. & Geophysics, 1, 4, 1950.
144. RAPPE, G. : Effect of manuring, especially with nitrogenous fertilizers, on the productive capacity of potatoes for setting (in Swedish) - Svenska Mosskulturföreningens Tidskrift, Årg. 50, 1, 1936.
145. RAPPE, G. : Growth variations on pasture land during the different parts of the season. Some results of 6 years' cutting tests (in Swedish) - Svenska Vall- och Mosskulturföreningens Kvartalskrift, Årg. 6, 1, 1944.
146. RAPPE, G. : Some observations concerning pasture growth in relation to water supply (in Swedish) - Svenska Vall- och Mosskulturföreningens Kvartalskrift, Årg. 8, 4, 1946.
147. RAPPE, G. : Graphical representation of pasture growth and weather (in Swedish) - Svenska Vall- och Mosskulturföreningens Meddelanden, 13, 1946.
148. RAPPE, G. : Suitable use of different soils as a means for levelling the unevenness of pasture supply (in Swedish) - Svenska Vall- och Mosskulturföreningens Kvartalskrift, Årg. 9, 1, 1947.
149. RAPPE, G. : Some growth features in pastures (in Swedish) - Svenska Vall- och Mosskulturföreningens Meddelanden, 14, 1948.
150. RAPPE, G. : Seasonal variations in the rate of pasture regrowth after grazing - Plant and Soil, 3, 4, 1951.
151. RAPPE, G. : Nitrification in eight farm soils - Oikos, Acta oecologica Scandinavica, 4, 1, 1952-1953.
152. ROBERTSON, G.W. : Probability table of spring frost at Central Experimental Farm - Ottawa, 1952.
153. ROBERTSON, G.W. : Probability table of fall frost at Central Experimental Farm - Ottawa 1952.
154. ROBERTSON, G.W. : Some agro-meteorological problems in Canada - Royal Met. Soc. Canadian Branch, 4, 2, 1-18, 1953.
155. ROBERTSON, G.W. : Evaporation measurements at branch units of the Experimental Farms Service throughout Canada - Bull. of Division of Field Husbandry, Soils and Agr. Engineering Dept. of Agriculture, Canada, 1-19, 1953.
156. ROBERTSON, G.W. : Agro-meteorology - Cereal News, Central Experimental Farm, Ottawa, 1, 9, 8-12, 1953.
157. ROBERTSON, G.W. : Latent Evaporation: Its concepts, measurement and application - Bull. of Division of Field Husbandry, Soils and Agricultural Eng. Dept. of Agriculture, Canada, 1-21, 1954.
158. ROBERTSON, G.W. : Agro meteorology. A National Crop-Weather Program - Presented at a conference to discuss the status of investigational work at the Experimental Substations. Fort Simpson, N/W/T/ Whitehorse, Y. and T. & Fort Vermilion, Alberta, Ottawa, 13 January, 1955.
159. ROBERTSON, G.W. : The standardization of the measurement of evaporation as a climatic factor - Technical note no 11, World Met. Org., Geneva, nr 42, 16, 1-10, 1955.
160. ROBERTSON, G.W. : The measurement of latent evaporation using the Central Experimental Farm plastic mount - Division of Field Husbandry, Soils and Agr. Eng.; Division of Illustration Stations, Centr. Exp. Farm, Ottawa, 1 - 10, 1956.
161. ROBERTSON, G.W. : Procedure for keeping a meteorological soil moisture budget for irrigation purposes - Division of Field Husbandry, Soils and Agricultural Engineering, Central Experimental Farm, Ottawa, 1-4, 1956.
162. ROBERTSON, G.W. : The use of climatic data in irrigation planning - Proc. of the Second Annual Meeting of the Can. Soc. of Agronomy, Univ. of Toronto, 31-38 1956.

163. ROBERTSON, G.W. : Estimating irrigation water requirements from meteorological data - Proc. of the First Ontario Irrigation Conf., Ontario, Agricultural College, Guelph, 1-7, 1956.
164. ROBERTSON, G.W. : Agricultural Bioclimatology - Areal Report: Symposium Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956.
165. ROBERTSON, G.W. : Agricultural Meteorology. A brief literature review and bibliography - Experimental Farms Service, Ottawa, 1-59, 1956 (co-author).
166. ROBERTSON, G.W. and HOLMES, R.M.: Estimating irrigation water requirements from meteorological data - Experimental Farms Service, Ottawa, 1-19, 1956.
167. ROGERS, W.S. : Rootstock effect on colour and size of apples - E-Malling Res. Sta. Ann. Rept., 16-32, 1926 (May 1927).
168. ROGERS, W.S. : A soil moisture meter depending on the "Capillary Pull" of the soil, with illustrations of its use in fallow land, grass orchards and irrigated orchards - J. Agric. Sci., 25, 326-43, 1935.
169. ROGERS, W.S. : Root Studies. VI. Apple roots under irrigated conditions with notes on use of a soil moisture meter - Journal of Pomology and Horticultural Science, 13, 3, 1935.
170. ROGERS, W.S. : Soil factors in relation to root growth - Int Congr. Soil.Sci., 3, 1, 249-253, 1935.
171. ROGERS, W.S. : The relation of soil moisture to plant growth, illustrated by moisture meter experiments with strawberries - E.Malling, Res.Sta. Ann. Rept. for 1935, 111-120, May 1936.
172. ROGERS, W.S. : Soil moisture studies. I. Water utilisation by apple trees in cultivated and grass orchard compared with fallow land - E. Malling. Res. Sta. Ann. Rept. 1936, 105-109, May 1937.
173. ROGERS, W.S. and SRIVASTAVA, D.N.: Soil moisture studies: II. Moisture variation in a pear orchard - E. Malling Res. Sta. Ann. Rept. 1936, 110-113, May 1937.
174. ROGERS, W.S. : An apple irrigation trial, using soil moisture meters - E. Malling Res. Sta. Ann. Rept. 1938, 1939.
175. ROGERS, W.S. : Root studies VIII: Apple root growth in relation to rootstock, soil, seasonal and climatic factors - J. Pomol. and Hort. Sci., 17, 1939.
176. ROGERS, W.S. : Root studies IX: the effect of light on growing apple roots; a trial with root observation boxes - J. Pomol. and Hort. Sci., 17, 1939.
177. ROGERS, W.S. : Making the best use of water in food production - J. Roy. Hort. Soc., 66, 279-281, 1941 and E. Malling Res. Sta. Ann. Rept., 1940, 82-83, 1941.
178. ROGERS, W.S. and MOORE, M.H.: Sun scald of fruits - Gdnrs.Chron. 122, 66-67 and reprinted with addendum E. Malling Res. Sta. Ann. Rept. 1942, 50-53, 1943.
179. ROGERS, W.S. : Bark splitting of fruit trees - Gdnrs.Chron., 113-187, 1943.
180. ROGERS, W.S. : Strawberries, straw and frost - Fruitgrower, 95, 393-394, 1943.
181. ROGERS, W.S. : Frost damage to fruit: a note on the present position of research in England - Annual report of the East Malling Research Station for 1948 1949.
182. ROGERS, W.S. and MODLIBOWSKA, I.: Low temperature injury to fruit blossom. III. Water sprinkling as an anti-frost measure - Rep. E. Malling Res. Sta. for 1949, 63-68, 1950.
183. ROGERS, W.S. and BEAKBANE, A.B.: The effect of netting and hessian anti-frost screens and of brick walls on air temperature - Rep. E. Malling Res. Sta. for 1950, 60-64, 1951.
184. ROGERS, W.S. : Protection from spring and autumn frosts by continuous water sprinkling - Grower, 36, 13, 626-628, 1951 and Rep. E. Malling Res. Sta. for 1951, 197-198, 1951.
185. ROGERS, W.S. : Fruit plant roots and their environment - Rep. 13th Int. Hort. Cong. 1952.
186. ROGERS, W.S. : Some aspects of spring frost damage to fruit trees and its control - Rep. 13th. Int. Hort. Cong., 1952.
187. ROGERS, W.S. and GOODE, J.E.: Irrigation requirements of fruit orchards - Rep. E. Malling Res. Sta. for 1952, 171-173, 1953.
188. ROGERS, W.S., MODLIBOWSKA, I., RUXTON, J.P. and SLATER, C.H.W.: Low temperature injury to fruit blossom. IV. Further experiments on water-sprinkling as an anti-frost measure - J. Hort. Sci., 29, 2, 126-141, 1954.
189. SMITH, L. : Relation of Polyploidy to heat and X-ray effects in the cereals - Journal of Heredity, Washington, D.C., 34, 5, 130-134, 1943.
190. SMITH, L. : A comparison of the effects of heat and X-rays on dormant seeds of cereals, with special reference to polyploidy - Journal of Agricultural research, Washington D.C., 73, 4, 137-158, 1946.
191. SMOLIK, L. : Influence of the direction of the potatoes row upon the air temperature - Sbornik, CAZ, 1941.

192. SMOLIK, L. : Influence of the direction of the potatoes row upon the soil temperature - Sbornik CAZ, 17, 276, 1942.
193. STRUZKA, V. : Improvement of the climate for agricultural purposes (in Czechoslov.) - Vesmír, 1, 1953.
194. STRUZKA, V. : The wind (Czech.) - A chapter - 22 pages - of the book: Klika, Novak, Gregor: Praktikum fytocenologie, ekologie, klimatologie a pudoznalství, Praha, 1954.
195. SUBRAMANIAN, V.K. and MALLIK, V.K.: Flowering of wheat in relation to air temperature - Ind. Journal of Met. & Geophysics.
196. SUBRAMANIAN, V.K. : Drought resistance of cotton - Ind. Cotton Growing Rev., 7, 4, 1953.
197. SUBRAMANIAN, V.K., MALLIK, A.K. and VENKATARAMAN: Complete enumeration experiment in Groundnut Crop for developing the optimum shape and size of samples for crop-weather observations - Proc. Ind. Acad. Sci., 40, 1954.
198. TAKASU, : Microclimatic study in tea plantation at the time of spring frost (in Japanese) - 1954.
199. TRUMBLE, H.C. : The climatic control of agriculture in South Australia - Trans. Roy. Soc. of S. Aust., 61, 41-62, 1937.
200. TRUMBLE, H.C. : Climatic factors in relation to the agricultural regions of southern Australia - Trans. Roy. Soc. of S. Aust., 63, 36-43, 1939.
201. TURRELL, F.M. and KLOTZ, L.J.: Rind structure and composition in water spot of navel orange - California Citrograph, 25, 2, 45, 56, 57, 1939.
202. TURRELL, F.M. and CUENO, F.: The production of H_2S gas by sulphur dusted citrus fruits - Bot. Soc. Amer. Abstract, 1940.
203. TURRELL, F.M. : Radiation and sulphur injury to lemon fruits - Amer. Jour. Bot., 31, 5-6, 1944.
204. TURRELL, F.M. : Estimating heights of Citrus trees - Proc. of the Am. Soc. for Hort. Science, 48, 1946.
205. TURRELL, F.M. : Chemical and physical factors in the sulfur burn in citrus - Scalacs 3, 9, 285, 1948.
206. TURRELL, F.M. : Specific gravity of Citrus fruits - Proc. of the Am. Soc. for Hort. Science, 52, 245-250, 1948.
207. TURRELL, F.M. : A study of the physiological effects of elemental sulphur dust on citrus fruits - Plant physiology, 25, 1, 13-62, 1950.
208. TURRELL, F.M. and SCOTT, F.M.: Effect of elemental sulfur dust on growth of citrus leaves, and its relation to the buffer capacity of the leaf-tissue fluid - Am. Journal of Botany, 38, 7, 560-566, 1951.
209. TURRELL, F.M., CHERVENAK, M. and VANSELOW, A.P.: Inorganic composition of exudate from Eureka lemons treated with elemental sulfur dust - Am. Journ. of Bot., 39, 10, 693-699, 1952.
210. TURRELL, F.M. : Effect of quality and intensity of solar radiation on injury of lemon fruit by sulphur treatment in the field - Plant Physiology, 28, 2, 151-176, 1953.
211. UNGER, K. : Agrarmeteorologische Studien I. Agrarmeteorologische Untersuchungen bei der Leistungsprüfung - Abh. des MHD der DDR, 3, 19, 1953.
212. UNGER, K. : Agrarmeteorologische Studien. II. Bodenfeuchtigkeitsbestimmungen unter einer Standardfläche und unter verschiedenen Pflanzenbeständen zur Charakterisierung der Dürrewirkung - Abh. des MHD der DDR, 3, 19, 1953.
213. UNGER, K. and FABIG, F.: Ueber den Einfluss der Temperatur auf die Erntezeit und Ertragsicherheit bei einigen Tomatensorten - Der Züchter, 23, 9, 257-266, 1953.
214. UNGER, K. : Zur klimatologischen Analyse der Kartoffel-Abbau und Gesundheitslagen - Angew. Met., 2, 26-32, 1954.
215. UNGER, K. and SÖRGEL, G.: Ueber den Befallsverlauf von Mycosphaerella pinodes (Berk et Blox), STONE, Ascochyta pinodella JONES und Ascochyta pisi LIB. an Erbsen am natürlichen Standort - Der Züchter, 24, 2/3, 56-68, 1954.
216. UNGER, K. and MÜLLER, H.J.: Ueber die Bedeutung der Zusammenhänge zwischen Witterung und Blattlausflug für die Probleme des Kartoffelabbaus - Forschungen und Fortschritte, 29, 8, 229-238, 1955.
217. UNGER, K. and SCHNEIDER, A.: Ueber Zusammenhänge zwischen der Reifeentwicklung von Gemüseerbsen und bestimmten mikrometeorologischen Faktoren - Der Züchter, 26, 1/2, 12-22, 1956.
218. UNGER, K. : Zum Einfluss der klimatischen Standortfaktoren auf die Blühzeit und den Blühverlauf einiger Gemüseerbsensorten - Ztschr. für Acker- und Pflanzenbau, 102, 1, 69-80, 1956.
219. WILLIAMS, C.B. : The cotton plant in relation to temperature and rainfall - Min. Agr. Egypt. Tech. & Scien. Service Bull., 32, 1-5, 1923.
220. WILLIAMS, C.B. : Cotton growing in relation to climate in Egypt and the Sudan - Min. Agr. Egypt. Tech. & Scient. Service Bull., 47, 1-31, 1925.

AGRICULTURAL PHENOLOGY

1. KNOCH, K. : Bericht über den phänologischen Reichsdienst und Vorlage der phänologischen Karten des Reichsgebiets - Organisation Météorologique Intern., Leiden, 36, 77-79.
2. MARCELLO, A. : Osservazioni fito-genologiche in pascoli alpini - Atti del R. Ist. Ven. di Scienze ed Arti, 93, 1933-34.
3. MARCELLO, A. : Osservazioni fito-fenologiche in stazioni glacio-nivali - Atti del R. Ist. Ven. di Scienze Lettere ed Arti, 94, 1934-35.
4. MARCELLO, A. : Bioclina e panificazione - L'Italia agricola, 72, 6, 1935.
5. MARCELLO, A. : Nuovi criterii per osservazioni fito-fenologiche - Nuovo Giornale Botanico Italiano, n.s., 42, 1935.
6. MARCELLO, A. : Reazioni morfologiche e fenologiche di alcune piante in funzione dell'ambiente - Nuovo Giornale Botanico Italiano, n.s., 42, 1935.
7. MARCELLO, A. : Osservazioni fito-fenologiche raccolte nel 1933 - Nuovo Giornale Botanico Italiano, n.s., 41, 1935.
8. MARCELLO, A. : Vetta e valico alpino. Osservazioni fito-fenologiche - Atti del R. Ist. Ven. di Scienze Lettere ed Arti, 94, 1934-35.
9. MARCELLO, A. : Erborizzazioni ienali nelle Alpi - Nuovo Giornale Botanico Italiano, n.s., 43, 1936.
10. MARCELLO, A. : Livelli di fioritura e pascoli alpini - Nuovo Giornale Botanico Italiano, n.s., 44, 1937.
11. MARCELLO, A. : Un problema di ecologia alpina Erica carnea L. - Atti dell'Ist. Ven. di Scienze Lettere ed Arti, 103, 1943-44.
12. MARCELLO, A. : Sintesi di una nuova tecnica fito-fenologica - Volume per le onoranze al Prof. Giuseppe Gola, Torino, 1947.
13. MARCELLO, A. : Ecologia e sinfenologia - Nuovo Giornale Botanico Italiano, n.s., 43, 1950.
14. MARCELLO, A. : Rilevamenti fito-fenologici all'isola d'Elba - Nuovo Giornale Botanico Italiano, n.s., 58, 1951.
15. MARCELLO, A. : Cicli fenologici nell'ambiente e loro rilevamento - Nuovo Giornale Botanico Italiano, n.s., 58, 1951.
16. MARCELLO, A. : Per la ricostituzione della Rete Fenologica Italiana - Nuovo Giornale Botanico Italiano, n.s., 58, 1951.
17. MARCELLO, A. : Pour l'institution d'un réseau phytophénologique international au service de l'Agriculture - Atti della Confédération européenne de l'Agriculture - Atti della Confédération européenne de l'Agriculture, 1951.
18. MARCELLO, A. : Un nuovo studio sinfenologico sull'Europa centrale di F. Schnelle - Nuovo Giornale Botanico Italiano, n.s., 59, 1952.
19. MARCELLO, A. : Piante e bioclina a Venezia - Minerva Medica Anno 43, 2, 79, 1952.
20. MARCELLO, A. : Su alcune osservazioni ecologiche e sinfenologiche di Leonardo da Vinci - Atti dell'Ist. Ven. di Scienze Lettere ed Arti, 111, 1952-53.
21. MARCELLO, A. : Il concetto di interazione in sinfenologia - Atti dell'Ist. Ven. di Scienze Lettere ed Arti, 111, 1952-53.
22. MARCELLO, A. : La nuova fenologia - Annali dell'Accademia Italiana di Scienze Forestali, 2, 1953.
23. MARCELLO, A. : De la méthode en Symphénologie - La Ricerca Scientifica Anno 25°, 6, 1954.
24. MARCELLO, A. : Atlante Fenologico per il servizio di rilevamento della Rete Fenologica Italiana - Accademia delle Scienze Forestali, Firenze, 1954.
25. MARCELLO, A. : Rete Fenologica Italia. Relazione riassuntiva sull'attività svolta durante gli anni 1952-53 - Nuovo Giornale Botanico n.s., 60, 1954.
26. MARCELLO, A. : Vicenda climatica ed avventura poliana - Volume nel VII° centenario della nascita di Marco Polo. Ist. Ven. di Scienze Lettere ed Arti - Venezia, 1955.
27. MARCELLO, A. : Rete Fenologica Italiana. Relazione dell'attività svolte nel 1954 - Nuovo Giornale Botanico Italiano n.s., 61, 1955.
28. MCMILLAN, C. : Nature of plant community. IV. Phenological variation within five woodland communities under controlled temperatures - Amer. Jour. Bot. 44, 154-163, 1957.
29. NOMURA, K. : Studies on the phenological type of the biological phenomena in Japan - Physiology and Ecology, 4, 1-2, 1950.
30. PASCALE, A.J. : Mapa fenológico del trigo en la República Argentina - Meteoros, 2, 1-2, 1952.
31. PASCALE, A.J. : Mapa fenológico del maíz en la República Argentina - Meteoros, 3, 4, 1953.

32. PRIMAULT, B. : Cinquant d'observations phénologiques systématiques en Suisse - Annalen der Schw. Meteorol. Zentralanstalt, Zürich, 1956.
33. PRIMAULT, B. : Contribution à l'Etude des réactions végétales aux éléments météorologiques. (L'Apparition du printemps dans le Canton de Neuchâtel de 1951 à 1954) - Bull. de la Soc. Neuchâteloise des Sciences naturelles, 80, 115-162, 1957.
34. RAMDAS, L.A. and MALLIK, A.K.: Phenology in India - Northern India Science Ass.
35. SAUBERER, F. : Ueber die Agrarmeteorologie und die Phänologie in Oesterreich - Wetter und Leben, 7, 257, 1955.
36. WILLIAMS, C.B. : Phenology - the study of the seasons - The advancement of Science, 11, 43, 267-270, 1954.

FOREST BIOCLIMATOLOGY

1. ALLEWELDT, G. : Der Einfluss von Photoperiode und Temperatur auf Wachstum und Entwicklung von Holzpflanzen unter besonderer Berücksichtigung der Gattung Vitis - Vitis, 1, 159-180, 1957.
2. AULITZKY, H. : Forstmeteorologische Untersuchungen an der Wald- und Baumgrenze in den Zentralalpen - Archiv f. Met. Geophys. Bioklimat., Wien, B, 3,3, 1953.
3. AULITZKY, H. : Ueber mikroklimatische Untersuchungen an der oberen Waldgrenze zum Zwecke der Lawinenvorbeugung - Wetter und Leben, 6, 5/6, 1954.
4. AULITZKY, H. : Waldkrone, Kleinklima und Aufforstung - Zentralblatt für das ges. Forstwesen, 73, 1/2, 1954.
5. AULITZKY, H. : Die Bedeutung meteorologischer und kleinklimatischer Unterlagen für Aufforstungen im Hochgebirge - Wetter und Leben, Wien, 7, 8/11, 1955.
6. AULITZKY, H. : Forest Bioclimatology. Areal report: Symposium Int. Soc. Bioclim. & Biomet. Paris, 29-31 August 1956.
7. AULITZKY, H. : Waldbaulich-ökologische Fragen an der Waldgrenze - Zentralblatt f.d. ges. Forstwesen, 75, 1, 18-33, 1958 (10 Abb.).
8. BOYKO, H. : On forest types of the semi-arid areas at lower latitudes - Palest Jnl. of Bot., Rehovot Series, 5, 1-21, 1945.
9. BOYKO, H. : A laurel forest in Palestine - Pal.J. of Bot., Reh.Ser., 6, 1-13, 1947.
10. BOYKO, H. : The decisive rôle of sun-radiation in the distribution of Quercus Aegilops L. - Pal.J. of Bot. Reh.Ser., 6, 20-22, 1947.
11. CIESLAR, A. : Die Bedeutung klimatischer Varietäten unserer Holzarten für den Waldbau - Zentralbl. f. das Ges. Forstw., 33, 1-19, 1907.
12. COTTAM, W.P. : Prevernal leafing of aspen in Utah mountains - Arnold Arboretum Jnl. 35, 239-248, 1954.
13. CRITCHFIELD, W.B. : Geographic variation in PINUS CONTORTA - Maria Moors Cabot Foundation (Harvard Univ.), 3, 1957.
14. DAVIDSON, H. and HAMMER, C.L.: Photoperiodic responses of selected woody ornamental shrubs - Mich. Agr. Expt. Sta. Quart. Bull., 40, 327-343, 1957.
15. DOWNS, R.J. and BORTHWICK, H.A.: Effects of photoperiod on growth of trees - Bot. Gaz., 117, 310-326, 1956.
16. ENGLER, A. : Einfluss der Provenienz des Samens auf die Eigenschaften der forstlichen Holzgewächse - Schweiz. Zentralanstalt f. Forstl. Versuchsw. Mitt., 8, 81-236, 1905.
17. ERMICH, K. : Forest and Climate - Wszechswist, 75-80, 1949.
18. ERMICH, K. : Climate Indices for Forest Economy - Instytut Badawczy Leśnictwa PW-RiL, Warszawa, 36, 1-23, 1951.
19. ERMICH, K. : The influence of the climatic factors upon the growth of English Oak, (Quercus robur L.) and Scotch Pine (Pinus sylvestris L.) - Prace Rolniczo-Leśne PAU., 68, 1-61, 1953.
20. ERMICH, K. and MERGENTALER, J.: The dependence between the growth of trees and the solar climate - Rocznik Dendrologiczny PTB, 9, 215-224, 1953.
21. ERMICH, K. and MERGENTALER, J.: The dependence of the diameter growth of trees from Tatra Mountains on the climatic fluctuations - Acta Societatis Botanicorum Poloniae, 24, 245-273, 1955.
22. ERMICH, K. and MERGENTALER, J.: Ecological investigations in two forest communities of the Kalwaria Zebrzydowska, Part I.: Thermic relations - Fragmenta Floristica et Geobotanica, 2, 28-49, 1956.
23. ERMICH, K. and ZURZYCKA, A.: Ecological investigations in two forest communities of the Kalwaria Zebrzydowska. Part II.: Humidity and light relations, Osmotic pressure - Fragmenta Floristica et Geobotanica, 2, 50-71, 1956.

24. ERMICH, K. and ZURZYCKA, A.: The light condition in the Patsch of Mountain Pine of Tatra - *Fragmenta Floristica et Geobotanica*, 3, 1957.
25. FAIRBAIRN, W.A. : Classification and description of vegetation types of Niger Botany , French W. Africa - Imperial Forestry Institute Paper 23, 1943.
26. FAIRBAIRN, W.A. and ANDERSON, M.L.: Division of Scotland into climatic sub-regions as an aid to silviculture - Forestry Department, Univ. of Edinburgh, Bulletin no. 1, 1955.
27. GAUSSEN, H. : Le milieu physique et la forêt au Portugal - Toulouse, 1940.
28. GAUSSEN, H. : Les résineux d'Afrique du Nord - Ecologie, Paris, 1952.
29. GIACOBBE, A. : L'Ecologia dell'Abies alba. Nota IV. L'eliofilia nel l'Appennino - *Travaux du Laboratoire forestier de l'Université de Toulouse*, 1951.
30. GIACOBBE, A. : L'écologie du Sapin (Abies alba). Note V: Le climagramme pour l'Alpennin - *Récueil des Travaux du Laboratoire de Botanique de la Fac. des Sc. Université de Montpellier*, no. 6, 1953.
31. GIACOBBE, A. : Les effets écologiques de l'aridité du climat sur la végétation méditerranéenne; héliophilie et configuration morphologique des Forêts - *Plant Ecology, Proc. of the Montpellier Symposium UNESCO*, 1954.
32. GLOCKER, KRUESMANN and FORTMANN: Beiträge zur Kenntnis der Industriefestigkeit der Gehölze - *Schriften zur Förderung des Gartenbaues*, 10, 1957.
33. GREHN, J. : Ueber Spaltungserscheinungen und photoperiodische Einflüsse bei Kreuzungen innerhalb der Sektion Leuke Duby - *Ztschr. Forstgenet. Forstpflanzenz.*, 1, 61-69, 1952.
34. GRULL, H. and WETTSTEIN, W.: Photo- und thermoperiodische Einflüsse auf das Wachstum der Kiefer (Pinus silvestris) - *Holzforschung*, 9, 81-84, 1955.
35. GRUNOW, J. : Erfassung zusätzlicher Niederschlagsmengen am Waldboden aus Nebelablagerungen - *Wetter und Leben*, Wien, 7, 262-263, 1955.
36. HAARLØV, N. : Measurements of temperature in Bark and Wood of Sitka Spruce (Picea sitchensis) with special reference to temperature in galleries of Hylesinus (Dendroctonus) micans (in Danish) - *Det Forstlige Forsøgs-va sen i Danmark, Copenhagen*, 21, 1-91, 1952.
37. HOFFMAN, D. : Die Rolle des Photoperiodismus in der Forstpflanzenzüchtung - *Ztschr. Forstgenetik. Forstpflanzenz.*, 2, 44-47, 1953.
38. HUECK, K.C.A. : Die Araukarienwälder des nördlichen Patagoniens - *Ztschr. für Weltforstwirtschaft*, 15, 1952.
39. HUECK, K.C.A. : Distribuição e habitat natural do Pinheiro do Brasil - *Bol. Facult. Cienc. e Letras Univ. S. Paulo, Botanica, Sao Paulo*, 10, 1953.
40. HUECK, K.C.A. : Problemas e importancia pratica da Fitosociologia no Estado de Sao Paulo - *Secretaria da Agricultura, Sao Paulo*, 1953.
41. HUECK, K.C.A. : Der Anden-Erlenwald in der Provinz Tucuman - *Angew. Pflanzensociologie, Festschrift Aichinger, Klagenuft*, 1954.
42. HUECK, K.C.A. : Waldbäume und Waldtypen aus N.W. Argentinien - *Fritz Haller Verlag, Berlin*, 1954.
43. HUECK, K.C.A. : Pflanzengeographisch-forstwirtschaftliche Probleme aus dem nordwestlichen Argentinien - *Ztschr. f. Weltforstwirtschaft*, 1954.
44. HUECK, K.C.A. : Plantas e Formacoes organogenicas das Duras Paulistas - *Secr. da Agricultura, Sao Paulo*, 1955.
45. HUECK, K.C.A. : O mapeamento geografico e sua importancia pratica para a silvicultura - *Anuario Brasileiro de economia florestal, Rio*, 8, 1955.
46. HUECK, K.C.A. : Nouvelles cartes de la végétation sud-américaine et leur signification pour l'agriculture et la sylviculture - *Ann. Biol., Paris*, 31, 1955.
47. HUECK, K.C.A. : Bosques chaquenos e extracao de tanino no Brasil - *Revista Brasileira de Geografia, Rio*, 17, 1955.
48. HUECK, K.C.A. : Die natürliche Pflanzendecke Brasiliens als Grundlage für Land- und Forstwirtschaft - *Staden-Jahrbuch, Sao Paulo*, 4, 1956.
49. HUECK, K.C.A. : Araukarienwald und Kampproblem - *Der naturw. Unterricht, Frankfurt*, 9, 6, 1956.
50. HUECK, K.C.A. : Mapa fitogeografico do Estado de Sao Paulo - *Bolet. Paulista de Geografia, Sao Paulo*, 22, 1956.
51. HUECK, K.C.A. : Waldgeographische Gliederung des brasilianischen Amazonasbeckens - *Forstw. Zentralblatt*, 75, 11/12, 505-568, 1956.
52. HUSTICH, I. : The Scotch pine in northern most Finland and its dependence on the climate in the last decades - *Acta Botanica Fennica*, 42, 1-75, 1948.
53. IRGENS-MOLLER, F. : Ecotypic response to temperature and photoperiod in Douglasfir - *Forest Sci.*, 3, 79-83, 1957.
54. KIRA, T. : Forest zones of Japan - *Sapporo & Tokyo*, 1949.
55. KIRA, T. : Ecological situation of the deciduous conifer forest climax in Eastern Siberia - In *"Explorations in the Great Khingan Mountains"* edited by Dr. K. Imanishi, 476-497, 1952. The Mainichi Press, Tokyo.

56. KNAPP, R. : Beiträge zur Kenntnis der Wälder im atlantischen Klimabereiche Deutschlands - Geobot. Mittlg., 2, 21-40, 1952.
57. KOCH, H.G. : Durchmesserzuwachsserhebungen an Waldbäumen zum Nachweis von Geländeklimaeinflüssen - Archiv für Forstwesen, Akademieverlag, Berlin, 5, 9/10, 16 pp., 1956.
58. KOCH, H.G. : Jahresringauszählungen an Waldbäumen zum Nachweis witterungsbedingter Zuwachsschwankungen - Annalen der Meteorologie, 8, 1/2, 21-33, 1957.
59. KOZLOWSKI, T.T. and WARD, R.C.: Seasonal height growth of conifers - Forest Sci., 3, 61-66, 1957.
60. KOZLOWSKI, T.T. and WARD, R.C.: Seasonal height growth of deciduous trees - Forest Sci., 3, 168-174, 1957.
61. KRAMER, P.J. : Some effects of various combinations of day and night temperatures and photoperiod on the height growth of loblolly pine seedlings - Forest Sci., 3, 45-55, 1957.
62. KRECMER, V. : Dew as a meteorological, pedological, physiological and ecological factor and dew in forestry - Lesnicka, Praha, 30, 4, 1951.
63. KRECMER, V. : The effect of forest on precipitations - Meteorologické Zprávy, Praha, 5, 1952.
64. KRECMER, V. : Methods of research and other problems of the influence of covering agricultural crops on the environment and growth of forest tree seedlings in the first year of their life - Práce výzkumných ústavů lesnických CSR, 3, 1953.
65. KRECMER, V. : To the question of covering agricultural crops in forestry - Meteorologické Zprávy, Praha, 8, 1-2, 1955.
66. KRECMER, V. : Climatological function of forests - 3rd nat. hydrometeorological Conference, September 1954, Praha, Docum., Praha, 1956.
67. LANGLET, O.: Studier over tallens fysiologiska variabilitet och dess samband med klimatet. Ett bidrag till kännedomen om tallens ekotyper. (Studien über die physiologische Variabilität der Kiefer und deren Zusammenhang mit dem Klima. Beiträge zur Kenntniss der Ökotypen von PINUS SILVESTRI L. - Medd. Skogsforsöksanst., 29, 219-470, 1936.
68. LANGLET, O. : Photoperiodismus und Provenienz bei der gemeinen Kiefer (Pinus silvestris L.) - Medd. Skogsforsöksanst., 33, 298-330, 1943.
69. LEMEE, G. : Recherches sur l'évolution forestière post-glaciaire en Auvergne. III. Modifications forestières et climatiques dans les Monts Dore d'après l'analyse pollinique - Rev.Sc.nat. d'Auv., 8, 84-125, 1942.
70. LEMEE, G. : Recherches sur l'économie de l'eau chez les sous-arbrisseaux xéromorphes de landes - Ann. Sc. nat. Bot., IIe série, 7, 53-85, 1946.
71. LEMEE, G. : L'histoire forestière et le climat contemporain des lignites de Savoie et de la tourbe wurmienne d'Armoir d'après l'analyse pollinique - Trav. Labor. de Géol., Grenoble, 29, 167-180, 1951.
72. LEMEE, G. : L'économie de l'eau chez quelques Graminées vivaces du Sahara septentrional - Vegetatio, 5/6, 534-541, 1954.
73. LEMEE, G. : L'évolution de la forêt française au cours du Quaternaire d'après les analyses polliniques - Rev. forst., 6, 442-460, 1955.
74. LEMEE, G. : Influence de l'alimentation en eau et de l'ombrage sur l'économie hydrique et la photosynthèse du cacaoyer - L'Agron. trop.no. 5, 592-603, 1955.
75. MEISSNER, O. : Jahresringe der Bäume und Klimaschwankungen - Bioklim.Beiblätter, 10, 2/3, 88, 1943.
76. MOREAU, R.E. : Eco-climatic data for closed evergreen forest in tropical Africa - J. Linn. Soc. Zool., 39, 285-293, 1935.
77. MORELLO, J. : Multiplicación de arbustos en las mesetas araucanas del valle de Santa María (Provincia de Tucumán) - Bol.Soc.Arg.Bot., 3, 4, 207-217, 1951.
78. MORELLO, J. : Regiones áridas de la Argentina. II. Balance de agua de los arbustos resinosos de follaje permanente - Rev.Agr.Norocste Arg., 1955.
79. NÄGELI, W. : Lichtmessungen im Freiland und in geschlossenen Altholzbeständen - Mitteilungen der Schweizerischen Anstalt für das forstlichen Versuchswesen, 21, 2, 1940.
80. NÄGELI, W. : Untersuchungen über die Windverhältnisse im Bereich von Windschutzstreifen - Mitteilungen der Schw. Anstalt f.d. forstlichen Versuchswesen, 23, 1, 1943.
81. NÄGELI, W. : Weitere Untersuchungen über die Windverhältnisse im Bereich von Windschutzstreifen - Mitteilungen der Schw. Anstalt f.d. forstlichen Versuchswesen, 24, 2, 1946.
82. NÄGELI, W. : Untersuchungen über die Windverhältnisse im Bereich von Schilfrohrwänden - Mitt. der Schw. Anst.f.d. forstl. Versuchswesen, 29, 2, 1953.

83. NITSCH, J.P. : Growth responses of woody plants to photoperiodic stimuli - Amer. Soc. Hort. Proc., 70, 512-525, 1957.
84. OLMSTED, C.E. : Experiments on photoperiodism, dormancy, and leaf age and abscission in sugar maple - Bot. Gaz., 112, 365-393, 1951.
85. OVINGTON, J.D. : A comparison of rainfall in different woodlands - Forestry, 27, 41-53, 1954.
86. OVINGTON, J.D. and MADGWICK, H.A.J.: A comparison of light in different woodlands - Forestry, 28, 141-146, 1955.
87. PASZYNSKI, J. : Die Niederschläge im Obergerbiet und ihr Zusammenhang mit Höhe und Bewaldung - Prace Geograficzne, 4, 90, 1955.
88. PEREIRA, H.C. : Interception of rainfall by Cypress plantations - E.African Agric.J. 18, 73-76, 1952.
89. PEREIRA, H.C. : The physical importance of forest cover in the East African Highlands - E. African Agric. J., 19, 233-236, 1954.
90. PHILIPPIS, A. de : Classificazioni ed indici del clima, in rapporto alla vegetazione forestale italiana - Nuovo Giornale Botanico Italiano, 1937. (Translated by the Soil Conservation Service, U.S.A.).
91. PHILIPPIS, A. de : Ecologie forestière et Phytoclimatologie - Unasylva, 1, 1951.
92. PHILLIPS, J.E. : Effect of day length on dormancy in tree seedlings - Jour. Forestry, 39, 55-59, 1941.
93. PLATT, R.B. and THOMAS, J.: Maximum temperature tolerances of loblolly pine and sweet gum seedlings under varying soil, moisture and light conditions - Journ. of the Tennessee Acad. of Sci., 29, 186, 1954.
94. PLATT, R.B. : Vertical and horizontal gradients occurring within a woodland and adjacent old field in the Coastal Plain - Bull. of the Ecol. Soc. of America, 35, 60-61, 1954.
95. PRESCOTT, J.A. and LANE POOLE, C.E.: The climatology of the introduction of pines in the Mediterranean environment to Australia - Trans. Roy. Soc. of S.Aust., 71, 67-90, 1947.
96. PRIMAULT, B. : Contribution à l'étude de l'influence des éléments météorologiques sur l'accroissement des forêts - Thèse de Doctorat à l'E.P.F. Geofisica pura e applicata, Milan, 24, 1953.
97. PRIMAULT, B. : Quelques considérations sur les conditions météorologiques et l'accroissement des forêts - Journal forestier suisse. 104, 10, 1953.
98. PRIMAULT, B. and KUHN, W.: De l'opportunité d'utiliser les coefficients de corrélation en météorologie forestière - Geofisica pura e applicata, Milan, 28, 246-260, 1954.
99. PRIMAULT, B. : En réponse à une communication de l'UNESCO - La Forêt, 9, 11, 1956.
100. PRIMAULT, B. : Bäume als Indikatoren für schlechte Luft - Zürcher Woche, 43, 1956.
101. PURI, G.S. : The secular or pre-historical factor and its application to forest ecology and silviculture - Ind. Forest Ranger College Annual, 71-75, 1948.
102. PURI, G.S. : Forest ecology and evaporation measurements in India - Science and Culture, 15, 465-467, 1950.
103. PURI, G.S. : A single value climatic factor in forest ecology - Proc. 37th Ind. Sci. Cong., Abstr., Poona, 1950.
104. PURI, G.S. : The influence of forests on precipitation and the possibilities of rain making in forest areas of India - Contributed to Symposium on Artificial Rainfall, 1953.
105. PURI, G.S. : Seasonal variation in foliar composition of some Indian forest trees - Journ. Ind. Bot. Soc., 33, 382-393, 1954.
106. REDMOND, D.R. : Studies in Forest Pathology. XV. Rootlets, Mycorrhiza, and Soil temperatures in relation to Birch Dieback - Can. J. Bot. 33, 595-627, 1955.
107. SCHIELDT PAULSEN, H., GODSKE, C.L. and MARSTRANDER, J.H.: Forstmeteorologische undersøkelser i Os Prestegårdsskog sommeren 1944 - Bergen Museums Arbok, Nat. vit. rekke, 2, 1-16, 1945.
108. SCHIELDT PAULSEN, H. and GODSKE, C.L.: The Forest Precipitation - Universitete i Bergen Arbok, 8, 1949.
109. SMILEY, T.L. : Forest Bioclimatology - See Areal report Dr. D.H.K. Lee. Symposium Int. Soc. Bioclim. and Biomet. Paris 29-31 August 1956.
110. SMOLIK, L. : Microclima of a forest and of a grass land near the river Labe - Lesnická práce, 16, 10, 461, 1937.
111. SMOLIK, L. : Temperature in the snow cover upon the forest grass land - Sborník, CAZ. 1938.
112. SMOLIK, L. : Influence of the forest upon the climate - Slovo naroda, 1946.
113. TRANQUILLINI, W. and PISEK, A.: Transpiration und Wasserhaushalt der Fichte (Picea excelsa) bei zunehmender Luft- und Bodentrockenheit - Physiol. Plant, 4, 1, 1951.

114. TRANQUILLINI, W. and PISEK, A.: Assimilation und Kohlenstoffhaushalt in der Krone von Fichten- (*Picea excelsa* LINK) und Rotbuchenbäumen (*Fagus silvatica* L.) - "flora" oder allgemeine botanische Zeitung, 141, 237, 1954.
115. TRANQUILLINI, W. : Die Bedeutung des Lichtes und der Temperatur für die Kohlensäureassimilation von *PINUS CEMBRA* - Jungwuchs an einem Hochalpinen Standort - *Planta*, 46, 154-178, 1955.
116. TRANQUILLINI, W. : Vom Existenzkampf des Baumes im Hochgebirge - *Jahrb. Ver. Schutz Alpenpfl. u. Tiere*, 1956.
117. TURRELL, F.M. : The relation of internal surface to intercellular space in foliage leaves - *Science*, 92, 2385, 244, 1940.
118. VAARTAJA, O. : High surface soil temperatures. On methods of investigation and thermocouple observations on a wooded heath in the south of Finland - *Oikos*, 1,1, 6-28, 1949.
119. VAARTAJA, O. : On factors affecting the initial development of pine - *Oikos*, 2, 1, 89-108, 1950.
120. VAARTAJA, O. : Forest humus quality and light conditions as factors influencing damping off - *Phytopathology*, 42, 501-506, 1952.
121. VAARTAJA, O. : Temperature and evaporation at and near ground level on certain forest sites - *Can. Jour. Bot.*, 20, 160-183, 1954.
122. VAARTAJA, O. : Photoperiodic ecotypes of trees - *Canad. Jour. Bot.*, 32, 392-399, 1954.
123. VAARTAJA, O. : Factors causing mortality of tree seeds and succulent seedlings - *Acta Forestalia Fennica*, 62, 3, 1954.
124. VAARTAJA, O. : Photoperiodic responses in seedlings of northern tree species - *Can. Jour. Bot.*, 35, 133-138, 1957.
125. VAARTAJA, O. : Experimental evidence of photoperiodic ecotypes in several tree species - *Ecol. Soc. Amer. Bull.*, 38, 76, 1957.
126. VAARTAJA, O. : Photoperiodic responses in seedlings of northern tree species - *Can. Jnl. of Botany*, 35, 133-138, 1957.
127. VAARTAJA, O. : Evidence of photoperiodic ecotypes in trees - *Ecol. Monographs*, 29, 2, 91-111, 1959.
128. WAREING, P.F. : Photoperiodism in woody species - *Forestry*, 22, 211-221, 1948.
129. WAREING, P.F. : Photoperiodism in woody plants - *Ann. Rev. Plant Physiol.*, 7, 191-214, 1956.
130. WAREING, P.F. and ROBERTS, D.L.: Photoperiodic control of cambial activity in *ROBINIA PSEUDOACACIA* L. - *New Phytol.*, 55, 356-366, 1956.
131. WASSINK, E.C. and WIERSMA, J.H.: Daylength responses of some forest trees - *Acta Bot. Neerl.*, 4, 657-670, 1955.
132. WAXMAN, S. : The effect of the length of day on the growth of woody plants - *Plant Propagators Soc. Proc.*, 5, 47-50, 1956.
133. ZELAWSKI, W. : Further researches on the photoperiodic reaction in seedlings of the European larch (*LARIX EUROPAEA* D.C.) (Polish with English summary) - *Soc. Bot. Polon. Acta.*, 26, 79-103, 1957.

PHYSIOLOGICAL PHYTO-BIOCLIMATOLOGY

1. BORREGUERO, G.C. : Observaciones Fisiologicas en el genero *Fusarium* - *Rev. Clinic. Española*, 7, 23, 2, 124-129, 1946.

PART III

ZOOLOGICAL BIOCLIMATOLOGY

(1959)

Section A : General Zoological bioclimatology

Section B : Entomological bioclimatology

ZOOLOGICAL BIOCLIMATOLOGY
Section B: Entomological Bioclimatology

MICROCLIMATE, DIURNAL RHYTHMS AND THE CONQUEST
OF THE LAND BY ARTHROPODS.

by

Dr. J.L. Cloudsley-Thompson, F.L.S. (Great Britain) *

I. INTRODUCTION.

The conquest of the land by insects and other arthropods illustrates a number of fundamental points of behaviour in the responses of the animals to the simple stimuli provided by the bioclimatological conditions of their physical environment.

The distribution of animals is determined not only by a number of geological factors and by the physical conditions of the environment - light, temperature, moisture, wind, acidity or alkalinity and so on - but also by its biotic factors which include plants and other animals. In the present review, however, I shall be concerned only with interpreting the distribution of terrestrial arthropods in terms of their psychological responses and behavioural reactions to the former. The significance of these at once becomes apparent when one remembers that most of the oldest fossils found in geological strata are marine and the sea is a relatively stable environment, not subjected to violent changes in temperature, salinity, etc. In this medium were evolved all the major groups of animals and during succeeding epochs most of them managed to conquer the land to a greater or lesser extent.

Life on land entails a number of problems: larger animals require structural support; respiratory organs must become adapted for breathing air and there is no surrounding water into which toxic excretory compounds can diffuse. Small animals have a very large surface area in proportion to their mass; consequently the conservation of water and the maintenance of a fairly constant internal medium are especially important. Many terrestrial invertebrates such as insects, spiders and mites avoid becoming desiccated by the development of a thin layer of wax which is relatively impervious to water vapour. Of course, such a layer is also impervious to oxygen and carbon dioxide: a respiratory mechanism has therefore had to be evolved which permits gaseous exchange to take place whilst restricting water-loss to a minimum. The spiracles of insects and the lung books of spiders and other arachnids are normally kept closed by means of special muscles, and only when carbon dioxide in the body begins to accumulate are they opened to facilitate respiration. It can easily be shown experimentally by weighing that the rate of water-loss by evaporation from an insect or arachnid is greatly increased when 5 per cent carbon dioxide is present in the atmosphere, as this results in the respiratory apertures remaining fully open. Before moulting takes place a new wax layer is secreted beneath the old cuticle that is due to be cast off, so that the ecdysis is effected with a minimum loss of water (39, 43, etc.).

In addition to such morphological and physiological adaptations to life on land, insects and arachnids have evolved special excretory products - uric acid and guanine respectively - which are extremely insoluble. Consequently nitrogenous waste matter can be eliminated from the body in a dry state and no water is lost in the process.

Another method of avoiding desiccation on dry land is found in woodlice, centipedes, millipedes, springtails and many other small animals which remain most, if not all of the daytime in a damp or humid environment under stones, fallen leaves, and in crevices under the bark of trees.

The terrestrial arthropods can, therefore, be roughly divided on an ecological basis into two main groups. The first includes woodlice, centipedes, millipedes and their allies which, in dry air lose water rapidly by transpiration through their integuments (8, 23, 24, etc., 34). Consequently they are restricted by reflex behaviour mechanisms to damp, dark habitats which they leave only at night when the temperature falls and the relative humidity of the atmosphere increases (18, 19, 20, 22, 23, 25, 27, etc.).

The second group includes most insects and arachnids; these are comparatively independent of moist surroundings because their integument possesses an impervious layer of wax which prevents desiccation (13, 14, 15, 16, 20, 30, etc.).

* Note: Dept. of Zoology, University of London, King's College, Strand, London W.C. 2, Great Britain.

Each aspect of adaptation to life on land affects, and is affected by other aspects. For example, if the integument is rigid and provides support, then growth becomes impossible except by moulting and this limits size. Size is also limited where the respiratory system consists of tracheae and tracheoles. The physiology of nutrition and excretion are closely concerned with water conservation. Superimposed upon such basic morphological and physiological requirements are the numerous concomitants of behaviour and ecology, for orientation and behaviour mechanisms must also be evolved to retain organisms in environments to which they are suited, to find food, mate and, indeed, carry out the functions essential to their continued existence.

II. WOODLICE AND THE LAND HABITAT

Woodlice show a diurnal rhythm of activity and tend to wander abroad during the night, when the temperature drops and the humidity of the atmosphere increases. This rhythm is correlated with alternating light and darkness and not with fluctuating temperature and humidity although these factors of the environment are probably of greater importance in their daily lives (11). When taken into the open, they are stimulated by light and drought so that they run actively until, by chance, they reach some dark damp spot where they come to rest. They can run directly away from light, but one searches in vain to find any directed orientation towards moist air or damp surfaces. The animals run aimlessly, turning first one way then the other. But, as the air becomes damper, their speed decreases (36,42).

This behaviour can be illustrated by a simple analogy with motor cars. Suppose a procession of cars is travelling at a steady speed of 50 m.p.h. outside a town. On entering the restricted zone the speed falls to 20 m.p.h. which is retained through the urban area. Now it will be seen that the cars are closer together. If the front vehicle is halted at traffic lights, an aggregation will occur which disappears when the cars speed up again in the open country. The time interval between the passage of one car and the next will be the same throughout the town and the country but, because they move more slowly in the built-up area, they will automatically be closer together. So it is with woodlice which move more slowly and consequently aggregate in damper air.

In addition to this simple ORTHOKINESIS or variation in linear velocity, woodlice tend to turn more and more frequently as they pass through a zone of moist air, until they finally come to rest. This is a KLINOKINESIS similar to that found in *PARAMECIUM CAUDATUM* which aggregates in regions of slight acidity. It is by such purely mechanical reactions that woodlice aggregate under bark and stones. Experiments have shown that the animals are not in any way attracted towards one another. They merely gather in the same place as a result of behaviour reactions which drive them away from light and from dry places (35, 42).

By means of choice-chamber apparatus it has been found that the intensity of the humidity response of the woodlice is less in darkness than in light and still less in the nocturnal phase; it increases with desiccation. Movement away from light is more marked in animals which have been in darkness for some time, and is still initially shown in dry air whereas control animals move towards the light when humidity is sufficiently reduced. Finally, the active phase is correlated with increased sensitivity to external conditions such as the carbon dioxide content of the air.

These experimental results can be related to the nocturnal ecology of the species. Woodlice are often to be seen wandering in dry places at night: as we have seen, they spend the daytime under stones, logs and in other damp, dark situations where environmental conditions are relatively constant. If nightfall involved only a reduction of the factors which normally operate to restrict them to their daytime habitats, it seems unlikely that they would have any occasion to leave them. The endogenous diurnal rhythm or physiological "clock", however, will engender locomotory activity in some individuals at least at nightfall when they are especially sensitive and easily disturbed. Even if the majority are exposed to daylight only occasionally, this may be sufficient to maintain their periodicity and keep it in phase with the 24-hour cycle (11).

Field observations on *P. SCABER* have indicated a shortening in the time of activity as the days lengthen to midsummer and also a lack of any clear relationship between activity and changes in temperature or relative humidity (2).

The decrease in the intensity of the humidity response at night enables woodlice to walk in dry places where they are never to be found during the day, and the increased photo-negative response after they have been conditioned to darkness ensures that they get under cover promptly at daybreak and thus avoid many potential predators. On the other hand, if their daytime habitat should dry up, the woodlice are not restrained there until they die of desiccation since they tend to become photopositive in dry air and thus are able to wander in the light until they find some other damp hiding place and again become photo-negative (11).

Even the desert woodlouse, *HEMILEPISTUS REAUMURI*, has a 24-hour periodicity and water relations essentially similar to those of woodlice from temperate regions. Although this species can withstand hot, dry conditions for some considerable time there is no "critical temperature" to indicate the presence of a cuticular layer of water-proofing wax (24).

The rate of water-loss by transpiration in a number of British species of woodlouse has been investigated by Edney (32,33,34). They have been shown to stand in the following order;
PHILOSCIA MUSCORUM > *ONISCUS ASELLUS* > *PORCELLIO SCABER* > *ARMADILLIDIUM VULGARE*.

At temperatures just above freezing the intensities of the reaction to humidity of all species are much reduced and do not differ significantly from one another. At higher temperatures (30° C) the humidity responses of *PH. MUSCORUM* and *O. ASELLUS* are again somewhat reduced, and it is suggested that this may be associated with a lower thermal death point and the need to effect a reduction in body temperature by evaporation (25).

The humidity responses of woodlice are shown to be correlated with saturation deficiency rather than with relative humidity. No specific sense organs are concerned with the appreciation of humidity and the response is probably engendered as a result of desiccation. It may result from the influence of increased concentration of the body fluids stimulating proprioceptors (42).

All species must spend the greater part of their time in an atmosphere that is saturated with water vapour, but there is considerable variation regarding their ability to withstand dry air, high temperatures and in the period of time during which they can venture into dry places.

PH. MUSCORUM is the most strongly photo-negative of the four species and there is a gradation through *O. ASELLUS* and *P. SCABER* to *A. VULGARE* which is the least so. *PH. MUSCORUM* has been shown by means of aktograph apparatus to be the most intensely nocturnal in habit, *A. VULGARE* the least. It is therefore suggested that the degree of nocturnal activity may be correlated with the ability to withstand water-loss by transpiration (25).

Finally, it has been shown by measurements with a sensitive anemometer that the nocturnal emergence of woodlice in nature is inhibited by wind. This may well be because air currents tend to remove the shell of moist air that surrounds the transpiring animal (29). The relationship between wind-speed and the number of woodlice wandering abroad at night applies only above a minimum threshold temperature of about 5.5° C, however.

III. THE WATER-RELATIONS OF MILLIPEDES

Somewhat similar responses have been shown to occur in millipedes which also lose water rapidly in dry air. Some species may tend initially to move towards dry places but the reaction is gradually reversed as desiccation proceeds (38). Orientation is again entirely "KINETIC" or non-directional and in an experimental chamber in which a choice of humidities is provided, both the time spent and the distance covered are greater on the moist side.

Millipedes possess a clear endogenous 24-hour periodicity and emerge from the comparatively constant conditions within their daytime retreats under stones, bark logs and fallen leaves, under the stimulus provided by some internal physiological "chronometer".

In the case of the small British millipedes *BLANIULUS GUTTULATUS* and *OXIDUS GRACILIS*, an exogenous diurnal rhythm is primarily a response to light and darkness, the degree of nocturnal activity being correlated with the stimulus of falling temperature at dusk (9, 10) and no endogenous activity has been observed. In the large West African species *OXYDESMUS PLATYCERCUS* and *OPHISTREPTUS* SP., however, aktograph experiments have demonstrated an endogenous rhythm independent of fluctuating light and temperature and persisting for as long as 19 days in constant light and temperature. Locomotory activity is stimulated both by increases and by decreases of temperature and it is suggested that temperature fluctuations may be of primary importance in the synchronisation of the animals' activities with day and night, for the effect of light on their activity is slight and is perhaps an insignificant factor in their natural gloomy habitat in tropical forests (10).

The response of millipedes to moisture has some economic importance, for under conditions of drought they may be forced to attack growing crops for the sake of water. Thus outbreaks of the "spotted snake-millipede", *BLANIULUS GUTTALATUS*, tend to be stimulated by a dry spell following a period suitable to the reproduction of the species when the soil is damp, undisturbed and rich in humus. It has been shown experimentally that humus and rotting substances have a texture which is preferred by millipedes to that of living plant tissues, and that the animals are attracted to dilute concentrations of sugars (8).

No doubt a moist season combined with the use of farmyard manure or following some crop producing a considerable amount of humus, will engender a great increase in the number of millipedes in the soil, particularly if the ground is not disturbed by ploughing etc. They can be beneficial at this stage in aiding the breakdown of the humus, but if the following season is dry, even for a short spell, they may be compelled to attack crops for the sake of moisture. Once an attack has been initiated, the attraction of sugars in the plant sap will prevent them from returning to their normal diet of humus and decomposing matter. It is unlikely that damage by millipedes to crops with tough exteriors such as potatoes and mongolds can ever be primary, for not only do their weak mouth parts prevent them from gaining access, but in addition they are not attracted to unbroken skins of potatoes, only to cut surfaces. Once an entrance has been achieved, however, through mechanical damage or the bites of wire worms and other pests, the millipedes will eat out the entire centre of a potato and the damage they cause is often followed by fungal

attack. Furthermore, the fact that single potatoes have been found containing over a hundred *BLANIULUS GUTTULATUS* of all ages while the remainder of the crop was unharmed shows that they must have been attracted to a damaged tuber and could not have bred there (7).

Like woodlice, millipedes avoid the light, but with the exception of a directed response or "TAXIS" in those forms that possess eyes, their reactions are non-directional. When illuminated they crawl around until, by chance, they find themselves in darkness where they come to rest.

Although to millipedes, as to woodlice, humidity is the most important factor of the environment, as we have seen these animals are not able to find their way directly to damp places; instead, they are merely repelled kinetically by drought. Nevertheless this stereotyped and curiously negative behaviour is surprisingly effective in preventing them from wandering away from their normal habitats: but it does raise the problem of how dispersal can take place and new habitats become colonised (19).

There are a number of cases on record of millipedes, sometimes accompanied by centipedes and woodlice, migrating in vast armies. Occasionally they have crossed railways and been squashed in such numbers that locomotives have been impeded and sand has had to be strewn on the lines before their driving wheels would grip. At other times cattle have refused to graze on invaded pastures, wells have been filled with drowned corpses and workmen cultivating the fields have become nauseated and dizzy from the odour of millipedes crushed by their hoes. Such mass migrations, however, are of rare occurrence and local in extent, so that their net effect on distribution is probably negligible (6,27).

The explanation of the problem of distribution lies in the fact that, as already mentioned, the restraining mechanisms are somewhat relaxed during the night when the temperature falls and the humidity of the air increases. Thus it is at night that these creatures are enabled to disperse themselves and overcome the restrictions inherent in the physiology of their integuments.

IV. THE PARADOX OF RESPIRATION AND IMPERMEABILITY

The conflict between the incompatible requirements of respiratory exchange and prevention of water-loss in arthropods possessing a discrete cuticular wax layer has been illustrated by a comparison of the spiders *AMAUROBIUS* (= *CINIFLO*) *FEROX* and *A. SIMILIS* (26). Although there is some overlap in their territories, *A. SIMILIS* tends to inhabit somewhat drier environments than *A. FEROX*.

Aktograph experiments indicate that both species are nocturnal in habit, over 90 per cent of their activity taking place during the hours of darkness. When their water-relations are considered, however, it is found that there is a critical temperature at approximately 35° C above which both species quickly lose water by evaporation in dry air, but below this *A. FEROX* loses water more rapidly through its lung-books than does *A. SIMILIS*. The rate is almost doubled in both species when 10 per cent carbon dioxide is present, as this keeps the lung-books open. Conversely the length of time or survival in air of 50 per cent relative humidity, and in dry air is longer in *A. SIMILIS* than in *A. FEROX*, death ensuing when from 20 to 25 per cent of total weight has been lost by evaporation.

A. SIMILIS "tires" more rapidly than *A. FEROX* when forced to run at full speed without stopping, but both species can run for long periods when supplied with oxygen. *A. FEROX* becomes anaesthetised more quickly in ether vapour and has the larger number of leaves in its lung-books. Its greater "stamina", therefore, depends upon a proportionately larger respiratory surface acquired at the expense of great dependence upon environmental humidity (26).

Diurnal rhythms of locomotory activity in animals are especially marked in desert regions where the climatic changes between day and night are particularly great (3, 4, 5, 18, 24, 37, 44 etc.). Microclimatic readings have indicated a considerable reduction in temperature fluctuations in the daytime retreats of deserticolous animals. For example, Williams (44) showed that in August whilst the surface of the sand in a wadi near Cairo varied from 17.5° to 58.2° C, at a depth of 18 cms. the range was only 29.3° to 37.8° C; whilst the effect of shading was to lower the surface temperatures some 20° C. Buxton (3) found that the surface of clay desert in southern Israel rose about 20° C beyond the shade temperature and quoted a daily range of temperature from -0.5° to 37.2° C in December at Bir Milgha in Southern Tripoli. Further examples of this kind can be obtained from the publications referred to above.

In addition to the considerable reduction in temperature extremes between day and night, only a few centimetres down a hole or within a small cave, a time-lag occurs in these sheltered habitats so that when their inhabitants emerge at dusk they may actually be leaving a warmer for a cooler environment.

In hot climates, survival depends upon avoiding desiccation and keeping cool. Consequently where water is in short supply, there must inevitably be a conflict between the requirements of conserving water for vital purposes and of transpiring it for cooling. Arthropods are too small to withstand transpiration for long and therefore can exist only by avoiding true desert conditions and their physiological

adaptations to desert conditions involve changes in degree rather than the evolution of new mechanisms. Thus, the behavioural responses and water-relations of the desert woodlouse, *HEMILEPISTUS REAUMURI*, and of centipedes such as *SCOLOPENDRA CLAVIPES* of North Africa are essentially similar to those of species from temperate regions.

Scorpions, however, present a more difficult problem. *EUSCORPIUS GERMANUS* from Italy and the Tunisian species, *SCORPIO MAURUS*, *BUTHUS OCCITANUS* and *ANDROCTONUS AUSTRALIS* are all markedly nocturnal and experimentally show well marked periodicity even under constant conditions. They can be arranged in a series as regards their rate of water-loss and time of survival in dry air and this series is in agreement with what is known of their ecology. The integuments of the three Tunisian species are extremely effective in preventing water-loss and their cuticular wax-layers have a critical temperature as high as 65° C. It is difficult, therefore, to ascribe any ecological function other than the avoidance of enemies to the markedly nocturnal behaviour of these animals which may be quite capable throughout much of the year of surviving the high temperatures and low humidity encountered on the surface of the desert sand during the daytime (24).

According to Vachon (40,41) however, scorpions were not originally characteristic of desert regions; they represent the remains of an ancient fauna which lived under quite different conditions of temperature and humidity and manage to survive in conditions of heat and drought because of their subterranean habitat and nocturnal habits. They are responsive to micro-climatic variations and each species seems to have to live and reproduce within characteristic ecological conditions varying within close limits (see also 27).

Desert beetles also show marked diurnal rhythms of locomotory activity and although the figures for water-loss from British and North African species are in the same order of magnitude it may be not only that there has been some adaptation of the water relations of the latter to desert life, but also that the types found in the desert are those whose physiology and behaviour are already such as would enable them to colonise arid regions (24). It must be remembered, however, that the ability to lose SOME water by evaporation and thus achieve reduction in temperature may be an important factor for all Arthropoda.

Buxton (4) comments upon the diurnal and seasonal appearance of desert beetles and has shown that many large Tenebrionidae, such as *ADESMIA* spp. which are conspicuous in North Africa and Israel at all times of the day in March and April, seek shelter under stones and bushes in May and June. Bodenheimer (1) has also shown that the time of maximum activity of desert beetles varies with the season.

It could scarcely be expected that the time of activity of terrestrial arthropods should be rigidly correlated with their rate of water-loss. Temperature tolerance, rate of water-loss in dry air, time of activity and the microclimatic conditions of the habitats in which they live are all factors of importance to be considered when assessing the adaptations of the various species to terrestrial life.

V. CONCLUSION

It would be a mistake to regard the absence of a cuticular wax layer as a primitive characteristic, although the forms that lack one are so restricted in their choice of environment that they cannot be regarded as entirely successful land animals. Rather, it seems that a particular method has been exploited for surviving the conditions of life on land.

The terrestrial ancestors of the arthropods may at first have been restrained by behaviour mechanisms to damp environments. Later some of their descendants, woodlice, centipedes and millipedes, exploited still further this form of terrestrial life; while others, the arachnids and insects, acquired waterproof integuments and the other physiological and morphological mechanisms that must accompany them. In the absence of more direct evidence, physiology and ecology can merely suggest the course that evolution may have followed.

The general speeding up of life in a medium of abundant oxygen and negligible viscosity has resulted in increased specialisation of the central nervous system, and only the size limit imposed by respiratory requirements and the necessity for growth by moulting may have prevented the arthropods from dominating the terrestrial environment to an even greater extent - possibly even to the exclusion of the vertebrates, including man.


VI. REFERENCES

1. BODENHEIMER, F.S. : Studies on the ecology of Palestinian Coleoptera. II Seasonal and diurnal appearance and activity - Bull. Soc. R. Ent., Egypt, 1-2, 211-41, 1934.
2. BRERETON, J.L. : The distribution of woodland isopods - Oikos 8, 85-106, 1957.
3. BUXTON, P.A. : Animal life in deserts. A study of the fauna in relation to the environment, London, 1923.
4. BUXTON, P.A. : Heat, moisture and animal life in deserts - Proc. Roy.Soc., B.96, 123-131, 1924.
5. BUXTON, P.A. : The temperature of the surface of deserts - J. Ecol. 12, 127-134, 1924.
6. CLOUDSLEY-THOMPSON, J.L. : The significance of migration in myriapods - Ann.Mag. Nat. Hist. (12), 2, 947-62, 1949.
7. CLOUDSLEY-THOMPSON, J.L. : Economics of the "spotted snake-millipede", BLANIULUS GUTTULATUS (Bosc.) - Ann.Mag.Nat.Hist. (12), 3, 1047-57, 1950.
8. CLOUDSLEY-THOMPSON, J.L. : The water-relations and cuticle of PARADESMUS GRACILIS (Diplopoda: STRONGYLOSOMIDAE) - Quart.J. Micr.Sci., 91, 453-64, 1950.
9. CLOUDSLEY-THOMPSON, J.L. : On the responses to environmental stimuli and the sensory physiology of millipedes (Diplopoda) - Proc.Zool.Soc.Lond., 121, 253-77, 1951.
10. CLOUDSLEY-THOMPSON, J.L. : Studies in diurnal rhythms, 1.- Rhythmic behaviour in millipedes - J. Exp. Biol., 28, 165-72, 1951.
11. CLOUDSLEY-THOMPSON, J.L. : Studies in diurnal rhythms. II. Changes in the physiological responses of the woodlouse ONISCUS ASELLUS to environmental stimuli - J. Exp. Biol., 29, 295-303, 1952.
12. CLOUDSLEY-THOMPSON, J.L. : Diurnal rhythms - Trans. IX. Int.Congr.Ent., 1, 305-10, 1952.
13. CLOUDSLEY-THOMPSON, J.L. : Diurnal rhythm of locomotory activity in isolated migratory locusts - Ent. Mon. Mag., 89, 233-5, 1953.
14. CLOUDSLEY-THOMPSON, J.L. : The significance of fluctuating temperatures on the physiology of insects - Entomologist, 87, 153-61, 1953.
15. CLOUDSLEY-THOMPSON, J.L. : Diurnal rhythms in animals - Sci. News, 28, 77-98, 1953.
16. CLOUDSLEY-THOMPSON, J.L. : Studies in diurnal rhythms. III. Photoperiodism in the cockroach PERIPLANETA AMERICANA (L.) - Ann. Mag. Nat.Hist.(12), 6, 705-12, 1953.
17. CLOUDSLEY-THOMPSON, J.L. : Studies in diurnal rhythms. IV. Photoperiodism and geotaxis in TENEBRIO MOLITOR L. (Coleoptera: TENEBRIONIDAE) - Proc. R. Ent. Soc. Lond. A., 28, 117-32, 1953.
18. CLOUDSLEY-THOMPSON, J.L. : The ecological significance of diurnal rhythms in terrestrial arthropods - Sci. Prog., 42, 46-52, 1954.
19. CLOUDSLEY-THOMPSON, J.L. : Problems of dispersal in some terrestrial arthropods - Advanc. Sci., 11, 73-5, 1954.
20. CLOUDSLEY-THOMPSON, J.L. : Arthropods and terrestrial life - Sci. News, 36, 95-108, 1955.
21. CLOUDSLEY-THOMPSON, J.L. : The design of entomological aktograph apparatus - Entomologist, 88, 153-61, 1955.
22. CLOUDSLEY-THOMPSON, J.L. : The effect of rock cover on the diurnal range of microclimatic conditions - Entomologist, 89, 213-5, 1956.
23. CLOUDSLEY-THOMPSON, J.L. : The ecological significance of diurnal rhythms in terrestrial arthropods - Proc. XIV Int.Congr. Ent, 415-7, 1956.
24. CLOUDSLEY-THOMPSON, J.L. : Studies in diurnal rhythms. VI. Bioclimatic observations in Tunisia and their significance in relation to the physiology of the fauna, especially woodlice, centipedes, scorpions and beetles - Ann.Mag.Nat.Hist. (12), 9, 305-29, 1956.
25. CLOUDSLEY-THOMPSON, J.L. : Studies in diurnal rhythms VII. Humidity responses and nocturnal activity in woodlice (isopoda) - J.Exp. Biol, 33, 1956.


26. CLOUDSLEY-THOMPSON, J.L. : Studies in diurnal rhythms, V-Nocturnal ecology and water relations of the British cribellate spiders of the genus CINIFLO Bl.-J. Linn. Soc. (Zool.), 43, 134-52, 1957.
27. CLOUDSLEY-THOMPSON, J.L. : Spiders, scorpions, centipedes and mites. The ecology and natural history of woodlice, "myriapods" and arachnids, London, 1958.
28. CLOUDSLEY-THOMPSON, J.L. : Water relations and diurnal rhythms in woodlice - Ann.Appl.Biol., 46, 117-9, 1958.
29. CLOUDSLEY-THOMPSON, J.L. : The effect of wind on the nocturnal emergence of woodlice and other terrestrial arthropods I-III - Ent.Mon.Mag., 94, 106-8, 184-5, 283-4, 1958.
30. CLOUDSLEY-THOMPSON, J.L. : Studies in diurnal rhythms, VIII. The endogenous chronometer in GRILLUS CAMPESTRIS L. (Orthoptera: GRILLIDAE) - J.Insect Physiol., 2, 275-80, 1958.
31. CLOUDSLEY-THOMPSON, J.L. and SANKEY, J.H.P.: Some aspects of the fauna of the district around the Etang de Berre, Bouches-du-Rhône, France - Ann.Mag.Nat. Hist. (12), 10, 417-24, 1957.
32. EDNEY, E.B. : The evaporation of water from woodlice and the millipede GLOMERIS - J. Exp. Biol., 28, 91-115, 1951.
33. EDNEY, E.B. : Woodlice and the land habitat - Biol. Rev., 29, 185-219, 1954.
34. EDNEY, E.B. : The water relations of terrestrial arthropods, Cambridge, 1957.
35. FRAENKEL, G.S. and GUNN, D.L.: The orientation of animals. Kineses, taxes and compass reactions, Oxford, 1940.
36. GUNN, D.L. : The humidity reactions of the woodlouse PORCELLIO SCABER (Latreille)- J. Exp. Biol., 14, 178-86, 1937.
37. KACHKAROV, O.N. and KOROVINE, E.P.: La vie dans les deserts - Ed. française par Th. Monod , Paris, 1942.
38. PERTTUNEN, V. : Reactions of Diplopods to the relative humidity of the air. Investigations on ORTHOMORPHA GRACILIS, IULUS TERRESTRIS and SCHIZOPHYLLUM SABULOSUM - Ann.Soc.Zool.Fen. "Vanamo", 16 (4), 1-69, 1953.
39. RICHARDS, A.G. : The integument of arthropods - Minneapolis, 1951.
40. VACHON, M. : Etudes sur les scorpions - Alger, 1952.
41. VACHON, M. : The biology of scorpions - Endeavour, 12, 80-9, 1953.
42. WALOFF, N. : The mechanisms of humidity reactions of terrestrial arthropods - J. Exp. Biol., 18, 115.-35, 1941.
43. WIGGLESWORTH, V.B. : Principles of insect physiology, 4th Ed., London, 1950.
44. WILLIAMS, C.B. : Some bioclimatic observations in the Egyptian desert, in CLOUDSLEY-THOMPSON, J.L. (ed.): Biology of Deserts, London: Inst. Biol., 18-24, 1954.

S U M M A R Y

A review is given of the adaptations of arthropods to life on land. The terrestrial Arthropoda are divided ecologically into two main groups. The first of these includes woodlice, centipedes, millipedes and their allies which lose water rapidly in dry air and consequently are restricted to damp, dark habitats which they leave only at night when the temperature falls and the relative humidity of the air rises. The second group includes most insects and Arachnida: these are comparatively independent of moist surroundings because their integument possesses an impervious layer of wax which prevents desiccation.



1. Woodlice on a wall at night




2. The desert woodlouse (*HEMILEPISTUS REAUMURI*)
(much enlarged)



3. Apparatus for microclimatic measurement
Left, electrical thermometer employing thermistors;
right, electrical resistance hygrometer.



4. The wall spider (*AMAUROBIUS FEROX*, male)
(enlarged)



5. The desert scorpion
(*ANDROCTONUS AUSTRALIS*)




FIG. 6 Rate of water-loss in dry air at different temperatures, and corresponding saturation deficiencies, from a woodlouse (*Porcellio*), millipede (*Oxidus*), spider (*Lycosa*) and insect (*Pieris* larva). In the woodlouse and millipede the rate of water-loss is proportional to the saturation deficiency of the atmosphere, but in the spider and insect it is negligible below about 30° C., the critical temperature at which their epicuticular wax-layers become porous. Rate of water-loss is expressed in milligrams per square centimetre of surface area per hour.

Section C : Veterinary bioclimatology

Z O O L O G I C A L B I O C L I M A T O L O G Y
Section C: Veterinary Bioclimatology

CORRELATION OF BODY TEMPERATURE AND OF RESPIRATION RATE
OF SWINE WITH ENVIRONMENTAL TEMPERATURE

by

T.L. Noffsinger and F.N. Andrews (U.S.A.) *

A B S T R A C T

Correlation analyses were carried out to determine relation of body temperature and respiration rate of three breeds of swine to the environmental temperature. Coefficients of regression were determined for a prediction equation of the form:

$$y = a + b_1 x_1 + b_2 x_2 + b_3 x_3 + b_4 x_4 + b_5 x_5$$

where, y = body temperature

x_1 = respiration rate

x_2 = body weight

x_3 = sex

x_4 = environmental temperature

$x_5 = x_1 x_4$

I. I N T R O D U C T I O N

It is generally conceded that limited fluctuations in environmental temperature does not materially alter body temperature of warm-blooded animals (1, 2, 4). Such fluctuations activate thermoregulatory apparatus to counteract the external change in environment. Animals differ in their ability to dissipate heat due to their hereditary make-up, state of health and nutritional regime. Heat exchange in the live animal theoretically can be altered by change in body surface, change in color and amount of body covering, vasodilation or constriction, hormonal change influencing metabolic rate, increased vaporization through sweating and by acceleration of respiration rate.

It has generally been assumed that swine are affected more by summer temperatures than any of the other farm animals. The hog is a nonsweating animal and in addition, with increasing weight, the fatty covering is thought to present a barrier to the loss of body heat.

A review of the literature indicates there is insufficient knowledge of the fundamental physiology of swine and farm animals in general in regards to adapting to unfavorable environmental temperature to make a valid comparison of breeds.

The objects of this experiment were to:

- (1) determine the correlation of body temperature with respiration rate and environmental temperature in three different breeds of swine;
- (2) evaluate the coefficients of the regression equation for predicting body temperature of the three breeds - Duroc Jersey, Chester White and Yorkshire

II. E X P E R I M E N T A L P R O C E D U R E

This experiment was conducted during two summer periods - June through August 1955 and 1956 - on the Livestock Experimental Farm of Purdue University at Lafayette, Indiana. Data were obtained on

*) T.L. Noffsinger: Climatologist of the Land Study Bureau, Univ. of Hawaii, Honolulu 14, Hawaii.
F.N. Andrews : Professor of Animal Science, Purdue University, Lafayette, Indiana, U.S.A.
Statistical analyses by: Dr. M. Carter, Statistical Lab., Purdue Univ., Lafayette, Indiana, U.S.A.

body temperature and respiration rate for hogs of the three breeds over a range of environmental temperatures extending from 60° to 95° F (15.6° to 35° C).

Rectal temperatures were obtained on the resting hog by inserting a clinical fever thermometer to a depth of approximately 4 inches and reading to the nearest one-tenth degree Fahrenheit. Respiration rate was determined on the resting hog by a one minute visual count. The ambient temperature was then secured from the reading of a mercurial thermometer housed in a standard louvered instrument shelter located in the vicinity of the experimental pens. The thermometer was located approximately 5 feet above the ground. Pigs of each breed were subjected to identical environmental conditions.

During the 1955 trial, pigs had access to self-feeders and automatic waterers and were fed a ration of yellow shelled corn and protein supplement free choice. ¹⁾

The protein supplement was composed of: Soybean oil meal, 46.9%; meat and bone scraps, 20.0%; fish meal, 20.0%; alfalfa meal, 10.0%; salt, 2.0%; antibiotics 3.3 ²⁾, 0.85%; and fortafeed ³⁾ 0.25%.

Pigs used in the 1956 trials were self-fed with a complete mixed ration composed of: Ground yellow corn, 64%; ground oats, 15%; supplement V ⁴⁾, 19%; and mineral mixture ⁵⁾ 2%.

Individual weights, average feed consumption and feed efficiency were obtained at weekly intervals. The average weight of pigs of each breed increased from an initial weight of 75 lbs. to 200 lbs. at the end of the experiment.

Three breeds of hogs were represented in this work. There were 37 Duroc, 110 Chester White and 95 Yorkshire pigs in the experiment. During the period more than 400 individual readings were made to determine rectal temperature and respiration rate. These data were then analyzed to determine the relationship of rectal temperature to respiration rate, weight of pig, sex of pig and environmental temperature.

III. RESULTS

In the preliminary analysis of the data all combinations of pairs of the independent variables were plotted vs. rectal temperature to obtain an indication of the need for interaction terms in the regression equation.

This was done by plotting the points of one of the independent variables and rectal temperature for various values of the second independent variable. If these relationships between the first independent variable and rectal temperature changed depending on the value of the second independent variable, this was taken as an indication of the need for an interaction term.

Of the combinations of independent variable, only respiration rate and ambient temperature appeared to be important. In this case, the increase in rectal temperature associated with an increase in respiration rate was greater for high ambient temperatures than for low ambient temperatures.

By a regression analysis, coefficients were determined for an equation of the form:

$$y = a + b_1 x_1 + b_2 x_2 + b_3 x_3 + b_4 x_4 + b_5 x_5$$

where y = rectal temperature in degrees Fahrenheit

x_1 = respiration rate, respirations per minute

x_2 = body weight in pounds

x_3 = sex, where 1 = females and 2 = castrate males

x_4 = environmental temperature in degrees Fahrenheit

$x_5 = x_1 x_4$

- 1) Feed was available at all times and temperature and respiration data were collected only on those animals lying quietly in the shade. Individuals from all breeds were examined within a period of a few minutes. Data were collected between the hours of 8 a.m. and 4 p.m. local time.
- 2) Pfizer's 3.3 contained 3 grams of terramycin and 3 milligrams of Vitamin B12 per pound which supplied 50 grams of terramycin per ton and 50 milligrams of Vitamin B12 per ton.
- 3) Fortafeed (Lederle) vitamin supplement contained 2, 4 and 9 grams of riboflavin, calcium pantothenate and niacin, respectively, per pound of supplement.
- 4) Purdue Supplement V had the following composition: 40% soybean oil meal, 25% meat and bone scraps, 25% fish meal, 10% cottonseed meal.
- 5) Mineral mixture was composed of 40% ground limestone, 40% steamed bonemeal, 20% iodized salt.

Table I shows the simple correlation coefficients of each of the independent variables with rectal temperature for the Duroc, Chester White and Yorkshire breeds, respectively, plus values for all breeds combined.

Table 2 gives the square of the multiple correlation coefficients which indicate the relative amount of the variation of rectal temperature accounted for by the factor studied. For example, the value 0.7263 for Durocs indicates that approximately 73% of the variation of rectal temperature of the Durocs is accounted for by the change in respiration rate, weight, sex, ambient temperature and respiration rate X ambient temperature. The value 0.6864 indicates that approximately 69% of the variation of rectal temperature in the Durocs can be accounted for by variation in respiration rate and environmental temperature. A comparison of R^2 for Durocs with that for Yorkshire (see Table 4) shows a marked difference between breeds. Approximately 46% of the variation of rectal temperature in Yorkshire was accounted for by the five variables studied, but only 37% when x_5 (the interaction term) was removed. When respiration rate and environmental temperature were the only variables included in the regression equation, 34% of the rectal temperature variation was accounted for.

The tables of simple correlation coefficients (Tables 2, 3 and 4) show that respiration rate and rectal temperature of the Chester White and Duroc hogs are more highly correlated than are those variables in the Yorkshire hogs. Table 5 shows the regression coefficients and intercepts for the variables when data for all breeds were combined. The mean values of body temperature by breeds for the three temperature levels are shown in Table 6.

IV. DISCUSSION

Dukes (4) reporting on work by Palmer gives the average body temperature of the pigs as 102.5°F (39.2°C) with a range of 101.6° to 103.6° F (38.7° to 39.8° C). Brody (2) reports the temperature of the pigs as 103° F (39.4°C) and includes the pig along with cattle, sheep, goats, dogs and rabbits within a temperature range of 100° to 103° F (37.8° to 39.4° C).

Findlay (6) reviewing the findings of several investigators reports the range of rectal temperature for the pigs from 98.3° to 104° F (36.8° to 40.0°C). Heitman et al (7, 8) using a psychrometric room for temperature control studies on swine found that as the air increased from 40° to 115° F (4.4° to 46.1° C) the rectal temperature of the pig also increased. The heavier pigs showed a greater increase in rectal temperature with increasing environmental temperature.

Respiration rate in swine, as in other nonsweating species, rises rapidly with increasing environmental temperature. Brody (2) points out that in swine the respiration rate rises rapidly with increasing environmental temperature to compensate for the inability to lose heat by evaporation of sweat and to increase the vaporization rate from the respiratory passage. Dukes (4) reports the respiration rate of the resting mature hog at 8-18 rpm. Robinson and Lee (12) noted an increase in respiration when pigs were exposed to 75° and 80° F (23.9° and 26.7° C). At temperatures of 85°, 90° and 95° F (29.4°, 32.2° and 35.0° C) there was about a fourfold increase in the respiration rate.

On the basis of average body temperature at three levels of environmental temperature, Yorkshire pigs appeared to be no more sensitive to high environmental temperature than Durocs and Chester Whites. Since only 46% of the variation in body temperature of Yorkshire pigs was accounted for (see Table 4) by the variables studied, it appears that one or more important factors have been excluded. It had previously been noted that Yorkshire swine show a greater reduction in feed consumption in hot weather than do the other breeds. During a 56-day period, June 20 to August 15, Duroc, Chester White and Yorkshire pigs consumed an average of 5.50, 4.83 and 3.68 lbs. per day, respectively. It would appear that the Yorkshire's sensitivity to high temperature is mediated through reduced feed consumption, while the Duroc and Chester White pigs maintain high feed consumption but mainly relatively constant body temperature by decreased activity and increased respiration rate.

No endocrinology studies were made to explain the lower mean body temperature in an environment of 70° to 80° F (21.1° to 26.7° C) than in one of 60° to 70° F (15.6° to 21.1°C) although the data suggest an increased thyroid activity due to the lower environmental temperatures. The environmental temperature of 80° to 90° F (26.7° to 32.2° C) appears to be sufficiently high to obscure the reduced thyroid activity thus resulting in a higher body temperature.

V. REFERENCES

1. ANDREWS, F.N. : The Influence of the Environment on Reproduction in Female Farm Animals - Iowa State Col. J. Sci., 28, 9-18, 1953.
2. BRODY, SAMUEL : Bioenergetics and Growth - Reinhold Publ. Co., New York, 265-266, 1945.
3. BRODY, SAMUEL, and KIBLER, H.H.: Resting Energy Metabolism and Pulmonary Ventilation in Growing Swine - Res. Bul. Univ. of Missouri, 380, 26, 1948.
4. DUKES, H.H. : The Physiology of Domestic Animals - Comstock Publ. Co., Ithaca, New York, 473-474, 1948.
5. FOOTE, W.C., POPE, A.L., NICHOLS, R.E., and CASIDA, L.E.: The Effect of Variation in Ambient Temperature and Humidity on Rectal and Testis Temperature of Sheared and Unsheared Rams - J. Ani. Sci., 16, 1, 144-150, 1957.
6. FINDLAY, J.D. : The Effects of Temperature, Humidity, Air Movement and Solar Radiation on the Behavior and Physiology on Cattle and Other Farm Animals - Bul. no. 9, Hannah Dairy Res. Inst., 1950.
7. HEITMAN, H., and HUGHES, E.H.: The Effects of Air Temperature and Relative Humidity on the Physiological Well Being of Swine - J. Ani. Sci., 8, 171-182, 1949.
8. HEITMAN, H., HUGHES, E.H., and KELLY, C.F.: Effects of Elevated Ambient Temperature on Pregnant Sows - J. Ani. Sci., 10, 907-915, 1951.
9. IRVING, LAURENCE : Physiological Insulation of Swine as Bare Skinned Mammals - J. Applied Physiol., 9, 414, 1956.
10. LEE, D.H.K., and PHILLIPS, R.W.: Assessment of the Adaptability of Livestock to Climatic Stress - J. Ani. Sci., 7, 391, 1948.
11. McDOWELL, R.F., LEE, D.H.K., FOHRMAN, M.H., and ANDERSON, R.S.: Respiratory Activity as an Index to Heat Tolerance in Jersey and Sindi X Jersey (F_1) Cross-bred Cows - J. Ani. Sci., 12, 573-582, 1953.
12. ROBINSON and LEE : Cited by Findlay, 1950.

*
* * *

TABLE 1. SIMPLE CORRELATION COEFFICIENTS BETWEEN SELECTED VARIABLES ¹⁾

Variables	Duroc Jersey	Chester White	Yorkshire	All Breeds Combined
r_{01}	.83	.78	.54	.74
r_{02}	.24	-.17	-.28	-.07
r_{03}	-.10	-.09	.00	-.21
r_{04}	.44	.65	.52	.58
r_{05}	.83	.80	.57	.75

1) 1, respiration rate; 2, body weight; 3, sex; 4, environmental temperature; 5, $x_1 x_4$.

TABLE 2. REGRESSION COEFFICIENTS ¹⁾ (Duroc trial)

Explained Variation 2) R^2	Respiration b_1	Weight b_2	Sex b_3	Environmental Temperature b_4	Interaction b_5	Intercept a
0.7263	- 0.0347	0.0013	-0.5940	- 0.0275	0.0007	0.4536
0.7111	0.0217	0.0025	-0.6019	0.0229	-----	0.9937
0.7023	- 0.3504	-0.0100	-----	- 0.3534	0.0066	0.5330
0.7258	- 0.0366	-----	-0.5714	- 0.0246	0.0007	0.4475
0.7019	- 0.3347	-----	-----	- 0.3804	0.0065	0.5406
0.6864	0.2252	-----	-----	0.1689	-----	0.6778

- 1) Coefficients showing rate of change of body temperature with respiration rate b_1 , weight b_2 , sex b_3 , environmental temperature b_4 , and interaction b_5 .
 2) Percent of variation in body temperature accounted for by the variables shown.

TABLE 3. REGRESSION COEFFICIENTS ¹⁾ (Chester White trial)

Explained Variation 2) R ²	Respiration b ₁	Weight b ₂	Sex b ₃	Environmental Temperature b ₄	Interaction b ₅	Intercept ■
0.6611	- 0.3588	-0.0233	-0.9864	0.0137	0.0059	0.2253
0.6401	0.1310	-0.0363	-0.8607	0.3498	-----	-0.1536
0.6593	- 0.3532	-0.0225	-----	0.0456	0.0057	0.2113
0.6570	- 0.4096	-----	0.9177	-0.0103	0.0064	0.2329
0.6554	- 0.4029	-----	-----	-0.0007	0.0063	0.2196
0.6284	0.1353	-----	-----	0.3324	-----	-0.5418

- 1) Coefficients showing rate of change of body temperature with respiration rate b₁, weight b₂, sex b₃, environmental temperature b₄, and interaction b₅.
 2) Percent of variation in body temperature accounted for by the variables shown.

TABLE 4. REGRESSION COEFFICIENTS ¹⁾ (Yorkshire trial)

Explained Variation 2) R ²	Respiration b ₁	Weight b ₂	Sex b ₃	Environmental Temperature b ₄	Interaction b ₅	Intercept ■
0.4636	- 0.1173	-0.0046	-0.1346	-0.0035	0.0015	0.3377
0.3702	0.0085	-0.0090	-0.1838	0.0506	-----	-0.6634
0.4602	- 0.1189	-0.0039	-----	-0.0052	0.0015	0.3343
0.4564	- 0.1262	-----	-0.0912	-0.0066	0.0016	0.2982
0.4548	- 0.1263	-----	-----	-0.0075	0.0016	0.2999
0.3390	0.0923	-----	-----	0.5155	-----	-0.2101

- 1) Coefficients showing rate of change of body temperature with respiration rate b₁, weight b₂, sex b₃, environmental temperature b₄, and interaction b₅.
 2) Percent of variation in body temperature accounted for by the variables shown.

TABLE 5. REGRESSION COEFFICIENTS ¹⁾ (All Breeds Combined)

Explained Variation 2) R ²	Respiration b ₁	Weight b ₂	Sex b ₃	Environmental Temperature b ₄	Interaction b ₅	Intercept ■
0.6028	- 0.0399	-0.0030	-0.4242	0.0035	0.0007	0.2388
0.5823	0.0159	-0.0013	-0.3830	0.0390	-----	-0.5642
0.5785	- 0.3239	-0.0244	-----	0.0438	0.0058	0.1969
0.5988	- 0.0401	-----	-0.4115	-0.0006	0.0009	0.2292
0.5176	- 0.3268	-----	-----	0.0100	-----	0.1899
0.5600	0.1702	-----	-----	0.3275	-----	-0.7353

- 1) Coefficients showing rate of change of body temperature with respiration rate b₁, weight b₂, sex b₃, environmental temperature b₄, and interaction b₅.
 2) Percent of variation in body temperature accounted for by the variables shown.

TABLE 6. BODY TEMPERATURE

Environmental Temperature	Breed	Number Measurements	Body Temperature	
			Degrees Fahrenheit	Degrees Centigrade
60°-70° F (15.6°-21.1° C)	Duroc	86	104.1	40.0
	Chester White	63	103.3	39.6
	Yorkshire	69	102.6	39.2
70°-80° F (21.1°-26.7° C)	Duroc	68	103.3	39.6
	Chester White	25	102.7	39.3
	Yorkshire	36	102.1	38.9
85°-95° F (29.4°-35.0° C)	Duroc	81	104.2	40.1
	Chester White	25	104.3	40.2
	Yorkshire	13	103.9	39.9

Z O O L O G I C A L B I O C L I M A T O L O G Y
Section C: Veterinary bioclimatology

THE INFLUENCE OF CLIMATIC FACTORS ON THE HEAT BALANCE AND
MILK YIELD OF THE COW IN RELATION TO THE DESIGN OF FARM
BUILDINGS IN THE NETHERLANDS

by

Dr. C.C. Oosterlee (Netherlands) *

I. INTRODUCTION

During the war many farm buildings in the Netherlands were destroyed and since farms had also to be built in the new land won from the sea (the polders) the need arose for an investigation on the influence of climatic factors on the heat balance and milk yield of the cow so that such new buildings could be efficiently designed for optimum milk production.

In 1954 the author was asked to make such an investigation on behalf of the Organisation for Applied Scientific Research (T.N.O.) and this paper deals with the methods used and the results obtained.

II. THE EFFECTS OF CLIMATE ON THE COW

The general effects of climate on cattle have been reviewed by Findlay and Beakley (2) who showed, in general, that cows are better adapted to low than to high temperatures. Worstell and Brody (7) have shown that the comfort zone for cattle lies between 0° and 15.5°C. Two reasons can be adduced for the ill-adaption of cattle to hot environments. First, the high level of heat production, e.g. 10,000 Kcal/day for a non-lactating, non-gestating cow or 20,000 Kcal/day for a cow yielding 20 kg milk per day. Second, cows are a poorly-sweating species of mammalia, skin moisture vaporisation attaining a maximum at only 17° C ambient temperature (4).

III. METHODS

Two experimental methods are available for studying the influence of climate on the milk yield during winter months. The first is to identify the climate within barns and ascertain their influence on the cow and its production. The second method is more indirect and consists of correlating periods of low temperature e.g. during frost, with the amount of milk sent to the factories.

Fig. 1 and 2 show the relation between the supply of milk to the factories for the whole country and the mean weekly temperature. During some years there was an indication, that temperatures below zero influenced the supply of milk. Other factors, for instance the time of calving, especially at the beginning of February, had much more influence, but in relation to other years the period of February 1956, associated with very low temperatures, had a significant effect on the supply of milk. The supply of milk during the first week in February with temperatures below zero was 3.7% too low in relation to the supply of milk during January 1955 and the supply of milk during the last week of January which did not have low temperatures.

This 3.7% fall indicates that during the first week of February 1956, the supply of milk was 2,284,000 kg/week too low, or about 0.5 kg/cow/day.

It is not known whether this effect was due to climatic factors (low temperatures in the stable), or to changes in the rations (the latter is the case in the Netherlands, if the silage is frozen and cannot be given to the cows).

* Afd. Veeteelt, Duivendaal 3a, Landbouw Hogeschool (Agricultural University), Wageningen, Netherlands.

This method can therefore give only an indication as opposed to directly measuring the effect of climate.

The non-evaporative heat loss was measured with a Hatfield J.L.C. heat flow disc (low temperature type) (6). To ascertain the influence of temperature and humidity on the milk yield, the cows in the barns were divided into two groups, one group with a high level, the other with a low level of production. Using the table for "standard cow production", the yield per cow was normalized on age and stage of lactation.

If the groups are big enough they are comparable. The table for standard cow production is based on the fact, that the highest yield of a cow in the Netherlands is generally reached at 8 yrs and 2 mths. This is taken as 100%. Each part of the lactation curve of a cow can be taken as a percentage of the top level of production. Doeksen and Heyboer (1) drew up this table in 1954 using hundreds of Dutch-Friesian cows. The basis of this method was that the influence of climatic factors on a cow with a high level of production, and therefore with a much higher heat production and heat loss, is not the same as that on a cow with a low level of production. In barns with, for instance, a high temperature, the cow with a high level of production is more likely to lose its heat balance than a cow with a low level of production. The depressing effect of high temperatures will be greater on the cow with the high level of production. Because of the normalisation tables, changes in milk production, due, for instance, to the quality of the food, will be equal in the two groups. Influences due to climatic factors would tend to depress the production of the group with a high level of production and not the one with a low level of production.

To check the accuracy of the heat flow disc, it was tested in a respiration (climatic) chamber of the Laboratory of Animal Physiology of the Agricultural College at Wageningen.

The heat loss to the ground was measured using a plastic sac filled with water at a constant temperature of 30° C. The heat flow to the ground beneath the sac was measured with the disc.

IV. THE MEASUREMENTS WITH THE HEAT FLOW METER

Using the heat flow disc the non-evaporative cooling can be measured at several environmental temperatures, so ascertaining at which temperature the cow is in thermo-neutrality. The basis of these measurements may be seen in Fig. 3 based on the measurements of Kibler and Brody 1950. The non-evaporative cooling is zero when the skin temperatures equal environmental temperatures. Decreasing the environmental temperature increases the non-evaporative heat loss. Below a certain temperature increase in non-evaporative heat loss cannot continue unless the body temperature decreases. Consequently the non-evaporative heat loss is limited (point P). This point represents thermo-neutrality since heat production should be equal or nearly equal to heat loss.

By measuring the non-evaporative cooling of the trunk, it is possible to ascertain the temperature corresponding to point P.

Before this was done the heat flow meter was tested. The non-evaporative cooling of the surface of the pelvis between the tuber coxae and the tuber ischiadicum, used as the place for routine measurements, was compared with the heat loss of other places on the surface of the trunk. No significant differences were found.

The non-evaporative cooling was then measured at several temperatures on more than 60 cows. It was found that the non-evaporative cooling is limited to about 8° - 9° C. Temperatures below zero did not increase the non-evaporative cooling but low temperatures and wind did. Figs. 4 and 5 show these results.

In the respiration chamber the sun, of the non-evaporative cooling measured with the heat flow meter and corrected for total surface, and the evaporative cooling, condensed water, were measured at various temperatures and were found to be comparable to the heat production as shown in Table 1.

The milk yield of cows housed in the open shed was unaffected during periods of very low temperatures and the animals were apparently healthy although temperature gradients between body and environment were about 50° C. In an attempt to objectively ascertain the comfort of the cow, living in an environment with temperatures of - 10° C, the following experiment was carried out. During a very severe frost period (Febr. 1956), an open shed was divided into three parts and 5 heating elements switched on successively above each part. Every 15 min. it was noted which cows preferred the warmth of the heating elements. It was found that when the air temperature was about -10° C the cows with the lowest heat production i.e. heat loss in the zone of thermo-neutrality preferred the heat of the elements significantly more than cows with higher heat production. When the conditions were particularly severe, e.g. low temperatures and wind, it was found that the "boss cows" scared away the other animals.

V. HEAT BALANCE AND THE FUNCTION OF THE SKIN, THE COAT AND CONDUCTION TO THE GROUND

Although on many farms the cows are clipped in the autumn, on others this does not occur. To obtain information about the insulating properties of the coat in relation to the heat balance of the animals, some measurements with the heat flow meter were carried out.

The non-evaporative cooling of six cows with a "normal" coat (end of November) was measured with the heat flow meter. The procedure was repeated after clipping a small part of the coat and after complete clipping of the animal. The results are given in Table 2. The increase of the non-evaporative cooling after clipping a very small part of the coat of 6 cows averaged 18.2 kcal/m²/hour. After complete clipping of the animals the non-avaporative cooling returned to the value obtained before clipping. In terms of non-evaporative cooling/cow/kcal/m²/hour, the animal with the lowest heat loss had the best insulation.

The return of the non-evaporative cooling, after complete clipping of the animals to the normal value indicates, that the animals can regulate the non-evaporative cooling adequately by vasoconstriction.

Non-evaporative cooling was also found to drop about 20 kcal/m²/hour after injection of 4 ml adrenaline (1 : 1000, 32° C), subcutaneously, under the disc. The temperature of the environment was about 12° C. Two animals were used.

During the experiments in the respiration chamber it was found that the non-evaporative cooling was higher when an animal was lying than when standing. This unexpected result was found to be due to the very good insulation of the floor of the respiration chamber. Measurements have proved that a lying animal does not lose enough heat if the ground beneath is well-insulated. This is compensated for by an increase in the non-evaporative cooling through the skin to the air.

This finding suggests that in a warm climate the insulation of the ground beneath should be poor and in a cold environment the ground should be well-insulated. Measurements with the heat flow meter showed that if a small layer of straw is put on the floor of a conventional barn, this can reduce the heatflow to the ground beneath from 150 kcal/m²/hr to 40 kcal/m²/hr.

VI. STABLE CLIMATE AND ITS INFLUENCE ON MILK PRODUCTION

To collect information about the role the climatic conditions play in the Dutch barns, the temperature and humidity were measured twice a day, in about 12 stables during the winter of 1954-1955 and 1955-1956.

It was found, that especially in spring, though also in autumn, "tropical" conditions are not uncommon. This can be seen in Fig. 6. These high temperatures, with corresponding high humidities, have a depressing effect on the milk yield of a cow with a high level of production (20 kg milk/day or more). As previously stated, this was found with the help of the tables for "standard cow production". Before these tables were used, they were tested. Fig. 7 assesses the value of the tables. Two groups of cows, one with a high level of production and one with a low level of production on an ordinary Friesian farm were used. It can be seen, that during the time the animals were in the pasture the difference between the two groups was constant. Only during the time the animals were in the stable there was a difference. It can also be seen, that during pasture time changes in the ration are nearly equal for both groups.

In the stables where the temperatures were measured comparisons between the standard cow production of groups with a high level of production and groups with a low level of production were made. It was found that the production of the group with a high level of production decreased when the animals were exposed to temperatures higher than 16-17° C. Higher temperatures with high humidities adversely affected the persistency of the lactation curve of the cows.

The lower temperatures during the period with frost in February 1956 had a depressing effect on the milk yield of all cows in the open shed and in some conventional barns during the first two days. The milk yield of the cows in the open shed did not further decline, but in two closed barns, where the animals were able to move freely, the milk production decreased further during the following days when the temperatures were much higher than in the open sheds. There was a marked correlation between the fluctuations in milk production of the cows in these two independent groups ($r = \pm 0.69$). It was found that the temperatures obtaining during the previous 5 days were the cause of these fluctuations. Lee, McMillan, McDowell and Fohrman (5) found the same delayed effect under summer conditions in the U.S.A. 't Hart and Kemp (3) also found a delay in the influence of the temperature of the previous 5 days when there was an outbreak of grass tetany in the Netherlands.

In conventional barns there is an indication that the milk yield of the cows with a low level

of production decreases during periods of frost.

The difference between the decrease of the milk yield of the cows in the two barns which were closed and where the animals were free to move and that of the undisturbed production of the cows in the open shed with much lower air temperatures is striking. The direct solar radiation on the cows in the open shed may be an important factor. During the experiment using heating elements it was noted that during the day the cows stood almost all day in the sun.

VII. SUMMARY

The results and methods of an investigation on the influence of climatic factors on the heat-balance and milk yield of the cow are described. The non-evaporative cooling was measured at several environmental temperatures with use of a heat flowmeter. The observation was made, that to about 8-9° C the increase of the non-evaporative cooling of the trunk (at decreasing environmental temperatures) was limited.

In spring and autumn the climatical conditions in the stables in the Netherlands were in some cases "tropical". These high temperatures, with corresponding high humidities, have a depressing effect on the milk yield of a cow with a high level of production (20 kg milk/day or more).

VIII. REFERENCES

1. DOEKSEN, J. and HEYBOER, S.E.: Het omrekenen van melkvee tot standaardkoeien - Verslagen van Landbouwkundige onderzoekingen, 58,7, 1952.
2. FINDLAY, J.D. and BEAKLEY, W.R.: Environmental physiology of farm animals - Progress in the physiology of farm animals, 1, 252-298, 1954.
3. HART, M.L.'t and KEMP, A. : De invloed van de weersomstandigheden op het optreden van kopziekte bij runderen - Tijdschrift v. Diergeneesk. 81,84-95, 1956.
4. KIBLER, H.H. and BRODY, S. : Influence of temperature 5-95° F on evaporative cooling from the respiratory and exterior body surfaces in Jersey and Holstein cows - Mo.Agr.Exp.Sta.Res.Bull., 461, 1950.
5. LEE, D.H.K., McMILLAN, H.W., McDOWELL, R.E. and FOHRMANN, M.H.J.: Temperature conditions critical to milk production in Holstein cattle at Beltsville - J. Anim. Sci., 13, 1024, 1954.
6. STAFFORD, H. : A heat flowmeter - Proc. Physiol.Soc.J. Physiol., 111, 16-19, 1949.
7. WORSTELL, D.M. and BRODY, S. : Comparative physiological reactions of european and indian cattle to changing temperatures - Mo.Agr.Exp.Sta. Res. Bull., 515, 1953.

Table 1. Comparison of the heatproduction with the evaporative cooling and the non-evaporative cooling (part of the skin between the tuber coxae and the tuber ischiadicum).

Chamber temp. °C	Rel. humidity %	Heatproduction Cal/hr		Evapera- tive cooling Cal/hr	Non-eva- porative cooling Cal/hr	Total heatloss Cal/hr	Body temp.	
		standing and lying	standing				10 hr	17 hr
15	74	804	844.2	232.2	587.7	819.9	38.6	38.2
11	81	797	836.8	186.1	633.7	819.8	38.5	38.8
5	79	772	810.6*	200.1**	648.5	848.6	38.8	38.8
25	75	780	819	427.4	387.2	814.6	38.8	38.8

* The lowest heat production at 5°C is difficult to understand but, as during the first experiments the animal was not completely accustomed to the chamber, it was rather restless and this explains the slightly higher heat production.

** The measured evaporative cooling (condensed water) is too high. This is due to the fact that the temperature of 5°C was the limite of cooling capacity of the chamber; this caused a certain error.

Table 2. Influence of the coat on the non-evaporative cooling of six Dutch - Friesian cows. The measurements were carried out at a stable temperature of 12.6°C. The total surface is calculated from the weight the formule $0 = 0.15 G^{0.56}$

Number of cow	Non-evaporative cooling before shaving		Non-evaporative cooling after shaving 15 cm2 of the coat		Increase of non-evaporative cooling		Non-evaporative cooling after complete shaving		Increase of non-evaporative cooling	
	Kcal/hr		Kcal/hr		Kcal/hr		Kcal/hr		Kcal/hr	
Identical twin:	per m2	Total surface	per m2	Total surface	per m2	Total surface	per m2	Total surface	per m2	Total surface
8714	109.4	521.8	131.1	625.3	21.7	103.5	110.3	526.8	0.9	4.0
53493	113.6	586.1	129.5	668.2	15.9	82.1	118.4	610.0	4.8	24.9
S. 321	104.7	513.0	128.2	628.1	23.5	115.1	104.3	512.0	-0.4	-0.9
19	112.4	551.9	126.2	619.6	13.8	67.7	118.1	579.8	5.7	27.9
20	110.2	556.5	127.1	641.8	16.9	85.3	115.5	583.2	5.3	26.7
4994	109.4	547.0	126.7	633.5	17.3	86.5	113.6	568.0	4.2	21.0
Average	109.9	546.0	128.1	636.0	18.2	90.0	113.3	563.2	3.4	17.2




Fig. 1. Relation between the supply of milk to the factories for the whole country and the average weekly temperature during the winters 1948-1949 up to 1955-1956.




Fig. 2. Relation between the supply of milk to the factories for the whole country and the average weekly temperature during the winters 1953-1954 up to 1956-1957.




Fig. 3. The relationship between surface temperature and non-evaporative cooling, theoretically to be expected at different environmental temperatures.
 Q.P.R. = non-evaporative cooling
 Q¹.P¹.R¹ = skin temperatures
 XY = relation between environmental temperatures and skin temperatures.




Fig. 4. The non-evaporative cooling (surface of the skin between tuber coxae and tuber ischiadicum) of 20 M.R.IJ. cows housed in an open shed at Leersum and 16 Dutch-Friesian cows in a loose house at Wageningen.




Fig. 5. The non- evaporative cooling (surface of the skin between tuber coxae and tuber ischiadicum) of 13 M.R.I.J. cows in an open shed at Diessen.




Fig. 6. Climograph for cattle in Europe and Asia (modified, Wright 1954). The climograph for a stable at Rijperkerk is drawn in the figure (hatched). The points represent the highest temperatures, with the corresponding humidities, found in the stables during the period of the investigation. The lines with the weak curves represent the limits of the comfort zone of men C, and of cows (Kibler and Brody 1953): A and B.
1. Lhasa, Tibet: Yak; 2. Delhi, India: Large Indian Zebu; 3. Cochin, South-India: Dwarf Indian Zebu; 4. Nuwera Eliya, Ceylon: imported European cattle. Hatched: stable climate in a Dutch farm.




Fig. 7. The standard cow production of a group cows with a high level production, a group with a low level production and the difference between the groups, during some months of 1950 for a Friesian farm. Two groups of 9 cows were used. The real production of the group with a high level production averaged at the beginning of March 24.3 kg milk/day, the group with a low level 12.4 kg milk/day. The broken line indicates the time the cows went into the pasture..

Section D : World literature

ZOOLOGICAL BIOCLIMATOLOGY

Section D: World literature*

GENERAL ZOOLOGICAL BIOCLIMATOLOGY

1. ADAMS, L. : An analysis of a population of snowshoe hares in northwestern Montana - Ecol. Monogr., 29, 141-170, 1959.
2. BROOKS, F.A. : Zoological Bioclimatology - (See Areal report Dr. D.H.K. Lee. Symposium Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956).
3. CLOUDSLEY-THOMPSON, J.L.: The relationship between reproductive rate and environmental resistance in British birds - Ann.Mag.Nat.Hist., 12, 5, 989-1019, 1952.
4. CLOUDSLEY-THOMPSON, J.L.: Maintenance activities in the behaviour of animals - The School Science review, 37, 134, 1956.
5. DAUBERT, K. and PETERS, H.: Studie über biosphär. Einflüsse auf die Abweichungen von Cichliden (Teleostei) - Zoolog. Anzeiger 1956 (In press).
6. DELAMARE DEBOUTEVILLE, Cl.: Microfaune du sol dans les pays tempérés et tropicaux - Actuelles Sc. et Industrielles, Hermann et Cie, Paris, 1060, 1-360, 1951.
7. DELAMARE DEBOUTEVILLE, Cl. and PAVLIAN, R.: Faune des nids et des Terriers en Basse Côte-s'Isoire - Encycl. Biog. et écol., 8, 1-116, 139 figs., 1952.
8. DESMARAIS, A. : Effet de l'ablation partielle du foie sur la résistance au froid d'animaux adaptés - Revue Canadienne de Biologie, 2, 453, 1943.
9. FOLK, G.E. and SCHELLINGER, R.R.: The diurnal rhythm of body temperature of the hamster - Anat. Rec., 120, 787, 1954.
10. FOLK, G.E. : Modification by light and feeding of the 24-hour rhythm of activity in rodents - Proc. Fifth.Conf. on Biol. Rhythms, Stockholm, 1955.
11. FOLK, G.E., GRINDELAND, R. and FARRAND, R.L.: Factors influencing the life span of hamsters - Proc. Iowa Acad. Sci., 64, 1957.
12. FREGLY, M.J. : Zoological Bioclimatology - (See Areal report Dr. D.H.K. Lee. Symposium Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956).
13. FRY, F.E.J. : The summer migration of the cisco. *Leucichthys artedii* (Le Sueur) in Lake Nipissing, Ontario - U.of T. Biol.; Ser. No. 44, Ont.Fish. Res. Lab. Pubn., 55, 1-91, 1937.
14. FRY, F.E.J. : The position of fish and other higher animals in the economy of lakes - Am. Assoc. Advancement Sci. Pubn., 10, 132-142, 1939.
15. FRY, F.E.J. : Movements of drift cards in Georgian Bay in 1953 - J. Fish. Res. Bd. Canada, 13, 1-5, 1956.
16. FRY, F.E.J. a.o. : Yield of the year classes of smallmouth bass of the decade of 1940 in South Bay Manitoulin Island - Trans.Am.Fish.Soc, 85, 1956.
17. HOCK, R.J. : Alaskan zoogeography and Alaskan amphibians - Science in Alaska, 1953 (in press)
18. HOCK, R.J. : The annual cycle of Alaskan ground squirrels - Science in Alaska, 1954 (in press).
19. HUNGERFORD, K.E. : Ruffed Grouse Populations and Cover Use - Trans.N.American Wildlife Conf., 16, 216-224.
20. HUNGERFORD, K.E. : The Ecology and Management of the Idaho Ruffed Grouse - Doctoral Dissertation, University of Michigan, published in Microfilm Edition, 1951.
21. HUNGERFORD, K.E. : Dew and Game - Paper presented in a symposium at the Ecological Society Meeting jointly with AAAS, December 1954.
22. IRVING, L. and PANEAK, S.: Biological reconnaissance along the Ahlasuruk River, east of Howard Pass, Brooks Range, Alaska, with notes on the Afifauna - J. of the Washington Academy of Sciences, 44, 7, 1954.
23. JUSATZ, H.J. : Klimatologische Beachtungen an Kaninchen (Aussprache-Bemerkung Mit-telrhein. Studienges. Bad Nauheim, II.1.1933) - Zschr.f.d.ges. phy-sikal. Ther., 45, 157-158, 1933.
24. LACK, D. : The natural regulation of animal numbers - Oxford University Press, 18, 343, 1954.

* References received from members by Secretariat of I.S.B.B. since 1957.

25. LANDSBERG, H.E. : Bird migration and Pressure patterns - Science, 108 (2817), 708-709, 1948.
26. McFARLANE, W.V. : The life cycle of STEGODEXAMENE ANGUILLAE n.sp., an Allocreadiid trematode from N.Z. - Parasitology, 41, 1-10, 1951.
27. McFARLANE, W.V. : Bionomics of two trematode parasites of N.Z. eels - J.Parasitol., 38, 391, 1952.
28. McFARLANE, W.V. : Experimental approaches to the functions of tropical livestock - Unesco Symposium on Microclimatology, 1956.
29. MOREAU, R.E. : A contribution to tropical African bird-ecology - J. Anim. Ecol. 3, 41-69, 1934.
30. MOREAU, R.E. : Breeding seasons of birds in East African evergreen forests - Proc. Zool. Soc. London, 631-653, 1936.
31. MOREAU, R.E. : The breeding seasons of African birds - Ibis, 92, 223-267, 1950.
32. MORRISON, P.R. and TIETZ, W.J.: Observations on food consumption in four Alaskan mammals - Arctic, 6, 52-57, 1953.
33. MORRISON, P.R. : Zoological Bioclimatology - (See Areal report Dr. D.H.K. Lee. Symposium Int. Soc. Bioclim. and Biomet., Paris 29-31 August, 1956).
34. NEVENS, W.B. : A comparison of sampling procedures in making pasture yield determinations - Journ. Dairy Sci., 28, 171-185, 1945.
35. PRUITT Jr., W.O. : An analysis of some Physical factors affecting the local distribution of the shorttail shrew (*Blarina brevicauda*) in the northern part of the lower Peninsula of Michigan - Ann Arbor University of Michigan, July 1953.
36. PRUITT Jr., W.O. : Notes on the Shorttail Shrew (*Blarina brevicauda kirtlandi*) in Northern Lower Michigan - The Am. Midland Naturalist, 52, 1, 236-241, 1954.
37. PRYCHODKO, W. : Über die Verkehropfer freilebender Tiere auf den Autobahnen im Südosten von Michigan - Säugetierkundliche Mitteilungen, 5, 4, 151-156, 1957.
38. SAPIN-JALOUSTRE : Biogéographie antarctique - Revue économique française, 67, 3, 2-6.
39. SCHEIN, M.W. : Zoological Bioclimatology - (See Areal report Dr. D.H.K. Lee. Symposium Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956).
40. SCHMIDT-NIELSEN, K. : Zoological Bioclimatology - (See Areal report Dr. D.H.K. Lee. Symposium Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956).
41. SCHUA, L. : Die Fluchtreaktion von Goldhamstern aus elektrischen Feldern - Naturwissenschaften, 40, 514, 1953.
42. SCHUA, L. : Die Wirkung von luftelektrischen Feldern auf Tiere - Verhandl. des Dtsch. Zool. Ges. in Tübingen 1954, 435, 1954.
43. SCHUA, L. : Die Wirkung luftelektrischer Felder auf Lebewesen - Umschau, 15, 468, 1954.
44. SHALEM, N. : L'influence de la rosée et des brouillards sur la répartition des escargots - Journal de Conchyliologie, 89, 95-107, 1949.
45. SIDOROWICZ, J. : Geographical variation of the squirrel *SCIURUS VULGARIS* L. in Poland - Acta Theriologica, 2, 7, 141-157, 1958. (In English).
46. SOLOMON, M.E. : The natural control of animal populations - J. Anim. Ecol., 18 (1), 1-35, 1949.
47. TAYLOR, W.E.A. and MALEWSKI, A.M.: The factor of topographical cues in bird homing experiments - Ecology, 36, 4, 1955.
48. UNGEHEUER, H. and JÖCHLE, W.: Psychisches Verhalten von Ratten unter extremen Haltungsbedingungen in Abhängigkeit von Wettereinflüssen - Ztschr. für Naturforschung, 12, 1, 1957.
49. WILLIAMS, C.B., FISHER, R.A. and CORBET, A.S.: The relation between the number of species and the number of individuals in a random sample of an animal population (Part. 2 Results obtained by means of a light trap at Rothamsted. pp 44-53 by C.B.W.) - Journ. anim.Ecol., 12, 42-58, 1943.
50. ZAWIDZKA, E. : Geographical distribution of the dark phase of the squirrel *SCIURUS VULGARIS FUSCOATER ALTUM* in Poland - Acta Ther., 2, 8, 159-174, 1958 (In English).

PHYSIOLOGICAL ZOOLOGICAL BIOCLIMATOLOGY

1. ASSENMACHER, I. : Evolution de la neurosécrétion hypothalamo-neurohypophysaire au cours de l'organogénèse du Canard domestique - Séances de la Société de Biologie, séance 13 July 1957, 151, 7, 1301, 1957.
2. ASSENMACHER, I. : Répercussions de lésions hypothalamiques sur le conditionnement génital du Canard domestique - Séances de l'Académie des Sciences, 245, 210-213, 1957.

3. ASSENMACHER, I. : Nouvelles données sur le rôle de l'hypothalamus dans les régulations hypophysaires gonadotropes chez le Canard domestique - Séances de l'Acad. des Sciences, 245, 2388-2390, 1957.
4. ASSENMACHER, I. and BENOIT, J. : Quelques aspects du contrôle hypothalamique de la fonction gonadotrope de la préhypophyse - Pathophysiologia Diencephalica, Symp. Int. Milan, May 1956, Wien, 402-427, 1958.
5. BEHMANN, F.W. : Die Regulierung der Wärmeproduktion bei Auskühlung homoiothermer Tiere - Arch. exp. Path. u. Pharmacol., 228, 126, 1956.
6. BONTKE, E. and BEHMANN, F.W. : Steuerbare Hypothermie durch intravasale KÜhlung beim Hund - Ber. ges. Physiol. u. Pharmacol., 172, 124, 1955.
7. BENOIT, J. : Activation sexuelle obtenue chez la Canard par l'éclairement artificiel pendant la période de repos génital - C.R. Ac.Sc., 199, 1671, 1935.
8. BENOIT, J. : Stimulation du développement testiculaire par l'éclairement artificiel - C.R.S.B., 118, 664, 1935.
9. BENOIT, J. : Rôle des yeux dans l'action stimulante de la lumière sur le développement testiculaire chez le Canard - C.R.S.B., 118, 669, 1935.
10. BENOIT, J. : Rôle de l'hypophyse dans l'action stimulante de la lumière sur le développement testiculaire chez le Canard - C.R.S.B., 118, 672, 1935.
11. BENOIT, J. : Nouvelles expériences relatives à la stimulation par la lumière du développement testiculaire chez le Canard - S.R. Ac.Sc., 201, 359, 1935.
12. BENOIT, J. : Influence de la lumière naturelle sur la croissance testiculaire chez le Canard au cours de la reprise sexuelle saisonnière - C.R.S.B., 120, 131, 1935.
13. BENOIT, J. : Stimulation par la lumière artificielle du développement testiculaire chez des Canards aveuglés par section du nerf optique - C.R.S.B., 120, 133, 1935.
14. BENOIT, J. : Stimulation par la lumière artificielle du développement testiculaire chez des Canards aveuglés par énucléation des globes oculaires - C.R.S.B., 120, 136, 1935.
15. BENOIT, J. : Maturité sexuelle et ponte obtenues chez la Cane domestique par l'éclairement artificiel - C.R.S.B., 120, 905, 1935.
16. BENOIT, J. : Sur la croissance du testicule du Canard immature déclenchée par l'éclairement artificiel. Etude histologique - C.R.S.B., 120, 1323, 1935.
17. BENOIT, J. : Hypophysectomie et éclairage artificiel chez le Canard mâle - C.R.S.B., 120, 1326, 1935.
18. BENOIT, J. : Activation du fonctionnement hypophysaire par des radiations lumineuses chez le Canard domestique - Revue de Physiothérapie, 12e année, 2, 86-94, 1936.
19. BENOIT, J. : Rôle de la préhypophyse dans le conditionnement de l'activité génitale du Canard, démontré par l'hypophysectomie - Arch. Portugaises des Sc. Biol., 5, 279-287, 1936.
20. BENOIT, J. : Activation par l'éclairement artificiel des glandes génitales des deux sexes chez le Canard domestique au repos sexuel - Bull. de la Soc. de Sexologie, 3, 438-455, 1936.
21. BENOIT, J. : Rôle de la thyroïde dans la gonadostimulation par la lumière artificielle chez le Canard domestique - C.R.S.B., 123, 243, 1936.
22. BENOIT, J. : Sur le caractère quantitatif de la réponse à l'excitation par la lumière artificielle du mécanisme gonadostimulant chez le Canard domestique - C.R.S.B., 123, 241, 1936.
23. BENOIT, J. : Stimulation of the hypophysis and genital glands in the duck by electric light. Effect of thyroidectomy on the testis and liver - Anat. Rec., 67, suppl. no. 1, 81, 1936.
24. BENOIT, J. : Relation between thyroid and growth of testes and penis when stimulated by electric light - Proc. Soc. Exp. Biol. & Med., 36, 782-784, 1937.
25. BENOIT, J. : Facteurs externes et internes de l'activité sexuelle. II. Etude du mécanisme de la stimulation par la lumière de l'activité testiculaire chez le Canard domestique. Rôle de l'hypophyse - Bull. Biol. de la France et de la Belgique, 71, 394-437, 1937.
26. BENOIT, J. : Action des facteurs externes sur l'hypophyse et les glandes génitales chez les Oiseaux - Colloque international au Collège de France du 10 au 19 juin 1937, 255-264. Fondation Singer Polignac Herman et Cie., Paris.
27. BENOIT, J. and OTT, L. : Action de lumières de différentes longueurs d'onde sur la gonadostimulation, chez le Canard mâle impubère - C.R.S.B., 127, 906, 1937.

28. BENOIT, J. : Action de divers éclairagements localisés dans la région orbitaire sur la gonadostimulation chez le Canard mâle impubère. Croissance testiculaire provoquée par l'éclaircement indirect de la région hypophysaire - C.R.S.B., 127, 909, 1938.
29. BENOIT, J. Rôle des yeux et de la voie nerveuse oculo-hypophysaire dans la gonadostimulation par la lumière artificielle chez le Canard domestique - C.R.S.B., 129, 231, 1938.
30. BENOIT, J. : Nouvelles recherches sur les voies nerveuses photoréceptrices et hypophysostimulantes chez le Canard domestique - C.R.S.B., 131, 89, 1939.
31. BENOIT, J. and OTT, L. : External and internal factors in sexual activity. Effect of irradiation with different wavelengths on the mechanisms of photostimulation of the hypophysis and on testicular growth in the immature duck - Yale J. of Biol. and Medicine, 17, 22-46, 1944.
32. BENOIT, J. : Reproduction. Caractères sexuels et hormones. Déterminisme du cycle sexuel saisonnier - Traité de Zoologie, P. GRASSE, 15, 384-478, 1950.
33. BENOIT, J., ASSENMACHER, I. and WALTER, F.X. : Réponses du mécanisme gonadostimulant à l'éclaircement artificiel et de la préhypophyse aux castrations bilatérales et unilatérales, chez le Canard domestique mâle, au cours de la période de régression testiculaire saisonnière - C.R.S.B., 144, 573, 1950.
34. BENOIT, J., ASSENMACHER, I., GROS, C. and WALTER, F.X. : La radiographie appliquée à l'étude macroscopique de l'évolution des testicules d'Oiseaux - C.R.S.B., 144, 1107, 1950.
35. BENOIT, J., WALTER, F.X. and ASSENMACHER, I. : Nouvelles recherches relatives à l'action de lumières de différentes longueurs d'onde sur la gonadostimulation du Canard mâle impubère - C.R.S.B., 144, 1206, 1950.
36. BENOIT, J., WALTER, F.X. and ASSENMACHER, I. : Contribution à l'étude du réflexe opto-hypophysaire gonadostimulant chez le Canard soumis à des radiations lumineuses de diverses longueurs d'onde - Congrès de l'Assoc. Physiologistes langue française mai 1950. Journal Physiologie, 42, 537-541, 1950.
37. BENOIT, J., MANDEL, P., WALTER, F.X. and ASSENMACHER, I. : Sensibilité testiculaire aux hormones gonadotropes hypophysaires chez le Canard domestique au cours de la période de régression testiculaire saisonnière - C.R.S.B., 144, 1400, 1950.
38. BENOIT, J., ASSENMACHER, I. and WALTER, F.X. : Activité gonadotrope de l'hypophyse du Canard domestique, au cours de la régression testiculaire saisonnière et de la prépuberté - C.R.S.B., 144, 1403, 1950.
39. BENOIT, J., and ASSENMACHER, I. : Quelques données relatives à la vascularisation de l'hypophyse du Canard domestique - Congrès Anatomistes suisses, sept. 1950, cf. Bull. Histol. appliquée, 27, 182, 1950.
40. BENOIT, J. and ASSENMACHER, I. : Etude préliminaire de la vascularisation de l'appareil hypophysaire du Canard domestique - Arch Anatomie Microscop. et de Morphol. expérimentale, 40, 27-45, 1951.
41. BENOIT, J. : Circulation porte tubéro-hypophysaire chez le Canard domestique - C.R.S.B., 145, 1112, 1951.
42. BENOIT, J. : La pars tuberalis de l'hypophyse du Canard. Ses rapports avec l'éminence médiane et la préhypophyse - C.R.Ass.Anat. Nancy, 65, 154-160, 1951.
43. BENOIT, J. and ASSENMACHER, I. : Dispositifs nerveux de l'éminence médiane, leurs rapports avec la vascularisation hypophysaire chez le Canard domestique - C.R.S.B., 145, 1395, 1951.
44. BENOIT, J. and ASSENMACHER, I. : Contribution à l'étude des relations hypothalamohypophysaires et de leur rôle dans la gonadostimulation chez le Canard domestique - J. Physiologie, 43, 643-645, 1951.
45. BENOIT, J., ASSENMACHER, I. and WALTER, F.X. : Différences de sensibilité de la rétine du Canard aux radiations colorées dans le réflexe pupillaire et dans le réflexe opto-sexuel - C.R.S.B., 146, 1027, 1952.
46. BENOIT, J. : Action de la lumière blanche et des radiations colorées, par l'intermédiaire de l'oeil, sur le système nerveux et en particulier sur le système neuro-végétatif - Rapport au Congrès d'Avignon: 3è Congrès technique National de Sécurité et d'Hygiène du Travail, 9 à 12 octobre 1952.
46. BENOIT, J. and ASSENMACHER, I. : Influence de lésions hautes et basses de l'infundibulum sur la gonadostimulation chez le Canard domestique - C.R. Ac.Sc., 235, 1547, 1952.
47. BENOIT, J. and ASSENMACHER, I. : Contribution à l'étude des relations de la substance Gomori positive avec le complexe hypophysaire et la gonadostimulation chez le Canard domestique - C.R.As.Sc., 236, 133, 1953.

48. BENOIT, J., ASSENMACHER, I. and MANUEL, S.: Pénétration, variable selon la longueur d'onde, des radiations visibles jusqu'à l'hypothalamus et au rhinencéphale, à travers la région orbitaire chez le Canard - C.R. Ac.Sc., 235, 1695, 1953.
49. BENOIT, J., ASSENMACHER, I. and MANUEL, S.: Pénétration des radiations visibles jusqu'à l'encéphale à travers la région orbitaire, chez le Canard. Sa mesure par un procédé photographique - C.R.S.B., 147, 40, 1953.
50. BENOIT, J., ASSENMACHER, I. and WALTER, F.X.: Dissociation expérimentale des rôles des récepteurs superficiel et profond dans la gonadostimulation hypophysaire par la lumière chez le Canard - C.R.S.B., 147, 186, 1953.
51. BENOIT, J., and ASSENMACHER, I.: Rôle des photo-récepteurs superficiels et profond dans la gonadostimulation par la lumière chez les Oiseaux - Journal de Physiol., 45, 34-37, 1953.
52. BENOIT, J. and ASSENMACHER, I.: Répercussions de la section du tractus porto-tubéral hypophysaire sur la gonadostimulation par la lumière chez la Canard domestique - C.R. Ac.Sc., 236, 2002-4, 1953.
53. BENOIT, J. and ASSENMACHER, I.: Rapport entre la stimulation sexuelle préhypophysaire et la neurosécrétion chez l'Oiseau - Convegno sulla Neurosecrezione, Pubblicazioni della Stazione Zoologica di Napoli, 24, 27-31, 1954.
54. BENOIT, J. and ASSENMACHER, I.: Action des facteurs externes et plus particulièrement du facteur lumineux sur l'activité sexuelle des Oiseaux - Rapport à la IIe Réunion des Endocrinologistes de langue française, 1953.
55. BENOIT, J. and ASSENMACHER, I.: Rapport entre la stimulation sexuelle préhypophysaire et la neurosécrétion chez l'Oiseau - Arch. Anat. micr. et de Morphologie exp., 42, 334-386, 1953.
56. BENOIT, J.: Réflexe opto-pituitaire-sexuel et relations hypothalamo-hypophysaires - Bull. de l'Académie Nationale de médecine, 138, 32-36, 1954.
57. BENOIT, J. and ASSENMACHER, I.: Sensibilité comparée des récepteurs superficiels et profonds dans le réflexe opto-sexuel chez le Canard - C.R. Ac.Sc., 239, 105-107, 1954.
58. BENOIT, J., TAUC, L. and ASSENMACHER, I.: Mesure photo-électrique de la pénétration trans-orbitaire des radiations visibles jusqu'au cerveau, chez le Canard domestique - C.R. Ac.Sc., 239, 451-453, 1954.
59. BENOIT, J., TAUC, L. and ASSENMACHER, I.: Nouveaux résultats de mesure photo-électrique de la pénétration de radiations lumineuses visibles jusqu'au cerveau, par le côté et le sommet de la tête, chez les Canards blancs et pigmentés - C.R. Ac.Sc., 239, 508-510, 1954.
60. BENOIT, J., ASSENMACHER, I. and BRARD, E.: Evolution testiculaire du Canard domestique maintenu à l'obscurité totale pendant une longue durée - C.R. Ac.Sc., 241, 251-253, 1955.
61. BENOIT, J., MESSERSCHMITT, J. and STORA, C.: Inégale croissance du tractus génital de la Souris blanche femelle impubère, selon l'époque de sa naissance dans l'année - Livre jubilaire de L. NOUNOURE, Arch. Anat., Histol. et Embryol., 37, 29-36, 1955.
62. BENOIT, J. and ASSENMACHER, I.: Le contrôle hypothalamique de l'activité préhypophysaire gonadotrope - J. Physiol., 47, 3, 427-567, 1955 and J. Physiol., 47, 61-66, 1955.
63. BENOIT, J., ASSENMACHER, I. and BRARD, E.: Apparition et maintien de cycles sexuels non saisonniers chez le Canard domestique placé pendant plus de trois ans à l'obscurité totale - Congrès de Physiol.; J. de Physiologie, 48, 388-391, 1956.
64. BENOIT, J. and ASSENMACHER, I.: Quelques aspects du contrôle hypothalamique de la fonction gonadotrope de la préhypophyse - Symposium internationale sul Diencefalo, Milano, 54-56, 1956.
65. BENOIT, J. and ASSENMACHER, I.: Nouvelles recherches sur les relations entre la neurosécrétion hypothalamique, le système porte hypophysaire et l'activité gonadotrope de la préhypophyse du Canard domestique - C.R. Ac.Sc., 242, 2986, 1956.
66. BENOIT, J., ASSENMACHER, I. and BRARD, E.: Etude de l'évolution testiculaire du Canard domestique soumis très jeune à un éclairage artificiel permanent pendant deux ans - C.R. Ac.Sc., 242, 3113-3115, 1956.
67. BENOIT, J. and ASSENMACHER, I.: Photorécepteurs rétiniens et encéphaliques impliqués dans la gonadostimulation par les radiations visibles chez le Canard domestique - Rapport 2nd Congress for Endocrinologists, Torino, 1957.
68. BENOIT, J.: Radiations lumineuses et activité sexuelle du Canard - Revue Suisse de Zool., 64, 30, 577-587, 1957.
69. BENOIT, J.: Rôle de l'oeil et de l'hypothalamus dans la stimulation par les radiations visibles des glandes génitales chez les oiseaux - Les Ann. d'oculistique, 191, 1, 1-14, 1958.

70. BIANCA, W. : Untersuchungen über die Methodik der Blutmengenbestimmung am lebenden Tier unter besonderer Berücksichtigung der Blutspeicher. Das Blutvolumen bei Pferd, Rind, Schaf und Ziege - Ztschr. für Tierzüchtung und Züchtungsbiologie, 1, 59, 1950.
71. BIANCA, W. : Die Bestimmung der Gesamtblutmenge in vivo unter Verwendung eines Pharmakons zur Entleerung der Blutspeicher - Cardiologia, 19, 5, 283, 1951.
72. BIANCA, W. : Zur Bestimmung des wahren Lebendgewichtes beim trächtigen Rind - Schweizer Landwirtschaftliche Monatshefte, 7, 243, 1951.
73. FOLK, G.E. and RIEDESEL, M. : Serum magnesium changes in hibernation - Nature, 177, 668, 1956.
74. FOLK, G.E. and FARRAND, R.L. : Characteristics of hibernation of the golden hamster - British Journal Am. Techn. Assoc., 7, 14, 1956.
75. FOLK, G.E. and FARRAND, R.L. : Factors influencing the nutritional status of the Hamster - Proc. of the Animal Care Panel, 7, 1956.
76. FOLK, G.E., FARRAND, R.L. and RIEDESEL, M.L. : Types of Mammalian Hibernation - Proc. Iowa Acad. Sci., 63, 724, 1956.
77. FOLK, G.E. and RIEDESEL, M.L. : Serum magnesium changes in mammals exposed to cold - Journal of Mammology, 38, 1957.
78. FOLK, G.E. : Biological Rhythms of Mammals in the Cold - The Am. Naturalist (Symposium volume). In press, 1957.
79. FOLK, G.E., DODGE, C.H. and FOLK, M.A. : Resting body temperatures of raccoons and domestic rabbits - Proc. of the Animal Care Panel, 8, 1957.
80. FOLK, G.E. : Twenty-four hour rhythms of Mammals in a cold environment - The Am. Naturalist, 91, 153-166, 1957.
81. FOLK, G.E. and CARTER, J.R. : Effects of three winters of hibernation upon thirteen-lined ground squirrels - Anat. Record, 127, 1957.
82. FOLK, G.E. and MELTZER, M.R. : A method for correlating bladder emptying and total activity in rodents - Journ. of Mammal., 39, 2, 1957.
83. FOLK, G.E. and FARRAND, R.L. : Responses of Hamsters during the first two weeks of Cold exposure - Fed. Proc., 16, 1957.
84. FRY, F.E.J. a.o. : Lethal limits of temperature for young goldfish - Rev. Can. Biol., 1, 50-56, 1941.
85. FRY, F.E.J. a.o. : Lethal temperature relations for a sample of young speckled trout, (*Salvelinus fontinalis*) - U. of T. Studies Biol. Ser. No. 54, Publ. Ont. Fish. Res. Lab., 66, 1-47, 1945.
86. FRY, F.E.J. a.o. : Influence of temperature on the asphyxiation of young goldfish (*Carassius auratus* L.) under various tensions of oxygen and carbon dioxide - Biol. Bull., 92, 217-224, 1947.
87. FRY, F.E.J. a.o. : The relation of temperature to oxygen consumption in the goldfish - Biol. Bull., 94, 66-77, 1948.
88. FRY, F.E.J. and BLACK, E.C. : The influence of carbon dioxide on the utilization of oxygen by some freshwater fish - Ont. Fish. Res. Lab. Univ. Toronto; & Edw. Martin Biol. Lab. Swarthmore Coll., Can. Zool., 32, 408-420, 1954.
89. GELINEO, S. : La thermogénèse des oiseaux après un séjour prolongé à différentes températures - C. r. Soc. Biol., 116, 672, 1934.
90. GELINEO, S. : Sur la thermogénèse de l'hibernant pendant l'été - C. r. Soc. Biol., 127, 1357-1359, 1938.
91. GELINEO, S. : Sur la thermogénèse de l'hibernant lors du passage de l'état de veille à l'état de torpeur - C. r. Soc. Biol., 127, 1360-1362, 1938.
92. GELINEO, S. : Contribution à l'étude de la thermorégulation et de la thermogénèse des hibernants. I. La thermorégulation et la thermogénèse du *Spermophile* (*Citellus citellus*) pendant l'été - Bull. Acad. r. serbe, B. 5, 197-217, 1939.
93. GELINEO, S. : Sur la thermorégulation et la thermogénèse du Rat durant un séjour prolongé aux températures provoquant l'hyperthermie - Bull. Acad. r. serbe, 6, 149-184, 1940.
94. GELINEO, S. and GELINEO, A. : Environmental temperature of the nest and the appearance of chemical thermoregulation in rats at the temperature of 21° C - Bull. de l'Acad. serbe des Sciences, 3, 149-153, 1951.
95. GELINEO, S. and GELINEO, A. : Sur la thermogénèse du Rat nouveau-né et la température du nid - C. r. Acad. des Sciences Paris, 232, 1031-1032, 1951.
96. GELINEO, S. and GELINEO, A. : Oxidation in rats the first days after birth - Bull. de l'Acad. serbe des Sci., 3, 119-123, 1951.
97. GELINEO, S. : La température du nid du rat et sa signification biologique - Bull. Acad. serbe des Sciences, 4, Sci. naturelles, 2, 197-210, 1952.
98. GELINEO, S. : Sur la thermorégulation chimique optimum du pigeon domestique - C. r. Soc. Biol., 147, 134, 1953.

99. GELINEO, S. : La naissance et le développement de la thermorégulation chimique chez le spermophile (*Citellus citellus*) - Bull. de l'Acad. serbe des Sci., 12, 1953.
100. GELINEO, S. and SOKIĆ, P.: Sur l'apparition de l'homeothermie chez le spermophile (*Citellus citellus* L.).- C.r. Soc. Biol., 147, 138, 1953.
101. GELINEO, S. : Influence du milieu thermique d'adaptation sur la dépense d'oxygène chez l'escargot *Helix pomatia* - Bull. de l'Acad. serbe des Sciences, 12, 1953.
102. GELINEO, S. : La pellagre expérimentale chez la spermophile (*Citellus citellus* L.) - Bull. de l'Acad. serbe des Sciences, 11, 97-103, 1954.
103. GELINEO, S. : Température d'adaptation et production de chaleur chez les Oiseaux de petite taille - Arch. de Sci. physiol., 9, 225-143, 1955.
104. GELINEO, S., GELINEO, A. and RAJEVSKAJA: La concentration de l'hémoglobine dans le sang chez quelques Oiseaux - C.r. So. Biol., 149, 1411-1413, 1955.
105. GELINEO, S. and GELINEO, A.: La concentration de l'hémoglobine dans le sang chez des poissons d'eau douce - C.r. Soc. Biol. 149, 1410-1411, 1955.
106. GELINEO, S. : Température d'adaptation et consommation d'oxygène chez le Rat blanc - C.r. Acad. des Sciences, Paris, 241, 367-369, 1955.
107. GELINEO, S. and GELINEO, A.: La consommation d'oxygène chez différentes espèces de Lézards à habitats différents - C.r. Soc. Biol., 149, 565-568, 1955.
108. GELINEO, S. and GELINEO, A.: Les échanges respiratoires de Lézards noir de Dalmatie adaptés à différentes températures - C.r. Soc. Biol., 149, 387-189, 1955.
109. HART, J.S., FRY, F.E.J. and WALKER, K.F.: Lethal temperature relations for a sample of young speckled trout (*Salvelinus fontinalis*) - Univ. of Toronto Stud. Biol. Ser. 54: Pub.Ont. Fish.Res.Lab., 66, 1946.
110. HART, J.S. : Lethal temperature relations for certain fish or the Toronto region - Trans.Royal Soc. of Can., 5, 51-71, 1947.
111. HART, J.S. and FRY, F.E.J.: The cruising speed of goldfish in relation to water temperature - Jr. Fish. Res. Bd. Can., 7, 169-175, 1948.
112. HART, J.S. and FRY, F.E.J.: The relation of temperature to oxygen consumption in the goldfish - Abstract Anat. Rec., 96, 4, 1946, Supp. December; Biol. Bull., 94, 66-77, 1948.
113. HART, J.S. : Interrelations of daily metabolic cycle, activity and environmental temperature of mice - Can.Res.D., 28, 293-307, 1950.
114. HART, J.S. : The effect of humidity and temperature on carbon dioxide production of deer mice and on heat transmission through their fur - Can.J. Res. D., 28, 280-284, 1950.
115. HART, J.S. : Average body temperature in mice - Science, 113, 325-326, 1951.
116. HART, J.S. : Calorimetric determination of average body temperature of small mammals and its variation with environmental conditions - Can.J. Zool., 29, 224-233, 1951.
117. HART, J.S. : Use of daily metabolic periodicities as a measure of the energy expended by voluntary activity of mice - Can.J. Zool., 30, 83-89, 1952.
118. HART, J.S. : Effects of temperature and work on metabolism, body temperature and insulation: results with mice - Can.J.Zool., 30, 90-98, 1952.
119. HART, J.S. : Energy metabolism of the white-footed mouse. *Peromyscus leucopus noveboracensis*, after acclimatization at various environmental temperatures - Can.J.Zool., 31, 99-105, 1953.
120. HART, J.S. : The relation between thermal history and cold resistance in certain species of rodents - Can.J. Zool., 31, 80-98, 1953.
121. HART, J.S. : The influence of thermal acclimatization on limitation of running activity by cold deer mice - Can.J. Zool., 31, 117-120, 1953.
122. HART, J.S. : Rate of gain and loss of cold resistance in mice - Can.J. Zool., 31, 112-116, 1953.
123. HART, J.S. : A comparison of some seasonal and temperature-induced changes in *peromyscus*: cold resistance, metabolism and pelage insulation - Can. J. Zool., 31, 528-534, 1953.
124. HART, J.S. and HEROUX, O. : Restraint hypothermia and its inhibition by cold acclimatization - Am. J. Phys., 177, 219-221, 1954.
125. HART, J.S. and HEROUX, O.: Effect of low temperature and work on blood lactic acid in deer mice - Am.J.Physiology, 176, 452-454, 1954.
126. HART, J.S. and HEROUX, O.: Comparison of four indices of adrenal activity in rats acclimated to 30°, 15°, 1° C. - Am J. Phys., 178, 445-448, 1954.
127. HART, J.S. and HEROUX, O.: Adrenal cortical hormone requirement of warm and cold acclimated rats after adrenalectomy - Am.J. Phys., 178, 449-452, 1954.
128. HART, J.S. and HEROUX, O.: Cold acclimatization and adrenal cortical activity as measured by eosinophil levels - Am.J. Phys., 178, 453-456, 1954.

129. HART, J.S. and HEROUX, O.: Exercise and temperature regulation in lemmings and rabbits - Can.J. Biochem. and Physiol., 33, 428-435, 1955.
130. HART, J.S. : Geographic variation of some physiological and morphological characters in certain freshwater fish - Univ. of Toronto Biol. Series no. 60, Pub. Ontario Fisheries Res. Lab., 72.
131. HART, J.S. : Seasonal changes in insulation of the fur - Can.J. Zool., (in press)
132. HOCK, M.J. and SCHOLANDER, P.F.: Body insulation in some arctic and tropical mammals and birds - Biol. Bull., 99, 225-236, 1950.
133. HOCK, R.J. and SCHOLANDER, P.F.: Heat regulation in some arctic and tropical mammals and birds - Biol. Bull., 99, 237-258, 1950.
134. HOCK, M.J. a.o. : Adaptation to cold in arctic and tropical mammals and birds in relation to body temperature, insulation and basal metabolic rate - Biol. Bull., 99, 259-271, 1950.
135. HOCK, R.J. : Nature of the hibernation of bats - Fed. Proc., 10, 65, 1951.
136. HOCK, R.J. : The metabolic rates and body temperatures of bats - Biol. Bull., 101, 289-299, 1951.
137. HOCK, R.J. : Rectal temperatures of the black bear during its "hibernation" - Science in Alaska (Proc. 2nd Alaska Sci. Conf.), 310-312, 1951.
138. HOCK, R.J. : Seasonal changes in the blood of ground squirrels - Science in Alaska, 137-138, 1952.
139. HOCK, R.J. : Photoperiod as stimulus for onset on hibernation - Fed. Proc., 14, 73-74, 1955.
140. HOCK, R.J. : Body temperature variations of non-hibernating Alaskan ground squirrels, Fed. Proc. 15. In press.
141. HOCK, R.J. : Physiological values in hibernation of vertebrates - Handb. Biol. Data, in press.
142. HOCK, R.J. : Body temperatures of Arctic animals - Handb. Biol. Data. In press.
143. IRVING, L., SCHOLANDER, P.F., HOCK, R.J., WALTERS, V. and JOHNSON, F.: Heat regulation in some arctic and tropical mammals and birds - Biol. Bull., 99, 2, 225-271, 1950.
144. IRVING, L., SCHOLANDER, P.F., HOCK, R.J. and WALTERS, V.: Adaptation to cold in arctic and tropical mammals and birds in regulation to body temperature, insulation, and basal metabolic rate - Biol. Bull., 99, 2, 225-271, 1950.
145. IRVING, L., SCHOLANDER, P.F., WALTERS, V. and HOCK, R.J.: Body insulation of some arctic and tropical mammals and birds - Biol. Bull., 99, 2, 225-271, 1950.
146. IRVING, L. : Physiological adaptation to cold in arctic and tropic animals - Federation Proc., 10, 2, 1951.
147. IRVING, L. : Respiration in some arctic and tropical lichens in relation to temperature - Am. J. of Botany, 39, 10, 707-713, 1952.
148. IRVING, L., CHATEFIELD, P.O. and LYMAN, C.P.: Physiological Adaptation to cold of Peripheral Nerve in the leg of the herring gull (*Larus Argentatus*) - The Am. J. of Phys., 172, 3, 1953.
149. IRVING, L., SCHOLANDER, P.F., FLAGG, W. and WALTERS, V.: Climatic adaptation in arctic and tropical poikilotherms - Phys. Zool., 26, 1, 1953.
150. IRVING, L., KROG, J.: Arctic and Subarctic Birds and Mammals - J. of Applied Phys., 6, 11, 1954.
151. IRVING, L., KROG, H. and MONSON, M.: The metabolism of some Alaskan animals in winter and summer - Phys. Zool., 28, 3, 1955.
152. IRVING, L. : Nocturnal decline in the temperature of birds in cold weather - The Condor, 57, 6, 362-365, 1955.
153. IWONSKI, B. : Untersuchungen über die physiologischen Schwankungen der Körpertemperatur, Puls- und Atemfrequenz der Ziege - Med. Vet. Diss., Wien, 1951.
154. KOPF, N. : Einfluss des Luftdruckes auf die Blutsenkungsgeschwindigkeit bei Rindern - Med. Vet. Diss., Wien, 1952.
155. MORRISON, P.R. : Acquired homiothermism in the pregnant sloth - J. Mamalogy, 26, 272-275, 1945.
156. MORRISON, P.R. : Observations on water loss and oxygen consumption in Peripatus - Biol. Bull., 91, 181-188, 1946.
157. MORRISON, P.R. : Temperature regulation in three Central American mammals - J. Cell. Comp. Physiol., 27, 125-138, 1946.
158. MORRISON, P.R. and PEARSON, O.P.: The metabolism of a very small mammal - Science, 104, 287-289, 1946.
159. MORRISON, P.R. : Oxygen consumption in several small wild mammals - J. Cell. Comp. Physiol., 31, 69-96, 1948.
160. MORRISON, P.R. : Oxygen consumption in several mammals under basal conditions - J. Cell. Comp. Physiol., 31, 281-292, 1948.
161. MORRISON, P.R. and RYSER, F.A.: The development of temperature regulation in the Alaskan red-backed vole, *Clethrionomys rutilus dawsoni* - Anat. Rec. 113, 1952.

162. MORRISON, P.R. and RYSER, F.A.: Weight and body temperature in mammals - Science, 116, 231-232, 1952.
163. MORRISON, P.R. and RYSER, F.A.: Cold resistance in the young ringnecked pheasant - The Auk, 71, 253-266, 1954.
164. MORRISON, P.R., RYSER, F.A. and STRECKER, R.L.: Growth and the development of temperature regulation in the tundra redback vole - J. Mammalogy, 35, 376-386, 1954.
165. MORRISON, P.R. and DAWE, A.R.: Characteristics of the hibernating heart - Amer.Heart J., 49, 367-384, 1955.
166. PASTERK, J. : Das Blutbild des Meerschweinchens nach grösseren Temperaturschwankungen - Med.Vet.Diss., Wien, 1951.
167. PRYCHODKO, W. : Effect of aggregation of laboratory mice (*mus musculus*) of food intake at different temperatures - Ecology, 39, 3, 500-503, 1958.
168. ROBINSON, K.W. and LEE, D.H.K.: Animal behaviour and heat regulation in hot atmospheres - Univ. Qld.Papers, Dept. Physiol., 1, 9, 1-8, 1946.
169. ROBINSON, K.W. and LEE, D.H.K.: The effect of the nutritional plane upon the reactions of animals to heat - J. anim. Sci., 6, 182-194, 1947.
170. ROBINSON, K.W. : Heat tolerances of Australian monotremes and marsupials - Austr. J. Biol. Sci., 7, 348-360, 1954.
171. ROBINSON, K.W. and MACFARLANE, W.V.: The influence of environmental temperature on the level of plasma antidiuretic substances in the rat - Austr. J. biol. Sci., 9, 130-138, 1956.
172. ROBINSON, K.W. and MORRISON, P.R.: The reaction to hgt atmospheres of various species of Australian marsupials and placental mammals - J. cell.comp.Physiol., 1956. In press.
173. SAPIN-JALOUSTRE, J. : Mammifères et Oiseaux de Terre Adélie - C.R. mensuels des seances de l'Académie des Sciences Coloniales, 11, 480-497, 1951.
174. SCHMIDT-NIELSEN, B. and SCHMIDT-NIELSEN, K.: Water metabolism of desert mammals - Phys. Reviews, 32, 2, 1952.
175. SCHMIDT-NIELSEN, K. : Animals and arid conditions: Physiological aspects of productivity and management - The future of Arid lands. Published by the American Ass. for the Advancement of Science, Washington D.C., 1956.
176. SCHMIDT-NIELSEN, K., SCHMIDT-NIELSEN, B., HOUP, T.R. and JARNUM, S.A.: The question of water storage in the stomach of the camel - Extrait de Mammalia, 20, 1956.
177. SCHMIDT-NIELSEN, K., SCHMIDT-NIELSEN, B., HOUP, T.R. and JARNUM, S.A.: Water balance of the camel - The Am.Journal of Phys., 185, 1, 1956.
178. SCHOLANDER, P.F. a.o.: Studies in the physiology of frozen plants and animals in the Arctic - J. Cell.Comp. Physiol.suppl. 1, 42, 1-56, 1953.
179. SCHUA, L. : Beobachtungen über das Auftreten von Winterschlaf bei Goldhamstern und über gewisse Wetterabhängigkeit derselben - Säugetierk. Mitteilungen, 2, 163, 1954.
180. SEALANDER, J.A. : Effect of environmental temperature and starvation on adrenal glands of the white-footed mouse - Amer.J. Physiol., 163, 92-95, 1950.
181. SEALANDER, J.A. : Survival of *Peromyscus* in relation to environmental temperature and acclimatization at high and low temperatures - Amer.Midl.Nat., 46, 257-311, 1951.
182. SEALANDER, J.A. : The relationship of most protection and huddling to survival of *Peromyscus* at low temperature - Ecology, 33, 63-71, 1952.
183. SEALANDER, J.A. : Food consumption in *Peromyscus* in relation to air temperature and previous thermal experience - J. Mamm., 33, 206-218, 1952.
184. SEALANDER, J.A. : Body temperatures of white-footed mice in relation to environmental temperature and heat and cold stress - Biol. Bull., 104, 87-99, 1953.
185. SOLLBERGER, A., ELFVIN, L.G. and PETREN, T.: Influence of some endogenous and exogenous factors on diurnal glycogen rhythm in chicken - Acta Anatomica, 25, 286-309, 1955.
186. STINSON, R.H. and FISHER, K.C.: Temperature selection in deer mice - Can. Journ. Zool., 31, 404-416, 1953.
187. WILBER, C.G. : The effects of increased environmental temperatures on the lipids in the blood of the turtle, *chrysemys picta* - St.Louis Univ.Studies Ser. C., 1, 1-20, 1950.
188. WILBER, C.G. : Fat usage in arctic compared with tropical zone mammals - med. Bull. St. Louis Univ., 4, 112-119, 1951.
189. WILBER, C.G. : The effect of ultra violet light on the protoplasm in *pelomyxa carolinensis* - Trans.Amer.Mic.Soc., 70, 265-271, 1951.
190. WILBER, C.G. : The lipids in the tissues of arctic animals - Rpt. 39, Chemical Corps Medical labor. Army Chemical Center, Maryland, 1954.

191. WILBER, C.G. : Heat stress and blood constituents in the painted turtle, *Chrysemys picta* - Rpt. 280, Chemical Corps medical Laboratories, Army Chemical Center Maryland, 1954.
192. WILBER, C.G. : Electrocardiogram of the opossum during exposure to cold - *Am.J.Phys.* 181, 352-355, 1955.
193. WILBER, C.G. and ROBINSON, P.F.: Effect of restraint on body temperature in Guinea Pigs - *J. of Appl.Phys.*, 12, 2, 214-216, 1958.
194. WILBER, C.G. and ROBINSON, P.F.: Aspects of blood chemistry in the white-tailed deer - *J. of Mamm.*, 39, 2, 309-311, 1958.
195. WILBER, C.G. : Influence of temperature on performance in Guinea Pigs - *The Am. J. of Phys.*, 190, 3, 457-458, 1958.
196. YAGI, N. : On the range of lethal body temperature of the rat - *Sci.Rep. Tokyo Bunrika Daigaku*, 22, 1934.
197. ZÖCHLING, F. : Untersuchungen über die Zusammenhänge zwischen Luftdruckschwankungen und Leukozytenzahlen bei Rindern - *Med.Vet.Diss.*, Wien, 1952.

ENTOMOLOGICAL BIOCLIMATOLOGY

1. ALBRECHT, F.O. : Limitation des effectifs chez Acridien: Influence de la sécheresse du sol sur les oeufs de *Nomadacris septemfasciata* (Serv.) - *Locusta*, 4, 1956.
2. ANDREWARTHA, H.G. and VEVERS STEELE, H.: Thrips investigation. 4. Some observations on the fluctuations in the numbers of Thrips *imagnis* Bagnall in the vicinity of Melbourne during the period 1932-1934 - *C.S.I.R. Journal*, 7, 4, 234-238, 1934.
3. ANDREWARTHA, H.G. : Thrips investigation. 5. On the effect of soil moisture on the viability of the Pupal Stages of Thrips *imagnis* Bagnall - *C.S.I.R. J.*, 7, 4, 239-244, 1934.
4. ANDREWARTHA, H.G. : The environment of the Australian plague locust (*Chortoicetes terminifera* Walk.) in South Australia - *Trans.Roy Soc. of S.Aust.*, 64, 76-94, 1940.
5. ANDREWARTHA, H.G. and BIRCH, L.C.: The influence of weather on grasshopper plagues in South Australia - *Journ. Agric. S.Aust.*, 45, 95-100, 1941.
6. ANDREWARTHA, H.G. : The distribution of plagues of *Austroicetes cruciata* Sauss. (Acrididae) in Australia in relation to climate vegetation and soil - *Trans. Roy. Soc. of S. Aust.*, 68, 315-326, 1944.
7. ANDREWARTHA, H.G. : Some differences in the physiology and ecology of locusts and grasshoppers - *Bull. Entom. Research*, 35, 379-389, 1945.
8. ANDREWARTHA, H.G. and DAVIDSON, J.: Annual trends in a natural population of Thrips *imagnis* Bagnall (Thysanoptera) - *J. Anim. Ecology*, 17, 193-199, 1948.
9. ANDREWARTHA, H.G. and DAVIDSON, J.: The influence of rainfall, evaporation and atmospheric temperature on fluctuations in the size of a natural population of Thrips *imagnis* Bagnall (Thysanoptera) - *J. Anim.Ecol.*, 17, 200-222, 1948.
10. ANDREWARTHA, H.G. : The distribution and abundance of animals - Chicago University, 1954. (Book).
11. BEIRNE, B.P. : The causes of the occasional abundance or scarcity of wasps - *Ent. Mon. Mag.*, 80, 121-124, 1944.
12. BEIRNE, B.P. : The origin and history of the British Macro-Lepidoptera - *Trans. R. Ent. Soc. Lond.*, 98, 273-372, 1947.
13. BEIRNE, B.P. : The origin and History of the British Fauna - Methuen & Co., London, 1952.
14. BEIRNE, B.P. : Natural fluctuations in abundance of British Lepidoptera - *Ent.Gaz.*, 6, 21-52, 1955.
15. BEIRNE, B.P. : Zoogeographical significance of distributional changes in British Macro-Lepidoptera - *Trans. R. Ent.Soc. Lond.*, 107, 117-123, 1955.
16. BEIRNE, B.P. : Effects of climate and weather on distribution and abundance of insects - Areal report: Symposium Int. Soc. Bioclim. and Biomet., Paris, 29-31 August 1956.
17. CHOUDHURI, J.S.B. : Experimental studies on the selection of oviposition sites by Locust *migratoria migratorioides* (R. et F.) - See ALBRECHT F.O., *Locusta*, 4, 1956.
18. CLOUDSLEY-THOMPSON, J.L.: Observations on the ecology of Arachnids in North-West Iceland - *Ann.Mag.Nat.Hist.*, 12, 1, 437-447, 1948.
19. CLOUDSLEY-THOMPSON, J.L.: The enemies of Miriapods - *Naturalist*, 137-141, 1949.

20. CLOUDSLEY-THOMPSON, J.L.: On spiders from North-East Greenland. 1948 - Ann.Mag.Nat.Hist., 12, 2, 398-400, 1949.
21. CLOUDSLEY-THOMPSON, J.L.: The significance of migration in Myriapods - Ann.Mag.Nat.Hist., 12, 2, 947-962, 1949.
22. CLOUDSLEY-THOMPSON, J.L.: On the responses to environmental stimuli and the sensory physiology of millipedes (Diplopoda) - Proc.Zool.Soc.London, 121, 253-277, 1951.
23. CLOUDSLEY-THOMPSON, J.L.: Studies in diurnal rhythms. I. Rhythmic behaviour in millipedes - J. Exp. Biol., 28, 165-172, 1951.
24. CLOUDSLEY-THOMPSON, J.L.: Studies in diurnal rhythms. II. Changes in the physiological responses of the woodlouse *Oniscus asellus* to environmental stimuli - J. Exp. Biol., 29, 295-303, 1952.
25. CLOUDSLEY-THOMPSON, J.L.: Behaviour of centipedes and millipedes. I. Responses to environmental stimuli - Ann.Mag.Nat.Hist., 12, 5, 417-434, 1952.
26. CLOUDSLEY-THOMPSON, J.L.: Diurnal Rhythms - Trans. IX.Int.Congr. Ent., 1, 305-310, 1952.
27. CLOUDSLEY-THOMPSON, J.L.: Diurnal Rhythms of locomotory activity in isolated migratory locusts - Ent.Mon.Mag., 89, 233-235, 1953.
28. CLOUDSLEY-THOMPSON, J.L.: The significance of fluctuating temperatures on the physiology of insects - Entomologist, 86, 183-189, 1953.
29. CLOUDSLEY-THOMPSON, J.L.: Studies in diurnal Rhythms. III. Photoperiodism in the cockroach *Periplaneta americana* (L) - Ann.Mag.Nat.Hist., 12, 6, 705-712, 1953.
30. CLOUDSLEY-THOMPSON, J.L.: Studies in diurnal Rhythms. IV. Photoperiodism and geotaxis in *Tenebrio molitor*, L. (Coleoptera: Tenebrionidae) - Proc. R. Ent.Soc. Lond.A., 28, 117-132, 1953.
31. CLOUDSLEY-THOMPSON, J.L.: Millipedes entering houses - The Entomologist's Monthly Magazine, 90, 1954.
32. CLOUDSLEY-THOMPSON, J.L.: Problems of dispersal in some terrestrial Arthropods - Advanc.Sci., 11, 73-75, 1954.
33. CLOUDSLEY-THOMPSON, J.L.: The Black-fly, simulium *Venustum*, biting man - The Entomologist, 87, 1100, 17-20, 1955.
34. CLOUDSLEY-THOMPSON, J.L.: Some aspects of the biology of centipedes and scorpions - Naturalist, 147-153, 1955.
35. CLOUDSLEY-THOMPSON, J.L.: Entomological Bioclimatology - See Areal report Dr. J.L.Cloudsley-Thompson. Symposium Int. Soc. Bioclim. and Biomet. Paris, 29 - 31 August 1956.
36. CLOUDSLEY-THOMPSON, J.L.: Studies in diurnal Rhythms. Diurnal Rhythms of activity in Terrestrial Arthropods - Nature, 178, 215, 1956.
37. CLOUDSLEY-THOMPSON, J.L.: The life histories of the British cribellate spiders of the genus *Ciniflo* Bl. (Dictynidae) - Ann.Mag.Nat.Hist., 12, 8, 787-794, 1956.
38. CLOUDSLEY-THOMPSON, J.L.: Notes on the Arachnide 1-28 - Ent.Mon.Mag. 1947-1956.
39. CLOUDSLEY-THOMPSON, J.L.: Studies in diurnal rhythms. VI. Bioclimatic observations in Tunisia and their significance in relation to the physiology of the fauna, especially woodlice, centipedes, scorpions and beetles - Annals and Magazine of Nat.Hist., 12, 9, 305, 1956.
40. CLOUDSLEY-THOMPSON, J.L.: Studies in diurnal Rhythms. VII. Humidity responses and nocturnal activity in woodlice (Isopoda) - The Journal of experimental Biology, 33, 576-582, 1956.
41. CLOUDSLEY-THOMPSON, J.L.: The effect of rock cover on the diurnal range of Microclimatic conditions - The Entomologist, 89, 1120, 1956.
42. CLOUDSLEY-THOMPSON, J.L.: Studies in diurnal rhythms. The ecological significance of diurnal rhythms in terrestrial arthropods - Sci.Prog., 42, 46-52, 1954.Proc. XIV Int.Congr.Zool., 415-417, 1956.
43. CLOUDSLEY-THOMPSON, J.L., and SANKEY, J.H.P.: Some aspects of the fauna of the district around Juniper Hall Nickleham, Surrey-III. - The Entomologist's Monthly Magazine, 92, 1956.
44. CLOUDSLEY-THOMPSON, J.L.: On the Arachnid fauna of Fair Isle - Ann.Mag.Nat.Hist., 12, 9, 830, 1956.
45. CLOUDSLEY-THOMPSON, J.L.: Studies in diurnal rhythms. V. Nocturnal ecology and water-relations of the British cribellate spiders of the genus *Ciniflo* BL. - Linnean Society's Journal Zoology, 43, 290, 1957.
46. CLOUDSLEY-THOMPSON, J.L.: Some comments on the natural control of animal populations with especial reference to insects - The Entomologist, 90, 1131, 195-204, 1957.
47. CLOUDSLEY-THOMPSON, J.L.: Institute of Biology Conversazione - Nature, 180, 319-320, 1957.
48. CLOUDSLEY-THOMPSON, J.L.: Studies in diurnal rhythms. Diurnal Rhythms in terrestrial Arthropoda - Acta Med. Scan. Suppl. (in press).
49. CLOUDSLEY-THOMPSON, J.L.: Studies in diurnal rhythms. VIII. The endogenous chronometer in *Gryllus campestris* L. (Orthoptera: Gryllidae) - J. Ins.Physiol., 2, 275-280, 1958.

50. CLOUDSLEY-THOMPSON, J.L.: Water Relations and Diurnal Rhythms in Woodlice - The Ann. of Appl. Biol., 46, 1, 117-119, 1958.
51. CLOUDSLEY-THOMPSON, J.L. and SANKEY, J.H.P.: Some aspects of the Fauna of the District around Juniper Hall, Mickleham, Surrey - IV - The Entomologist's Monthly Magazine, 94, 1958.
52. CLOUDSLEY-THOMPSON, J.L.: The evolution of twenty-four-hour periodicities in Terrestrial Arthropoda - Paper, XVth Int. Congress of Zoology, Sect. 10, Paper 21, 1958.
53. CLOUDSLEY-THOMPSON, J.L.: The effect of wind upon the nocturnal emergence of woodlice and other Terrestrial Arthropods I. - The Entomologist's Monthly Mag., 94, 106-108, 1958.
54. CLOUDSLEY-THOMPSON, J.L.: Observations on *Aepus Marinus* Stroem (Col.: Carabidae) - The Entomologist, 91, 1143, 169-170, 1958.
55. CLOUDSLEY-THOMPSON, J.L.: The behaviour of false-scorpions - The Naturalist, 96, 1958.
56. COLLINGWOOD, C.A.: Distribution of *Athous longicollis* and *Corymbites cupreus* (Col. Elateridae) in Warwickshire - M.A.F. Conference of Advisory Entomologists, Wireworm Survey, Notes no. , 1945.
57. COLLINGWOOD, C.A.: Carrot Fly Observations in Worcestershire - Worces. Agric. Chron., 16, 170-171, 1948.
58. COLLINGWOOD, C.A.: Ants in North Scotland - Ent. Rec., 61, 1949.
59. COLLINGWOOD, C.A.: Observations on Cabbage Stem Weevil - M.A.F. Conf. Adv. Ent. Occasional notes no. , 1949.
60. COLLINGWOOD, C.A.: *Formica sanguinea* Latr. in the Forest of Dean - Ent. Rec., 62, 91, 1950.
61. COLLINGWOOD, C.A.: Uncommon ant in Worcestershire - Ent. Mon. Mag., 86, 359, 1950.
62. COLLINGWOOD, C.A.: Distribution of ants in N.W. Scotland - Scot. Nat., 63, 45-49, 1951.
63. COLLINGWOOD, C.A.: Ants in Scotland - Ent. Rec., 63, 306-307, 1951.
64. COLLINGWOOD, C.A.: Insecticide Trials in West Midlands Orchards - Worces. Agric. Chron., 19, 297-301, 1951.
65. COLLINGWOOD, C.A., CROXALL, H.E. and JENKINS, J.E.E.: Observations on Brown Rot (*Sclerotinia laxa*) in relation to injury caused by Earwigs (*Froficula auricularia*) - Ann. Appl. Biol., 38, 833-842, 1951.
66. COLLINGWOOD, C.A.: Bionomics of *Metociscus paradoxus* - Ent. Rec., 65, 300, 1953.
67. COLLINGWOOD, C.A.: *Trichodes alveolaris* in England - Ent. Rec., 65, 301, 1953.
68. COLLINGWOOD, C.A.: Ant in Galloway - Ent. Rec., 65, 297-298, 1953.
69. COLLINGWOOD, C.A.: Ants in Berwickshire - Ent. Mon. Mag., 90, 23, 1954.
70. COLLINGWOOD, C.A.: Coleoptera in North Scotland - Ent. Rec., 66, 155-156, 1954.
71. COLLINGWOOD, C.A.: Uncommon Beetles in the West Midlands - Ent. Mon. Mag., 90, 197, 1954.
72. COLLINGWOOD, C.A.: Ants in a Country Garden - Ent. Mon. Mag., 90, 43, 1954.
73. COLLINGWOOD, C.A.: Ants in Dorset - Ent. Mon. Mag., 90, 43-44, 1954.
74. COLLINGWOOD, C.A.: The Hop Root Weevil - Plant. Path., 3, 63-65, 1954.
75. COLLINGWOOD, C.A.: Biology of *Epipolaeus caliginosus* F. (Col. Curculionidae) - Ent. Mon. Mag., 90, 169-172, 1954.
76. COLLINGWOOD, C.A. and CROXALL, H.E.: Carrot Fly Injury and "Canker" of Parsnips in the West Midlands - Plant Path., 3, 1954.
77. COLLINGWOOD, C.A.: Ants in S.W. Scotland - Ent. Rec., 67, 11-12, 1955.
78. COLLINGWOOD, C.A.: Ants in the S. Midlands - Ent. Gaz., 6, 143-149, 1955.
79. COLLINGWOOD, C.A. and SATCHELL, J.E.: Wood ants of the English Lake District - N.W. Nat., 1955.
80. COLLINGWOOD, C.A.: Ant hunting in France - Entomologist, 89, 105-108, 1956.
81. COLLINGWOOD, C.A.: A rare parasitic ant in France - Ent. Mon. Mag., 92, 197, 1956.
82. COLLINGWOOD, C.A. and SATCHELL, J.E.: Ants of the South Lake District - Journ. Soc. Brit. Ent., 5, 159-164, 1956.
83. COLLINGWOOD, C.A.: Aberrations in British ants of the genus *Formica* - Journ. Soc. Brit. Ent., 5, 193-196, 1956.
84. COLLINGWOOD, C.A.: *Formica cunicularia* Latr. (Hym. Formicidae) in the East Midlands - Ent. Mon. Mag., 92, 336, 1956.
85. COLLINGWOOD, C.A.: Distribution of ants allied to *Formica fusca* L. and *F. rufa* L. in Britain - Entomologist, 89, 291-294, 1956.
86. COLLINGWOOD, C.A.: The Glasshouse Red Spider Mite on Blackcurrants - Ann. Appl. Biol., 43, 144-147, 1956.
87. COLLINGWOOD, C.A.: Entomological Bioclimatology - See: Areal Report Dr. J.L. Cloudsley-Thompson. Symposium Int. Soc. Bioclim. and Biomet., Paris 29 - 31 August 1956.
88. COLLINGWOOD, C.A.: Coleoptera in N.W. Scotland - Ent. Rec., 69, 37-39, 1957.
89. COLLINGWOOD, C.A.: Myrmecophilous Beetles in the Midlands - Ent. Rec., 69, 9-14, 1957.
90. COLLINGWOOD, C.A.: *Myrmecichneumon subterraneus* Chev. (Col. Colydiidae) in Derbyshire - Ent. Mon. Mag., 93, 1957.

91. COLLINGWOOD, C.A. : A Collection of Ants in the Leicester City Museum - Ent. Rec. 69, 1957.
92. COLLINGWOOD, C.A. : British ants of the genus *Lasius* - Journ. Soc. Brit. Ent., 1957.
93. COLLINGWOOD, C.A. : A Survey of Irish Formicidae - 1957
94. COLLINGWOOD, C.A. and BROCK, A.M.: Spray Timing in Blackcurrants - Plant Path., 7, 1957.
95. COLLINGWOOD, C.A. and BRANGHAM, A.N.: Some ants of the Alpes Maritimes (in French) - Riviera Nat., 1957.
96. CORBET, P.S. : Seasonal regulation in British dragonflies - Nature, London, 174, 655, 1954.
97. CORBET, P.S. : A critical response to changing length of day in an insect - Nature, London, 175, 338, 1955.
98. CORBET, P.S. and TJØNNELAND, A.: Rhythmic flight activity of certain East African Thrichoptera - Nature London, 175, 1122, 1955.
99. CORBET, P.S. and TJØNNELAND, A.: The flight activity of twelve species of East African Thrichoptera - Univ. Bergen Arbok, 9, 1-49, 1955.
100. CORBET, P.S. : Entomological Bioclimatology - See Areal report Dr. J.L. Cloudsley-Thompson. Symposium Int. Soc. Bioclim. and Biomet., Paris, 29 - 31 August, 1956.
101. CORBET, P.S. : Environmental factors influencing the induction and termination of diapause in the Emperor Dragonfly, *Anax imperator* Leach (Odonata: Aeshnidae) - J. Exp. Biol. 1956. (in press).
102. CORBET, P.S. : The influence of temperature on diapause development in the dragonfly. *Lestes sponsa* (Hansemann) (Odonata: Lestidae) - Proc. R. Ent. Soc. Lond. (A), 1956 (in press).
103. CORBET, P.S. : The life-history of the Emperor Dragonfly, *Anax imperator* Leach, (Odonata: Aeshnidae) - J. Anim. Ecol. 1956 (in press).
104. CORBET, P.S. : The life-histories of two summer species of dragonfly (Odonata: Coenagruidae) - Ent. Gaz. 1956 (in press).
105. CORBET, P.S. : The life-histories of two spring species of dragonfly (Odonata: Zygoptera) - Ent. Gaz. 1956 (in press).
106. DAVEY, J.T. : A preliminary note on seasonal movements of the African migratory locust in the solitary phase - Locusta, 3, 1955.
107. DAVEY, J.T. : A method of marking isolated adult locusts in large numbers as an aid to the study of their seasonal migrations - Bull. of Entomological Research, 46, 4, 1956.
108. DAVEY, J.T. : The African Migratory Locust in the Central Niger Delta. I. Climate and Vegetation - Locusta, 5, 1957.
109. DAVIDSON, J. : Climate in relation to insect ecology in Australia. I. Mean monthly precipitation and atmospheric saturation deficit in Australia - Trans. Roy. Soc. of S. Aust., 58, 197-210, 1934.
110. DAVIDSON, J. : Climate in relation to insect ecology in Australia. 2. Mean monthly temperature and precipitation-evaporation ratio - Trans. Roy. Soc. of S. Aust., 59, 107-124, 1935.
111. DAVIDSON, J. : Rainfall-evaporation ratio in relation to locust and grasshopper outbreaks - Nature, 136, 298, 1935.
112. DAVIDSON, J. : Some aspects of climate and insect ecology - Journ. of Aust. Inst. of Agric. Sci., 1, 105-108, 1935.
113. DAVIDSON, J. : Climate in relation to insect ecology in Australia. 3. Bioclimatic zones in Australia - Trans. Roy. Soc. of S. Aust., 60, 88-92, 1936.
114. DUBIEF, J. and FOUREY, A.: De l'influence des éléments météorologiques de l'Afrique équatoriale sur les vols de sauterelles en Algérie - 63ème Congrès des Soc. Sav., Alger., 1930.
115. DUFFEY, E. : Aerial dispersal in a known spider population - Journal of Animal Ecology, 1956.
116. DUFFEY, E. : Entomological Bioclimatology - See Areal report Dr. J.L. Cloudsley-Thompson. Symposium Int. Soc. Bioclim. and Biomet., Paris, 29 - 31 August 1956.
117. DUVDEVANI, S. : Hessian fly in Palestine - Hassadeh, 1939.
118. EDNEY, E.B. and KEY, K.H.L.: Precocious adults resulting from the omission of the fifth instar in *Locusta migratoria migratorioides* R. and F. - Proc. R. Ent. Soc. London (A), 11, 55-58, 1936.
119. EDNEY, E.B. : A study of locomotor activity in *Locusta migratoria migratorioides* by the autograph method - Bull. Ent. Res., 28, 243-278, 1937.
120. EDNEY, E.B. : Notes on the behaviour and reactions to certain stimuli of the freshwater Jellyfish *Limnocyclus rhodesiae* Boul., - Occ. Pap. Nat. Mus. S. Rhodesia, 8, 1-11, 1939.
121. EDNEY, E.B. : The Heteronychiainae (Family Chrysidiidae) of South Africa - Occ. Pap. Nat. Mus. S. Rhodesia, 9, 29-126, 1940.

122. EDNEY, E.B. : Laboratory studies on the bionomics of the rat fleas *Xenopsylla brasiliensis* and *X-cheopsis*. 1. Certain effects of light, temperature and humidity on the rate of development and on adult longevity - Bull.ent.Res. 35, 399-416, 1945.
123. EDNEY, E.B. : An apparatus for handling small living insects during microscopic examination - Bull. ent.Res., 37, 83-87, 1946.
124. EDNEY, E.B. : Laboratory studies on the bionomics of the rat fleas *Xenopsylla brasiliensis* and *X.-cheopsis*. II. Water relations during the cocoon period - Bull.ent.Res., 38, 263-280, 1947.
125. EDNEY, E.B. : Laboratory studies on the bionomics of the rat fleas *Xenopsylla brasiliensis* and *X.-cheopsis*. III. Further factors affecting adult longevity - Bull.ent.Res., 38, 389-404, 1947.
126. EDNEY, E.E. : The Holonychinae (Family Chrysidae) of South Africa. Part I. Tribes Pseudochrysidini Bischoff; Parnopini Aaron; Allocoeliini Mocsary - Occ. Pap. Nat. Mus. S. Rhodesia, 13, 168-205, 1948.
127. EDNEY, E.B. : The evaporation of water from woodlice - Nature, 164, 321, 1949.
128. EDNEY, E.B. : The evaporation of water from woodlice and the millipede *Glomeris* - J. Exp. Biol., 28, 91-115, 1951.
129. EDNEY, E.B. : The body temperature of woodlice - J. Exp. Biol., 28, 271-280, 1951.
130. EDNEY, E.B. : The holonychinae (Family Chrysidae) of South Africa. Part II. Chrysidea Bischoff, Gonochrysis Licht. and Holochrysis Licht. - Occ. Pap. Nat. Mus. S. Rhodesia, 17, 403-452, 1952.
131. EDNEY, E.B. and DAVIES, M.E. : The evaporation of water from spiders - J. Exp. Biol. 29, 571-182, 1952.
132. EDNEY, E.B. : The body temperature of arthropods - Nature, 170, 586-587, 1952.
133. EDNEY, E.B. : The woodlice of Great Britain and Ireland - Proc. Linn. Soc. London, 164, 49-98, 1953.
134. EDNEY, E.B. : The Holonychidae (Family Chrysidae) of South Africa. Part III. Trichrysis Licht. - Occ. Pap. Nat. Mus. S. Rhodesia, 18, 532-539, 1953.
135. EDNEY, E.B. : The temperature of woodlice in the sun - J. exp. Biol., 30, 331-349, 1953.
136. EDNEY, E.B. : A synopsis of the British woodlice - Linn. Soc. Synopses of the British Fauna, 9, 1-55, 1954.
137. EDNEY, E.B. : The Holonychinae (Family Chrysidae) of South Africa. Part IV. Tetrachrysis Licht. - Occ. Pap. Nat. Mus. S. Rhodesia, 19, 543-623, 1954.
138. EDNEY, E.B. : The Holonychinae (Family Chrysidae) of South Africa. Part V. Pentachrysis Licht. and Hexachrysis Licht. - Occ. Pap. Nat. Mus. S. Rhodesia, 19, 624-673, 1954.
139. EDNEY, E.B. : Woodlice and the land Habitat - Biol. Rev. 29, 189-219, 1954.
140. EDNEY, E.B. and SPENCER, J.O. : The absorption of water by woodlice - J. Exp. Biol., 31, 491-196, 1955.
141. EDNEY, E.B. and SPENCER, J.O. : Cutaneous respiration in woodlice - J. Exp. Biol., 32, 256-269, 1955.
142. EDNEY, E.B. : Hymenoptera: Chrysidae in "South African Animal Life" - Ed. Hanström, B., Brink, P and Rudebeck, G., 3, 415-419, 1956.
143. EDNEY, E.B. : The water relations of Arthropods - Monographs in Experimental Biol. Cambridge, 5, 109 and 6, 1957.
144. EVANS, J.W. : Thrips investigation. 1. Seasonal fluctuations in number of Thrips *imaginis* Bagnall and associated blossom thrips - C.S.I.R. Journal, 6, 1933.
145. EVANS, J.W. : Thrips investigation. 2. Some factors that regulate the abundance of thrips *imaginis* Bagnall - C.S.I.R. Journal, 7, 61-69, 1934.
146. GOESSWALD, K. : Ueber die Frassstatigkeit von Forstschädlingen unter dem Einfluss von Altersunterschieden und der Einwirkung verschiedener Temperatur und Luftfeuchtigkeit und ihre praktische und physiologische Bedeutung. 1. Untersuchung an *Dendrolimus pini* L. - Ztschr. f. angew. Entom., 21, 2, 183-207, 1934.
147. GOESSWALD, K. : Physiologische Untersuchungen über die Einwirkung ökologischer Faktoren, besonders Temperatur und Luftfeuchtigkeit, auf die Entwicklung von *Diprion (Lophyrus) pini* L. zur Feststellung der Ursachen des Massenwechsels - Zeitschr. für Angew. Entom., 22, 3, 331-384, 1935.
148. GOESSWALD, K. : Zur Frage nach der Abhängigkeit der Entwicklung des Kiefernswärmers *Sphinx pinastri* L. von Temperatur und Luftfeuchtigkeit - Ztschr. f. angew. Entom., 22, 4, 521-532, 1936.
149. GOESSWALD, K. : Ueber den Einfluss von verschiedener Temperatur und Luftfeuchtigkeit auf die Lebensäusserungen der Ameisen. I. Die Lebensdauer ökologisch verschiedener Ameisenarten unter dem Einfluss bestimmter Luftfeuchtigkeit und Temperatur - Ztschr.f.wiss.Zoologie, 151, 337-381, 1938.

150. GOESSWALD, K. : Ueber den Einfluss von verschiedener Temperatur und Luftfeuchtigkeit auf die Lebensäusserungen der Ameisen. II. Ueber den Feuchtigkeits-sinn ökologisch verschiedener Ameisenarten und seine Beziehungen zu Biotrop, Wohn- und Lebensweise - Zeitschr. f. wiss. Zoologie, 154, 247-344, 1941.
151. GOESSWALD, K. : Zur Ameisenfauna des Mittleren Maingebietes mit Bemerkungen über Veränderungen seit 25 Jahren - Zool. Jahrb., Abtl. Syst., 80, 5/6, 507-532, 1951.
152. GOETZ, B. : Beiträge zur Analyse des Mottenfluges bei den Traubenwicklern *Clysia ambiguella* und *Polychrosis botrana* - Wein und Rebe, 23, 207-228, 1941.
153. GOETZ, B. : Neue Apparate zum Studium der Insektenphysiologie - Die Umschau in Wissenschaft und Technik, 49, 779-781, 1941.
154. GOETZ, B. : Freiland- und Laboratoriumsuntersuchungen über Ausschlüpfen, Eiablage und Nahrungsaufnahme bei den Traubenwicklern *Clysia ambiguella* und *Polychrosis botrana* - Wein und Rebe, 25, 135 - 153, 1943.
155. GOETZ, B. : Freilanduntersuchungen über Ausschlüpfen, Begattung und Eiablage beim Springwurmwickler *Sparganothis pilleriana* - Anzeiger f. Schädlingskunde, 19, 61-70, 1943.
156. GOETZ, B. : Tageszeit und Isektenaktivität - Naturwissenschaftliche Rundschau, 2, 257-261, 1949.
157. GOETZ, B. : Der Einfluss von Tageszeit und Witterung auf Ausschlüpfen, Begattung und Eiablage der Springwurmwicklers *Sparganothis pilleriana* Schiff - Z.f. angew. Entomologie, 31, 261-274, 1950.
158. GOETZ, B. : Entomological bioclimatology - Areal report: Symposium Int. Soc. Bioclim. and Biomet. Paris, 29-31 August 1956.
159. GREEN, G.W., SULLIVAN, W.G. and WELLINGTON, C.R.: Polarized light and body temperature level as orientation factors in the light reactions of some hymenopterous and lepidopterous larvae - Canad. J. Zool. 29, 339-351, 1951
160. GREEN, G.W. and HENSON, W.R.: A new type of coprometer for laboratory and field use - Canad. Entomol., 85, 6, 227-230, 1953.
161. GREEN, G.W. : Some laboratory investigations of the light reactions of the larvae of *Neodiprion americanus banksianae* Roh. and N. Lecontei (Fitch) (Hymenoptera: Diprionidae) - Canad. Entomol., 86, 5, 207-222, 1954.
162. GREEN, G.W. : Humidity reactions and water balance of larvae of *Neodiprion americanus banksianae* Roh. and N. Lecontei (Fitch) (Hymenoptera: Diprionidae) - Canad. Entomol., 86, 6, 261-274, 1954.
163. GREEN, G.W. : The functions of the eyes and antennae in the orientation of adults of *Neodiprion lecontei* (Fitch) - Canad. Entomol., 86, 8, 371 - 376, 1954.
164. GREEN, G.W. and FREITAS, A.S. de: Frass-drop studies of larvae of *Neodiprion americanus banksianae* Roh. and N. lecontei (Fitch) (Hymenoptera: Diprionidae) - Canad. Entomol., 87, 10, 427-440, 1955.
165. GREEN, G.W. : Temperature relations of ant-lion larvae (Neuroptera: Myrmeleontidae) - Canad. Entomol., 87, 10, 441-459, 1955.
166. GREEN, G.W. : Bioclimatic studies of forest insects - Areal report: Symposium Int. Soc. of Bioclim. and Biomet., Paris 29-31 August 1956.
167. GREENBANK, D.O. : Bioclimatic studies on the spruce budworm within the green river project - Areal report: Symposium Int. Soc. of Bioclim. and Biomet., Paris 29-31 August 1956.
168. GREENBANK, D.O. : The role of climate and dispersal in the initiation of outbreaks of the spruce budworm in New Brunswick. Part I: The role of climate - Can. J. Zool. (In press)
169. GREENBANK, D.O. : The role of climate and dispersal in the initiation of outbreaks of the spruce budworm in New Brunswick. Part II.: The role of dispersal - Can. J. Zool. (In press).
170. GRIFFITHS, J.F. : Entomological Bioclimatology - See Areal report Dr. J.L.Cloudsley - Thompson. Symposium Int. Soc. of Bioclim. and Biomet., Paris, 29-31 August 1956.
171. HADDOW, A.J. : Entomological Bioclimatology - See Areal report Dr. J.L.Cloudsley - Thompson. Symposium Int. Soc. of Bioclim. and Biomet., Paris, 29-31 August 1956.
172. HAINE, E. : Die Fauna des Grundwassers von Bonn mit besonderer Berücksichtigung der Crustaceen - Dissertation der Mathematisch-Naturwissenschaftlichen Fakultät der Rheinischen Friedrich-Wilhelm Universität, Bonn, 1-144, 1946.
173. HAINE, E. : Zur Frage der Ueberwinterung von *Myzodes persicae* Sulz. an Sekundärwäldern. I. Das Vorkommen des grünen Pfirsichlaus an Wintergemüse der Kölner Bucht und ihrer Randgebiete im ausgehenden Winter 1948/49 - Anz. f. Schädlgk., 23, 81-86, 1950.

174. HAINE, E. : Kurzer Bericht über einen Goldafterbekämpfungsversuch - Anz.f.Schäd-
lgk., 23, 163-165, 1950.
175. HAINE, E. : Zur Frage der Ueberwinterung von *Myzodes persicae* Sulz. an Sekundär-
wirten. II. *Myzodes persicae* Sulz. und andere an Kartoffeln vorkom-
mende Aphiden in den Gewächshäusern von Bonn - Anz. f. Schädlgk., 24,
97-105, 1951.
176. HAINE, E. : Studien zur Biologie des Zuckerrübenrüsslers (*Tanymecus palliatus* F.)
I. Welche Pflanzen vermögen *Tanymecus Palliatus* F. als Nahrung zu
dienen? - Anz. f. Schädlgk., 25, 36-40, 1952.
177. HAINE, E. : Weitere Bekämpfungsversuche mit *Euproctis chrysorrhoea* L. und *Diprion*
sertifer Geoffr. - Anz.f. Schädlgk., 25, 129-132, 1952.
178. HAINE, E. : Studien und Experimente zur Frage des Populations und Massenwechsels
und des Flugverhaltens virusübertragender Blattläuse - Anz.f. Schäd-
lgk., 27, 55-59, 1954.
179. HAINE, E. : Biologisch-ökologische Studien an *Rhopalosiphoninus latysiphon* D. -
veröffentl. durch das Bundesministerium für Ernährung, Landwirtschaft
und Forsten in der Reihe: Landwirtschaft, angewandte Wissenschaft,
Bonn, 1-58, 1955.
180. HAINE, E. : Aphid take-off in controlled windspeeds - Nature, 175, 474, 1955.
181. HAINE, E. : Zur Frage der Flugfähigkeit der Blattläuse: Beobachtungen über das
Flugstartverhalten und die Dauer der Flugfähigkeit der Männchen von
Periphyllus aceris acericola Wlk. - Anz.f. Schädlgk., 28, 67-70, 1955.
182. HAINE, E. : The flight activity of the sycamore aphid, *Drepanosiphon platanoides*
Schr. (Hemiptera Aphididae) - Journ.Anim. Ecol., 24, 388-394, 1955.
183. HAINE, E. : Häutung, Abflug und Landung der Blattläuse in Wechselwirkung auf die
Blattlauszahlen in der Luft - Mitt.Biol.Bundesanst., Berlin Dahlem
85, 1956.
184. HAINE, E. : Periodicity in aphid moulting and reproduction in constant tempera-
ture and light - Ztschr. Angew. Ent., 40, 1, 100-124, 1957.
185. HAINE, E., JOHNSON, C.G. and TAYLOR, L.R.: The analysis and reconstruction of diurnal flight
curves in alienicolae of *Aphis fabae* scop. - Ann.Appl.Biol., 45, 4,
682-701, 1957.
186. HAINE, E., JOHNSON, C.G., COCKBAIN, A.J. and TAYLOR, L.R.: Moulting rhythm in the alienico-
lae of *Aphis fabae* Scop. (Hemiptera: Aphididae) in the field - Ann.
Appl. Biol., 45, 4, 702-708, 1957.
187. HARVEY, G.T. : A relationship between Photoperiod and Cold-Storage Treatment in the
Spruce Budworm - Science, 128, 1205-1206, 1958.
188. HAUF, W.O. : Predicting mosquito emergence - Proc. of 40th Ann.Meeting of New
Jersey Mosquito Extermination Ass., Atlantic City, 52-57, 1953.
189. HAUF, W.O. : The effects of atmospheric pressure on the flight responses of *Aedes*
aegypti (L.) - Bull. of Ent. Research, 45, 3, 507-526, 1954.
190. HAUF, W.O. : The development of *Aedes* mosquitoes at Fort Churchill, Manitoba and
prediction of their emergence - Ecology
191. HAUF, W.O. : Physical environment and behaviour of the immature stages of *Aedes*
communis Deg. in subarctic. Canada - The Canadian Entomologist
192. HAUF, W.O. : Environmental Physiology of insects affecting man and livestock with
special reference to host parasite relationships - Areal report:Sym-
posium Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956.
193. HAUF, W.O. : Mosquito behaviour and weather: I. Relations of macrometeorological
factors to adult mosquito behaviour. II. Relations of acclimatic en-
vironment to mosquito activity. III. Relations of mosquito activity
to the human host under natural microclimatic conditions - In pre-
paration.
194. HENSON, W.R. : Entomological Bioclimatology - See Areal report Dr.D.H.K. Lee. Sym-
posium Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956.
195. HOSNY, M.M. : An attempt to forecast the average number of egg masses of the cotton
leaf worm, *Prodenia litura*, in Egypt in June by analysing preceeding
weather conditions - See Areal report Prof.L. Enberger. Symposium
Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956.
196. ISHIKURA and HIDETSUGU: On the types of the seasonal prevalence of rice stem borer moth in
Japan - Bull.Nat.Inst.Agric.Sci., C, 5, 67-80.
197. ISHIKURA and HIDETSUGU: Relation of the meteorological conditions to the occurrence of rice
stem borer - Studies of Agricultural Meteorol., 4, 36-106, 1948.
198. ISHIKURA and HIDETSUGU: Prediction of the occurrence of agricultural pest insects - Kawade
Pub. Co., Tokyo, 134 pp, 1949.
199. ISHIKURA and HIDETSUGU: Short review on the studies on the statistical prediction of the
date and number of rice borer moth appearance - Oyo-Dobutsu Zasshi
16, 1/2, 32-40, 1950.

200. JEFFREE, E.P. : A modified phenological approach in the determination of flowering times of beekeeping and other plants in Aberdeenshire - Proc.Intern. Beekeeping Congress Leamington Spa, Sept. 1951.
201. JEFFREE, E.P. : Acarine disease of the Honeybee and temperature - Nature, 175, 91, 1955.
202. JEFFREE, E.P. : Entomological Bioclimatology - See Areal report Dr. J.L. Cloudsley-Thompson, Symposium Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956.
203. JOHNSON, C.G. : Development, hatching and mortality of the eggs of *Cimex lectularius* L. (Hemiptera) in relation to climate, with observations on the effects of preconditioning to temperature - Parasitology, 2, 2, 127-173, 1940.
204. JOHNSON, C.G. : The longevity of the fasted Bed-bug (*C.lectularius* L.) under experimental conditions and particularly in relation to the saturation deficiency law of water-loss - Parasitology, 32, 3, 239-270, 1940.
205. JOHNSON, C.G. : The maintenance of high atmospheric humidities for entomological work with Glycerol-water mixtures - Ann. Appl. Biol., 27, 2, 295-299, 1940.
206. JOHNSON, C.G. : Insect survival in relation to the rate of water-loss - Biol.Rev. 17, 151-177, 1942.
207. JOHNSON, C.G. : The ecology of the Bed-bug, *Cimex lectularius* L., in Britain. Report on research 135-40 - J. Hyg. Camb., 41, 4, 345-456, 1942.
208. JOHNSON, C.G. and SOUTHWOOD, T.R.E.: Seasonal records in 1947 and 1948 of flying Hemiptera *Meteroptera* particularly *hygus pratensis* L., caught in nets 50 feet to 3000 feet above the ground - Proc.R. ent. Soc. Lond., A, 24, 128-130, 1949.
209. JOHNSON, C.G. : A suction trap for small airborne insects which automatically segregates the catch into successive hourly samples - Ann. Appl. Biol. 27, 80-91, 1950.
210. JOHNSON, C.G. : The infestation of a bean field by *aphis fabae* scop. in relation to wind direction - Ann. Appl. Biol., 37, 3, 441-450, 1950.
211. JOHNSON, C.G. : A comparison of suction trap, sticky trap and tow-net for estimating densities of small airborne insects - Ann. Appl. Biol., 37, 268-285, 1950.
212. JOHNSON, C.G. : The study of Wind-borne Insect populations in relation to terrestrial ecology. Flight periodicity and the estimation of aerial populations - Sc. Progress, 39, 153, 41-62, 1950.
213. JOHNSON, C.G. : Insect Aerobiology - New Biol. (enguin publication), 9, 1950.
214. JOHNSON, C.G. and EASTOP, V.F.: Aphids captured in a Rothamsted Suction Trap 5 feet above ground level, from June to November 1947 - Proc. R. ent.Soc. London, A, 26, 1-3, 17-24, 1950.
215. JOHNSON, C.G. and PENMAN, H.L.: Relationship of aphid density to altitude - Nature, 168, 337-338, 1950.
216. JOHNSON, C.G. : Changing number of *aphis fabae* flying at crop level in relation to current weather and population on the crop - A.A.B., Dec. 1952.
217. JOHNSON, C.G. : A new approach to the problems of the Spread of aphids and to insect trapping - Nature, 170, 147, 1952.
218. JOHNSON, C.G. : The rate of population level, flight periodicity and climate in the dispersal of aphids - Amsterdam, 1952.
219. JOHNSON, C.G. : The aerial dispersal of aphids - Discovery, Jan. 1953.
220. JOHNSON, C.G. : Aphid migration in relation to weather - Biol.Revs., 29, 87-118, 1954.
221. JOHNSON, C.G. and TAYLOR, C.E.: Wind direction and the infestation of bean fields by *aphis fabae* scop. - Ann. Appl. Biol., 41, 1, 107-116, 1954.
222. JOHNSON, C.G., SOUTHWOOD, T.R.E. and ENTWISTLE, H.: A method for sampling arthropods and molluscs from herbage by suction - Nature, 176, 559, 1955.
223. JOHNSON, C.G. and TAYLOR, L.R.: The measurement of insect density in the air - Lab.Practice 4, 5, 187-192 (pt.1) and 4, 6, 235-239 (pt 2), 1955.
224. JOHNSON, C.G. and TAYLOR, L.R.: The development of large suction traps for airborne insects - Ann. Appl. Biol., 43, 1, 52-61, 1955.
225. JOHNSON, C.G. : Entomological Bioclimatology - See Areal report Dr. J.L. Cloudsley-Thompson, Symposium Int. Soc. Bioclim. and Biomet. Paris, 29-31 August 1956.
226. KETTLE, D.S. : The effect of light on the biting activity of *chrysops silacea* (Diptera. Tabanidae) - Ann.trop.Med. Parasit., 47, 340-343, 1953.
227. KEVAN, D.K. McE. : Entomological Bioclimatology - See Areal report Dr. J.L. Cloudsley-Thompson, Symposium Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956.
228. KEVAN, D.K. McE. and SINGH, M.P.: Notes on three common British species of Agrotid Moth. I. Longevity and Oviposition - The Entomologist's Record, 68, 1956.

229. KEY, K.H.L. : An analysis of the outbreaks of the Australian Plague Locust (*Chortoicetes terminifera* Walk.) during the seasons 1937-38 and 1938-39 - Coun.Sci. Ind.Res. (Aust.) Bull., 146, 88, 1942.
230. KEY, K.H.L. : The outbreak of the Australian Plague Locust (*Chortoicetes terminifera* Walk.) in the season 1939-40, with special reference to the influence of climatic factors - Coun.Sci.Und.Res.(Aust) Bull., 160, 40, 1943.
231. KEY, K.H.L. : The general ecological characteristics of the outbreak areas and outbreak years of the Australian Plague Locust (*Chortoicetes terminifera* Walk.) - Coun.Sci.Ind.Res.(Aust) Bull., 186, 127, 1945.
232. KEY, K.H.L. : The taxonomy, phases and distribution of the genera *Chortoicetes* Brunn, and *Austroicetes* Uv. (Orthoptera: Acrididae) Canberra - Comm. Sci.Ind.Res.Org., 237, 1954.
233. KNOCH, K. : Die meteorologischen Verhältnisse bei dem Massenaufreten der Kriebelmücke - Hygienische Rundschau, 257-261, 289-293, 1921. Auszug: Das Wetter, 113-114, 1921.
234. KNOCH, K. : Die meteorologischen Verhältnisse an den Tagen mit Kriebelmückenschäden der Jahre 1921 und 1922 - Berliner Tierärztliche Wochenschrift, 39, 433-434, 1923.
235. KOCH, H.G. : Zur Phänologie der Bienenvölker - Zschr. f. Angewandte Meteorologie, (in press).
236. LARSEN, E.B. : The influence of temperature on the development of some species of Diptera - 1940.
237. LARSEN, E.B. : The importance of master factors for the activity of noctuids - 1943.
238. LARSEN, E.B. : The influence of the severe winters of 1939-42 on the soil fauna of Tipperne - 1949.
239. LARSEN, E.B. : Activity and migration of *Plusia gamma* L. - 1949.
240. LARSEN, E.B. : Observations on the activity of some Culicids - 1949.
241. MACFADYEN, A. : Entomological Bioclimatology - See Areal report Dr. J.L. Cloudsley-Thompson. Symposium Int. Sec. Bioclim. and Biomet., Paris, 29-31 August 1956.
242. MACLEOD, J. : The bionomics of *Ixodes ricinus* L. the sheep tick of Scotland - Parasitology, 24, 382-400, 1932.
243. MACLEOD, J. : Control of the sheep tick on hill pastures. A review of the possibilities, with some experimental data - Trans. Highl. Agric.Soc.Scot., 45, 1933.
244. MACLEOD, J. : *Ixodes ricinus* in relation to its physical environment: the influence of climate on development - Parasitology, 26, 282-305, 1934.
245. MACLEOD, J. : *Ixodes ricinus* in relation to its physical environment. II. The factors governing survival and activity - Parasitology, 27, 123-144, 1935.
246. MACLEOD, J. : *Ixodes ricinus* in relation to its physical environment. III. Climate and reproduction - Parasitology, 27, 489-500, 1935.
247. MACLEOD, J. : *Ixodes ricinus* in relation to its physical environment. IV. An analysis of the ecological complexes controlling distribution and activities - Parasitology, 28, 295-319, 1936.
248. MACLEOD, J. : The sheep tick (*Ixodes ricinus*) and its control in Great Britain - Cooper Tech. Bull, 2, 27, 1938.
249. MACLEOD, J. : The tick problem - Vet.Rec., 50, 1245-1250, 1938.
250. MACLEOD, J. : The seasonal and annual incidence of the sheep tick, *Ixodes ricinus*, in Britain - Bull. ent. Res., 30, 103-118, 1939.
251. MACLEOD, J. : The moisture balance of the living fleece in relation to blowfly myiasis, with some remarks on the Australian and British conceptions of susceptibility - Ann. Appl. Biol., 27, 379-392, 1940.
252. MACLEOD, J. : Recent work on the sheep tick and its bearing on control measures. Summary of paper to the Association of Applied Biologists, 14/2/'41. - Ann. Appl. Biol., 28, 296-297, 1941.
253. MACLEOD, J. and CRAUFURD-BENSON, H.J.: Observations on natural populations of the body louse *Pediculus humanus corporis* de Geer - Parasitology, 33, 278-299, 1941.
254. MACLEOD, J. : A survey of British sheep blowflies - Bull ent.Res., 34, 65-88, 1943.
255. MACLEOD, J. : A survey of British sheep blowflies. II. Relation of strike to host and edaphic factors - Bull.ent.Res., 34, 95-111, 1943.
256. MACLEOD, J. : The climatology of Blowfly Myiasis. I. Weather and Oviposition - Bull. ent.Res., 38, 285-303, 1947.
257. MACLEOD, J. : The distribution and dynamics of Ked populations, *Melophagus ovinus* Linn., Parasitology, 39, 61-68, 1948.
258. MACLEOD, J. : The climatology of Blowfly Myiasis. II. Oviposition and Daily Weather Indices - Bull. ent. Res., 40, 179-201, 1949.

259. McFARLANE, W.V. : Life cycle of *Coitocaicum anaspidis* Hickman a N.Z. digenetic trematode - Parasitology, 31, 172-183, 1939.
260. McFARLANE, W.V. : Blowfly strike in Marlborough, New Zealand - N.Z.J. Sci. and Techn., 23, 205A-213A, 1941.
261. McFARLANE, W.V. : The life cycle of the heterophyoid trematode *Telogaster opisthorchis* ng. nsp., T.R.S. N.Z., 75, 218-230, 1945.
262. McFARLANE, W.V. : Bionomics of two trematode parasites of N.Z. eels - J. Parasitol., 38, 39-397, 1952.
263. MUELLER, H.J. : Die intrazelluläre Symbiose bei *Fulgera europaea* und *Gixius nervosus* als Beispiele polysymbionter Zyklen - Verh. VII Internat. Kongr. für Entomol. Berlin 1938, 2, 877-894, 1939.
264. MUELLER, H.J. : Beiträge zur Biologie des Rapsglanzkäfers (*Meligethes aeneus* F.) - Z. Pflkrkh, 51, 9, 385-435, 1941.
265. MUELLER, H.J. : Weitere Beiträge zur Biologie des Rapsglanzkäfers, *Meligethes aeneus* F. (Winterlager und Massenbewegung im Frühjahr) - Z. Pflkrkh, 51, 12, 529-595, 1941.
266. MUELLER, H.J. : Bau und Funktion des Legeapparates der Zikaden - Z. Morph. u. Oekol., Tiere, 38, 534-629, 1942.
267. MUELLER, H.J. : Gestaltende Einflüsse des tierischen Organismus auf symbiontische Bakterien - Forsch. u. Fortsch., 18, 193-197, 1942.
268. MUELLER, H.J. : Saisondimorphismus bei Arten der Gattung *Euscelis* BRULLE (Homoptera Euscelidae) - Vorläufige Mitteilung *Bombus* 40, Hamburg, 1947.
269. MUELLER, H.J. : Zur Systematik und Phylogenie der Zikaden-Endosymbiosen - Biol.Zbl., 68, 343-368, 1949.
270. MUELLER, H.J. and EICHLER, Wd.: Erdflöhschäden in Sachsen-Anhalt 1948 - Nachr.bl.f.d.Dtsch. Pflschd., 3, 15-19, 1949.
271. MUELLER, H.J. : Ueber die Bedeutung der Winterwirte für die Bekämpfung der Schwarzen Bohnenlaus (*Doralis fabae* SCOP) - Nachr.bl. f.d. Dtsch. Pflschd., 5, 111-115, 1951.
272. MUELLER, H.J. : Ueber das Schlüpfen der Zikaden aus dem Ei - Zoologica, 103, 41, 53, 1951. (Abbild. auf 14 Taf.)
273. MUELLER, H.J. : Ueber die Ursachen der unterschiedlichen Resistenz von *Vicia faba* L., gegenüber der Bohnenblattlaus *Doralis fabae* SCOP. .III. Ueber das Wirtswahlvermögen der Schwarzen Bohnenblattlaus *Doralis fabae* SCOP.-Züchter, 21, 161-179, 1951.
274. MUELLER, H.J. : Ueber die intrazelluläre Symbiose der Peloridiide *Hemiodoeus fidelis* EVANS - Zool.Anz., 146, 150-167, 1951.
275. MUELLER, H.J. and UNGER, K.: Ueber die Ursachen der unterschiedlichen Resistenz von *Vicia faba* L. gegenüber der Bohnenblattlaus *Doralis fabae* SCOP. I. Der Massenwechsel von *Doralis fabae* SCOP in Abhängigkeit vom Witterungsverlauf 1949 in Quedlinburg - Züchter, 21, 1-30, 1951.
276. MUELLER, H.J. and UNGER, K.: Ueber die Ursachen der unterschiedlichen Resistenz von *Vicia faba* L. gegenüber der Bohnenblattlaus *Doralis fabae* SCOP. II. Ueber die Fluggewohnheiten, insbesondere das sommerliche Schwärmen, von *Doralis fabae* SCOP. und ihre Abhängigkeit von Tagesgang der Witterungsfaktoren - Züchter, 21, 76-79, 1951.
277. MUELLER, H.J. and UNGER, K.: Ueber den Einfluss von Licht, Wind und Temperatur auf den Befallsflug der Aphiden *Doralis fabae* SCOP. und *Myzodes persicae* SULZ sowie der Psyllide *Trioza nigricornis* FRST - Züchter, 22, 206 - 228, 1952.
278. MUELLER, H.J. : Schwarzstirn- und Rotkopfwürger im nordöstlichen Harzvorland - Beiträge zur Vogelkunde, 3, 138-145, 1953.
279. MUELLER, H.J. : Ueber die Ursachen der unterschiedlichen Resistenz von *Vicia fabae* L. gegenüber der Bohnenblattlaus *Doralis fabae* SCOP. IV. Das Zustandekommen des unterschiedlichen Initialbefalls - Züchter, 23, 176-189, 1953.
280. MUELLER, H.J. : Der Blattlausbefallsflug im Bereich eines Ackerbohnen- und eines Kartoffelbestandes - Beiträge z. Entomol., 3, 229-258, 1953.
281. MUELLER, H.J. and UNGER, K.: Studien über die Bedeutung von Witterung und Mikroklima für den Massenwechsel der Schwarzen Bohnenlaus (*Doralis fabae* SCOP) - Angew. Meteorolog. 1, 257-275, 1953.
282. MUELLER, H.J. : Der Saisondimorphismus bei Zikaden der Gattung *Euscelis* BRULLE (Homoptera Auchenorrhyncha) 3. Beitrag zur Biologie mitteleuropäischer Zikaden - Beiträge z. Entomol., 4, 1-56, 1954.
283. MUELLER, H.J. : Ueber den Einfluss der Tageslänge auf die Saisonformenprägung von *Euscelis plebejus* FALL. (Homoptera Auchenorrhyncha) - Verh. Dtsch. Zool. Ges. Tübingen, 1954, 307-316, 1955.

284. MUELLER, H.J. : Ueber die Massen- und Pertelanzucht der Jasside *Euscelis plebejus* Fall. (Homoptera Auchenorrhyncha) sowie anderer Pflanzensaftsauger - Beiträge z. Entomol., 4, 546-552- 1954.
285. MUELLER, H.J. and UNGER, K.: Ueber die Wirkung geländeklimatisch unterschiedlicher Standorte auf den Blattlaus-Befallsflug - Züchter, 24, 337-345, 1954.
286. MUELLER, H.J. : Ueber das Zahlenverhältnis der Geschlechter bei der Jasside *Strep-tanus marginatus* KBM - Wiss.Z. Univ.Halle, Math.-Nat., 4, 27 - 30, 1955.
287. MUELLER, H.J. : Die Bedeutung der Tageslänge für die Saisonformenbildung der Insekten, insbesondere bei den Zikaden - Ber. 7 Wanderversa.Dtsch. Entomol. Berlin 1954, 102-120, 1955.
288. MUELLER, H.J. : Zur Frage der Bruchus-Resistenz von Ackerbohnen - Z. Pflkrkh., 62, 598-602, 1955.
289. MUELLER, H.J. : Ueber die Vorflugzeit von *Doralis fabae* SCOP. und ihre Bedeutung für den Massenwechsel - Z. angew. Entomol., 38, 82-96, 1955.
290. MUELLER, H.J. : Die Saisonformenbildung von *Araschnia levana*, ein photoperiodisch gesteuerter Diapause-Effekt - Naturwiss., 42, 134-135, 1955.
291. MUELLER, H.J. and UNGER, K.: Ueber die Bedeutung der Zusammenhänge zwischen Witterung und Blattlausflug für die Probleme des Kartoffelabbaus - Forsch.und Fortschr., 29, 229-238, 1955.
292. MUELLER, H.J. : Experimentelle Studien an der Symbiose von *Coptosoma scutellatum* GEOFFR. (Hem.Heteropt.) - Z. Morphol. und Oekol. Tiere, 44,459-482, 1956.
293. MUELLER, H.J. : Zikaden, Aleurodiden, Psylliden. In SORAURER - Handbuch der Pflanzenkrankheiten, 5, 5.Aufl., 3 Lief., 150-359, 1956.
294. MUELLER, H.J. : Die Wirkung verschiedener diurnaler Licht-Dunkel Relationen auf die Saisonformenbildung von *Araschnia levana* - Die Naturwissenschaften, 43, 21, 503-504, 1956.
295. MUELLER, H.J. : Ueber die Wirkung von Umweltfaktoren auf die Variabilität saison-dimorpher Insekten, insbesondere der Gattung *Euscelis* - Verh. Dtsch. Zool.Ges. Hamburg, 1956 (in press).
296. MUELLER, H.J. : The taxonomic value of the male genitalia in the light of new studies on the seasonal forms of *Euscelis* - Verh. X Intern. Kongr. d. Entomol. Montreal 1956 (in press).
297. MUELLER, H.J. : Die Wirkung exogener Faktoren auf die zyklische Formenbildung der Insekten, insbesondere der Gattung *Euscelis* (Hom.Auchenorrhyncha) - 5. Beitrag (Habilitationsschrift in press).
298. MUELLER, H.J. : Ueber die Diapause von *Stenocranus minutus* FABR. (Homoptera Auchenorrhyncha) - Beiträge z. Entomol. (in press).
299. MUELLER, H.J. : Zur Problematik der Blattlausresistenz landwirtschaftlicher Kulturpflanzen - Sitzungsberichte der Dtsch. Akad. d. Landwirtschaftswissenschaften (in press).
300. MUELLER, H.J. : Ueber die Entwicklung erhöhten Randbefalls von Ackerbohnenbeständen durch *Aphis fabae* SCOP. - Zschr. f. Pflanzenkrankheiten und Pflanzenschutz, 64, 7/10, 593-599, 1957.
301. MUELLER, H.J. : Ueber die Ursachen der unterschiedlichen Resistenz von *Vicia Faba* L. gegenüber der Bohnenblattlaus, *Aphis* (*Doralis*) *Fabae* SCOP. V. Antibiotische Wirkungen auf die Vermehrungskraft - Ent.Exp. & Appl., 1, 181-190, 1958. (North Holland Publishing Co., Amsterdam).
302. MUELLER, H.J. : Ueber den Einfluss der Photoperiode auf Diapause und Körpergrösse der Delphacide *Stenocranus minutus* FABR. (Homoptera Auchenorrhyncha) - Zoologischer Anzeiger, 160, 11/12, 294-312, 1958.
303. NOMURA, K. : On the locality of the outbreak type of agricultural insect pests and climatic zone in Japan - Sci. Bull., Faculty of Agr., Kyushu Univ., 13, 1-4, 1951.
304. NOMURA, K. : The relationship between agricultural insect pest infestation and climate in Japan - Technical Bull. of Faculty of Horticulture, Chiba Univ., 1, 1953.
305. NØRGAARD, E. : On the biology of *Eresus niger*. - Ent.Med., 22, 1941.
306. NØRGAARD, E. : Økologiske undersøgelser over nogle danske jagtedderkopper (Summary: Ecological investigations on some Danish Lycosid spiders) - Flora og Fauna, 1945.
307. NØRGAARD, E. : Bidrag til danske edderkoppers biologi. I. *Lithyphantes albomaculatus*. (Summary: Contributions to the biology of Danish spiders. I. *Lithyphantes albomaculatus*) - Flora og Fauna, 1948.
308. NØRGAARD, E. : Notes on the biology of *Filistata insidiatrix* - Ent.Med. 26, 1951.
309. NØRGAARD, E. : On the ecology of two Lycosid spiders (*P. piraticus* and *L. pullata*) from a Danish sphagnum bog. - Oikos, 3, 1951.

310. PLATT, R.B. : A multiple factor environmental gradient control chamber for mosquito behavior reactions - JI. of the Tennessee Ac. of Science, 29, 184-185, 1954.
311. PLATT, R.B. and WHITHERSPOON, J.P.: Responses of females of *Anopheles quadrimaculatus* Say. to controlled gradients of light, temperature and humidity - JI. of the Tennessee Academy of Science, 187-188, 1954.
312. PLATT, R.B. : A field laboratory for the study of the natural behavior of adult mosquitoes in time along both vertical and horizontal gradients with respect to precisely determined microenvironmental conditions - J. of American Society of Tropical Medicine and Hyg., 5, 390, 1956.
313. PLATT, R.B. and COLLINS, C.L.: Temperature tolerances and behavior patterns of adult *Anopheles quadrimaculatus* (Say) populations by age and sex in moist and dry conditions, with ecological implications - JI. of American Soc. of Tropical Med. and Hyg., 5, 378, 1956.
314. PLATT, R.B., COLLINS, C.L. and WITHERSPOON, J.P.: Reactions of *Anopheles quadrimaculatus* Say to Moisture, Temperature and Light - Ecological Monographs, 27, 303-324. 1957.
315. RAINY, R.C. and WALEFF, Z.: Desert locust migrations and synethic meteorology in the Gulf of Aden area - J. of Animal Ecology, Cambridge, 17, 101-112, 1948.
316. RAINY, R.C. : Weather and the movements of locust swarms: a new hypothesis - Nature 168, 1057, 1951.
317. RAINY, R.C. and WALEFF, Z.: Flying locusts and convection currents - Anti-Locust Bulletin London, 9, 51-70, 1951.
318. RAINY, R.C. : Desert locust swarm movements and weather - Proc. R. Ent. Soc. Lond. (c) 17, 45-46 and 56-57, 1952.
319. RAINY, R.C. and ASHALL, C.: Note on the behaviour of desert locusts in a light-beam - Brit. J. Anim. Behaviour, Huntingdon, 1, 136-138, 1953.
320. RAINY, R.C. : Entomological Bioclimatology - See Areal report Dr. J.L. Cloudsley-Thompson. Symposium Int. Soc. Bioclim. and Biomet. Paris 29-31 August 1956.
321. RAMDAS, L.A. : Meteorological assistance in locust control - Report of the chairman of CAGM working group of the commission for Agricultural Meteorology, 2nd session, Warsaw, 1958, World Met.Org., no. CAGM-II/Doc.7.
322. REMAUDIERE, G. : Etude Ecologique de *Locusta Migratoria Migratorioides* Rech. et Frm. (Orth. Acrididae) dans la zone d'Inondation du Niger en 1950 - Locusta, 2, 1954.
323. RITCHIE, J.Y. : Entomological Bioclimatology - See Areal report Dr. J.L. Cloudsley-Thompson. Symposium Int. Soc. Bioclim. and Biomet. Paris, 29-31 August 1956.
324. ROSE, A.H. and BLAIS, J.R.: A relation between April and May temperatures and spruce budworm larval emergence - Canad. Ent. 86, 4, 174-177, 1954.
325. SANKEY, J.H.P. : Further records of british harvestspiders (Arachnida, opiliones) with a note on *Nelima fuscifrons* (simon), possibly new to Britain - The Entomologist, 86, 116-117, 1953.
326. SANKEY, J.H.P. and CHAPMAN, R.F.: The larger invertebrate fauna of three rabbit carcasses - The Journal of Animal Ecology, 24, 2, 395-402, 1955.
327. SANKEY, J.H.P. : Entomological Bioclimatology - See Areal report Dr. J.L. Cloudsley-Thompson. Symposium Int. Soc. Bioclim. and Biomet., Paris, 29-31 August 1956.
328. SCHUA, L. : Der Einfluss des Wetters auf das Verhalten der Honigbienen - Verhandl. d. dtsh. Zool. Marburg, 1950.
329. SCHUA, L. : Untersuchungen über den Einfluss meteorologischer Elemente auf das Verhalten der Honigbienen - Ztschr.f. vergl. Physiol., 34, 258, 1952.
330. SCHUA, L. : Stehen Erkrankungen und Totenfall der Bienen im Winter im Zusammenhang mit Witterungsfaktoren ? - Bull. 1 Sec.Fr. d'Union International pour l'étude d. ins. soc., 1955.
331. SHEPHERD, R.F. and HENSON, W.R.: Effects of radiation on the habitat temperatures of the lodgepole needle miner *Recurvata milleri* Busck (Gelechiidae Lepidoptera) - Can. Journal Zool., 30, 144-153, 1952.
332. SHEPHERD, R.F. : The influence of the climate on the behaviour and survival of forest insects - Areal report : Symposium Int. Soc. Bioclim and Biomet., Paris, 29 - 31 August 1956.
333. SIDDORN, J.W. : Entomological Bioclimatology - See Areal report Dr. J.L. Cloudsley-Thompson. Symposium Int. Soc. Bioclim. and Biomet., Paris, 29-31 August 1956.
334. SLATER, A.E. : Insects and Weather - Quarterly Journal of the Royal Meteorological Society, 71, 309-310, 1945.

335. SMITH, R.F. : The importance of the microenvironment in Insect Ecology - Journal Econ. Ent., 47, 2, 205-210, 1954.
336. SOLOMON, M.E. : Experiments on the effects of temperature and humidity on the survival of *Halotydeus destructor* (Tucker) Acarina fam. Pentheleidae - Aust. J. Exp. Biol. Med. Sci., 15, 1-16, 1937.
337. SOLOMON, M.E. : Tyroglyphid mites in stored products - Ecological studies 1940. Ann. Appl. Biol., 33, 82-97, 1946.
338. SOLOMON, M.E. and ADAMSON, B.: The powers of survival of storage and domestic pests under winter conditions in Britain - Bull. Ent. Res., 46, 311-355, 1955.
339. SOLOMON, M.E. : Entomological Bioclimatology - See Areal report Dr. J.L. Cloudsley-Thompson. Symposium Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956.
340. STARK, R.W., HENSON, W.R. and WELLINGTON, W.G.: Effects of the weather of the coldest month on winter mortality of the lodgepole needle miner - Can. Ent., 86, 1, 13-19, 1954.
341. STARK, R.W. : The role of climate in the epidemiology of the lodgepole needle miner - Areal report: Symposium Int. Soc. Bioclim. and Biomet., Paris, 29-31 August 1956.
342. SULLIVAN, C.M., HENSON, W.R. and WELLINGTON, W.G.: Investigations in physical ecology - For. Insect Invest., Bimonthly Progress Rep., Ottawa, Can., 5, 5, 1949.
343. SULLIVAN, C.M., WELLINGTON, W.G. and GREEN, G.W.: Polarized light and body temperature level as orientation factors in the light reactions of some hymenopterous and lepidopterous larvae - Canad. J. Zool., 29, 339-359, 1951.
344. SULLIVAN, C.R. and WELLINGTON, W.G.: The light reactions of larvae of the tent caterpillars *Malacosoma disstria* Hbn., *M. americanum* (Fab) and *M. pluviale* (Dyar). (Lepidoptera: Lasiocampidae) - Canad. Entomol., 85, 8, 297-310, 1953.
345. SULLIVAN, C.R. : Use of radioactive cobalt in tracing the movements of the white-pine weevil *Pissodes strobi* Peck. (Coleoptera: Curculionidae) - Canad. Entomol., 85, 8, 273-276, 1953.
346. SULLIVAN, C.R., WELLINGTON, W.G. and HENSON, W.R.: The light reactions of larvae of the spotless fall webworm, *Hyphantria textor* Harr. (Lepidoptera: Arctiidae) - Canad. Entomol., 86, 12, 529-542, 1954.
347. SULLIVAN, C.R. : Bioclimatic studies on forest insects - Areal report: Symposium Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956.
348. SULLIVAN, C.R. and SIMPSON, S.D.: The x-radiation tolerance of the white pine weevil *Pissodes strobi* Peck (Coleoptera: Curculionidae) - Canad. Entomol., 1956.
349. SULLIVAN, C.R. and BELYEA, R.M.: The white pine weevil: a review of current knowledge - For. Chron., 1956.
350. TAYLOR, C.E. : Entomological Bioclimatology - See Areal report Dr. J.L. Cloudsley-Thompson. Symposium Int. Soc. Bioclim. and Biomet., Paris, 29-31 August 1956.
351. TJØNNELAND, A. and CORBET, P.S.: Rhythmic flight activity of certain East African Trichoptera - Nature, 175, 1955.
352. TJØNNELAND, A. and CORBET, P.S.: The flight activity of twelve species of East African Trichoptera - Univ. Bergen, Arb. Naturv. R., 9, 1955.
353. UNGER, K. and MUELLER, H.J.: Ueber die Fluggewohnheiten, besonders das sommerliche Schwärmen, von *Doralis fabae* und ihre Abhängigkeit vom Tagesgang der Witterungsfaktoren - Der Züchter, 21, 3, 31-44, 1951.
354. UNGER, K. and MUELLER, H.J.: Der Verlauf des Massenwechsels von *Doralis fabae* SCOP in Abhängigkeit vom Witterungsverlauf 1949 in Quedlinburg - Der Züchter, 21, 1, 1-30, 1951.
355. UNGER, K. and MUELLER, H.J.: Ueber den Einfluss von Licht, Wind, Temperatur und Luftfeuchtigkeit auf den Befallsflug der Aphiden *Doralis fabae* SCOP. und *Myzodes persicae* SULZ sowie der Psyllide *Trioza nigricornis* FRST. - Der Züchter, 22, 7/8, 206-228, 1952.
356. UNGER, K. und MUELLER, H.J.: Studien über die Bedeutung von Witterung und Mikroklima auf den Massenwechsel der Schwarzen Bohnenlaus (*Doralis fabae* SCOP) - Angew. Met., 1, 9, 257-275, 1953.
357. UNGER, K. and MUELLER, H.J.: Ueber die Wirkung geländeklimatisch unterschiedlicher Standorte auf den Blattlausbefallsflug - Der Züchter, 11/12, 337-345, 1954.
358. VERZAR, F. and LEVTWILER, M.: Notiz über die Wirkung von grossen Höhen auf die Entwicklung von *Axolotl*. (*Ambystoma*) und *Rana-fusca*-Larven und die *Thyreocida* der ersteren - Höhenklima-Forschung des Basler Physiologischen Inst. II, 1948.
359. WATERHOUSE, F.L. : Entomological Bioclimatology - See Areal report Dr. J.L. Cloudsley-Thompson. Symposium Int. Soc. Bioclim. and Biomet. Paris 29-31 August 1956.
360. WATERS, W.E. : Entomological Bioclimatology - See Areal report Dr. D.H.K. Lee. Symposium Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956.

361. WELLINGTON, W.G. : Barotaxis in Diptera, and its possible significance to economic entomology - *Nature*, 154, 671-672, 1944.
362. WELLINGTON, W.G. : The effect of ground temperature inversions upon the flight of *Culex* sp. - *Canadian Entomologist*, 76, 223, 1944.
363. WELLINGTON, W.G. : A laboratory method for determining the minimum active temperatures of insects - *Canadian Entomologist*, 77, 135-136, 1945.
364. WELLINGTON, W.G. : Conditions governing the distribution of insects in the free atmosphere. I. Atmospheric pressure, temperature and humidity - *Canadian Entomologist*, 77, 7-15, 1945.
365. WELLINGTON, W.G. : Conditions governing the distribution of insects in the free atmosphere. II. Surface and upper winds - *Canadian Entomologist*, 77, 21-28, 1945.
366. WELLINGTON, W.G. : Conditions governing the distribution of insects in the free atmosphere. III. Thermal convection - *Canadian Entomologist*, 77, 44-49, 1945.
367. WELLINGTON, W.G. : Conditions governing the distribution of insects in the free atmosphere. IV. Distributive processes of economic significance - *Canadian Entomologist*, 77, 69-74, 1945.
368. WELLINGTON, W.G. : The effects of variations in atmospheric pressure upon insects - *Canadian Journal of Research (D)*, 24, 51-70, 1946.
369. WELLINGTON, W.G. : Some reactions of muscoid Diptera to changes in atmospheric pressure - *Canadian Journal of Research (D)*, 24, 105-117, 1946.
370. WELLINGTON, W.G. and HENSON, W.R.: Notes on the effects of physical factors on the spruce budworm - *Canadian Entomologist*, 79, 168-170, 195, 1947.
371. WELLINGTON, W.G. : The light reactions of the spruce budworm - *Canadian Entomologist* 80, 56-82A, 1948.
372. WELLINGTON, W.G. : The spruce budworm in relation to flowering condition of the host tree: measurements of the physical environment - *Dominion Department of Agriculture Forest Insect Investigations Bi-monthly Progress Report*, 4, 5, 1-2, 1948.
373. WELLINGTON, W.G. : Temperature measurements in ecological entomology - *Nature*, 163, 614-615, 1949.
374. WELLINGTON, W.G. : The effects of temperature and moisture upon the behaviour of the spruce budworm. I. The relative importance of graded temperatures and rates of evaporation in producing aggregations of larvae - *Scientific Agriculture*, 29, 201-215, 1949.
375. WELLINGTON, W.G. : The effects of temperature and moisture upon the behaviour of the spruce budworm. II. The responses of larvae to gradients of evaporation - *Scientific Agriculture*, 29, 216-229, 1949.
376. WELLINGTON, W.G. : Effects of radiation on the temperatures of insectan habitats - *Scientific Agriculture*, 30, 209-234, 1950.
377. WELLINGTON, W.G. : Variations in the silkspinning and locomotor activities of larvae of the spruce budworm at different rates of evaporation - *Transactions Royal Society of Canada*, 44 (Sect. V), 89-101, 1950.
378. WELLINGTON, W.G., FETTES, J.J., TURNER, K.B. and BELYEA, B.M.: Physical and biological indicators of the development of outbreaks of the spruce budworm - *Canadian Journal of Research (D)*, 28, 308-331, 1950.
379. WELLINGTON, W.G., GREEN, G.W. and SULLIVAN, C.R.: Polarized light and body temperature level as orientation factors in the light reactions of some hymenopterous and lepidopterous larvae - *Can. Journal of Zoology*, 29, 339-351, 1951.
380. WELLINGTON, W.G. : Air-mass climatology of Ontario north of Lake Huron and Lake Superior before outbreaks of the spruce budworm and the forest tent caterpillar - *Canadian Journal of Zoology*, 30, 114-127, 1952.
381. WELLINGTON, W.G. : Motor responses evoked by the dorsal ocelli of *Sarcophaga aldrichi* Parker, and the orientation of the fly to plane polarized light - *Nature*, 172, 1177-1179, 1953.
382. WELLINGTON, W.G. and SULLIVAN, C.R.: The light reactions of Larvae of the Tent Caterpillars, *Malacosoma disstria* Hbn., *M. americanum* (Fab.) and *M. pluviale* (Dyar). (Lepidoptera: Lasioidae). - *The Canadian Entomologist*, 85, 8, 297-310, 1953.
383. WELLINGTON, W.G. : Atmospheric circulation processes and insect ecology - *Canadian Entomologist*, 86, 312-333, 1954.
384. WELLINGTON, W.G. : Weather and climate in forest entomology. In *Recent studies in bioclimatology - Meteorological Monographs*, 2, 8, 11-18, 1954.
385. WELLINGTON, W.G. : Pole blight and climate - *Department of Agriculture Forest Biology Division Bimonthly Progress Report*, 10, 6, 2-4, 1954.

386. WELLINGTON, W.G., STARK, R.W. and HENSON, W.R.: Effects of the weather of the coldest month on winter mortality of the lodgepole needle miner, *Recurvaria* sp., in Banff National Park - The Canadian Entomologist, 86, 13-19, 1954.
387. WELLINGTON, W.G., SULLIVAN, C.R. and HENSON, W.R.: The light reactions of larvae of the spotless fall webworm, *Hyphantria textor* Harr. (Lepidoptera: Arctiidae) - The Canadian Entomologist, 86, 12, 529-542, 1954.
388. WELLINGTON, W.G. : Solar heat and plane polarized light versus the light compass reaction in the orientation of insects on the ground - Annals Entomological Society of America, 48, 67-76, 1955.
389. WELLINGTON, W.G. : Bioclimatic studies on insects - Areal report: Symposium Int. Soc. Bioclim. and Biomet., Paris 29-31 August, 1956.
390. WELLINGTON, W.G. : Behaviour. pp. 312-313 In entomology in Canada up to 1956: a review of developments and accomplishments compiled by R.Glen - Canadian Entomologist, 88, 290-371, 1956.
391. WELLINGTON, W.G. : The synoptic approach to studies of insects and climate - Annual review of Entomology, 2, 143-162, 1957.
392. WELLINGTON, W.G. : Individual differences as a factor in population dynamics: the development of a problem - In press.
393. WYNNE-EDWARDS, V.C. : Entomological Bioclimatology - See Areal report Dr. J.L. Cloudsley-Thompson. Symposium Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956.
394. YAGI, N. : Sur la coefficient de température de la pulsation de Ver à soie - Archivio Zoologico Italiano, Padova, 1930.
395. YAGI, N. : Temperature characteristics of pulsation and growth in MOSQUITO PUPAE as basis for determination of lower developmental threshold point - Bull. Imp. Ag. Exp. St., 3, 319, 1932.
396. YAGI, N. : On the isophenological region of the rice stem borer - Bull. Imp. Agr. Exp. St., 2, 3, 1934.
397. YAGI, N. : On the nocturnal activity of the moth of rice stem borer - Bull. Imp. Agr. Exp. St., 2, 4, 1934.
398. YAGI, N. : Note on the reaction of CULEX PIPIENS PAILLENS - Proc. Imp. Acad., 18, 5, 1941.
399. YAGI, N. : On the spectral range for pigment migration of the compound eye of CHILO SIMPLEX Butler (a rice stem borer), - Proc. Imp. Acad., 18, 5, 1941.
400. YAGI, N. : On the origin and migration of guanophore pigments in the compound eye of the scarabaeid beetle, *Copris ochus* Motsch - Jap. Journ. Zool., 11, 2, 1954.
401. YAGI, N. : On the future distribution of Paddy Stem Borer - Oyo-kontyu, 11, 3, 1955.
402. YASUE, Y and HAMADA, A.: Climatic factors in relation to the geographical distribution of the potato lady beetles, *Epilachna vigintioctomaculata* and *Epilachna sparsa* in Okayama prefecture, Japan - Nogaku-Kenkyu, 41, 162-172, 1954.
403. YASUE, Y. : Bioclimatological studies on the geographical distribution of the potato lady beetles, *Epilachna vigintioctomaculata* and *Epilachna sparsa* in Chugoku District, the Western parts of Honshu, Japan - Setonaikai-Kenkyu (A journal of the society for the study of the Inland Sea), 8, 1956.
404. ZOLOTAREVSKY, B. : History of the international organisation for control of the African migratory locust - Locusta, 1, 1954.

VETERINARY BIOCLIMATOLOGY

1. BIANCA, W. : Der Einfluss der Alpeng auf das Blutvolumen von Jungziegen - Schweizer Archiv für Tierheilkunde, 93, 7, 468, 1951.
2. BIANCA, W. : Der Einfluss der Alpeng auf die Körperbeschaffenheit und das Blut von Jungziegen - Schriften der Schweiz. Vereinigung für Tierzucht, 15, 1951.
3. BIANCA, W. : Die Akklimatisation von Haustieren - Landwirtschaftliches Jahrbuch der Schweiz., 66, 1952.
4. BIANCA, W. : Klimatologische Untersuchungen im Offenstall - Zeitschrift für Tierzucht und Züchtungsbiologie, 61, 1953.
5. BIANCA, W. : Einige physiologische Reaktionen des Rindes auf atmosphärische Hitze - Schweizer Archiv. für Tierheilkunde, 95, 8, 451, 1953.
6. BIANCA, W. : Die Hitzeresistenz von Schweizer Braunvieh - Schweiz. Landwirtschaftliche Monatshefte, 32, 1954.
7. BIANCA, W. : The effect of heat stress on the acid-base balance of the calf - Jnl. of Agricultural Science, 45, 4, 428, 1955.

8. BIANCA, W. : The effect of short terme exposures to high environmental temperature on the volume and hydration of the blood of the calf - *Journal of Physiology* , 130, 17, 1955.
9. BIANCA, W. : The effect of repeated short exposures to heat on the volume and hydration of the blood of the calf - *British Veterinary Journal*, 113, 6, 1957.
10. CENA, M. : Studies on the sorbtive properties of the skin of domesticated animals - *Medycyna Weterynaryjna*, 3, 1949. Summary in English.
11. CENA, M. and COURVOISIER, P.: Untersuchungen über die physikalischen Faktoren des Stallklimas unter besonderer Berücksichtigung der Abkühlungsgrösse - *Schweiz. Archiv für Tierheilkunde*, 5, 1949.
12. CENA, M. and COURVOISIER, P.: Die Bedeutung der ultraroten Wärmestrahlung für die Abkühlungsgrösse im Stall - *Schweizer Archiv für Tierheilkunde*, 7, 1949.
13. CENA, M. : Studies on the photo climatic conditions in the breeding of domesticated animals - *Medycyna Weterynaryjna* 10, 1949. Summary in English.
14. CENA, M. and COURVOISIER, P.: Die Reflexionseigenschaften des Felles der Schweizer Braunvierasse - *Schweizer Archiv für Tierheilkunde*, 2, 1950. Summary in English, German and French.
15. CENA, M. : Pomiary temperatury skóry kotów przy pomocy termoelementu. Measurements of the cats' skin temperature with thermoelements - *Odbitka z "Medycyny weterynaryjnej"* 7, 1950.
16. CENA, M. : Methodische Beiträge zu fotoklimatische Untersuchungen in der Tierumwelt - *Przegląd Meteorologiczny i Hydrologiczny Rocznik 1950/1951*.
17. CENA, M. : Records of the cooling power in animals environment - *Medycyna Weterynaryjna*, 4, 1951.
18. CENA, M. and AUGUSTYN, J.: Bemerkungen zur Photophilie der Hühner - *Przegląd Hodowlany*, 9, 1951.
19. CENA, M. : Comparative investigations on the absorption of sunradiation through the skin of Swiss-brown breed of cattle - *Archiv. Weterynaryjne*, 1, 1952.
20. CENA, M. : Vergleichende Untersuchungen der physikalischen Faktoren des Klimas in der Tierstallungen - *Prace Wroclawskiego Towarzystwa Naukowego Seria B.*, 53, 1952. Summary in English.
21. CENA, M. : Investigations on the thermal protection of furs of living rabbits - *Medycyna Weterynaryjna* 7, 1952.
22. CENA, M. : Das Klima der Tierstallungen - *Medycyna Weterynaryjna*, 10, 1952.
23. CENA, M., CZYSZEK, W. and SLOMKA, J.: Investigations on the influence of the colour on the absorption of the sun radiation - *Medycyna Weterynaryjna*, 6, 1952.
24. CENA, M. and CZAJKOWSKI, Z.: Examinations of the chemical content of air in animal dwellings - *Postępy Wiedzy Rolniczej*, 2, 1953.
25. CENA, M. : Untersuchungen über den Einfluss des Lichtes auf das Hausgeflügel - *Gazeta Obserwatora PIHM*, 5, 1953.
26. CENA, M. and KRAUSS, S.: Ways of the spreading of foot and mouth disease in Poland on the background of the environment - *Medycyna Weterynaryjna* 4 i 5, 1954.
27. CENA, M. : Messungen der Verstaubung in der Haustierumwelt - *Gazeta Obserwatora PIHM*, 6, 1954.
28. CENA, M. : Messungen der Verdunstung im Haustierumwelt - *Gazeta Obserwatora PIHM* 6, 1954.
29. CENA, M. : Examination of the content of air in animal dwellings - *Gazeta Obserwatora PIHM* 7, 1954.
30. CENA, M., CZAJKOWSKI, Z and MASTALERZ, P.: The polarymetric method of Temperature tests in the accomodation of animals - *Roczniki Nauk Rolniczych Tom 67*, 1, 1955.
31. CENA, M., and KRAUSS, S.: The course and distribution of equine infectious anaemia in Poland - *Medycyna Weterynaryjna* 2, 1955.
32. CENA, M. and RZYMKOWSKI, A.: Etudes sur l'éclaircissement solaire des étables - *Zeszyty Naukowe Akademii Górniczo Hutniczej, Kraków Architektura* 1, 1956. Summary in English.
33. CENA, M., JANOWSKI, T., OLPİŃSKA, K. and SLOMKA, J.: Investigations over the dusting in breeding environments - *Zeszyty naukowe wyższiej szkoły rolniczej we wroclawiu*, 4, 1956.
34. CENA, M. and CZAJKOWSKI, Z.: The measurements of evaporation in Pigsties - *Zeszyty naukowe wyższiej szkoły rolniczej we wroclawiu*, 4, 1956.
35. CENA, M. and JANOWSKI, T.: The results of zootechnical stocktaking in two villages - *Zeszyty Naukowe Wyższej Szkoły Rolniczej, Wroclaw*, 4, 1956.
36. CENA, M. : Bioklimatologiczne Fragen der Tierzucht - *Zeszyty problemowe posterpów nauk rolniczych. Zeszyt*, 9, 1957.
37. CRADDOCK, G.W. and FORSLING, C.L.: The influence of climate and grazing on spring-fall sheep range in southern Idaho - *U.S. Dept. Agr.Tech.Bull.*, 600, 1938.

38. CZAJKOWSKI, Z. and BALBIERZ, H.: Introductory investigations over the influence of breeding environment upon the quantity of haemoglobin and erythrocytes in pigs - *Zeszyty naukowe wyzszej szkoly rolniczej we wroclawiu*, 7, 1956.
39. DAVIDSON, J. : On the ecology of the growth of the sheep population in South Australia - *Trans. Roy. Soc. of S. Aust.*, 62, 141-148, 1938.
40. DAVIDSON, J. : On the growth of the sheep population in Tasmania - *Trans. Roy. Soc. of S. Aust.*, 62, 342-346, 1938.
41. FINDLAY, J.D. and YANG, S.H.: The sweat glands of Ayrshire Cattle - *J. Agric. Sci.*, 40, 126, 1950.
42. FINDLAY, J.D. : The effects of temperature, humidity, air movement and solar radiation on the behaviour and physiology of cattle and other farm animals - *Hannah Dairy research Institute Bulletin no. 9*, 1950.
43. FINDLAY, J.D. : Farm animals and high temperatures - *Brit. Agric. Bull.*, 6, 212, 1953.
44. FINDLAY, J.D. and BEAKLEY, W.R.: Environmental physiology of farm animals - Chapter 6 in *Progress in the physiology of farm animals*. Ed. J. Hammond. Butterworths, London, 1954.
45. FINDLAY, J.D. and BEAKLEY, W.R.: The effect of environmental temperature and humidity on the rectal temperature of Ayrshire Calves - *J. Agric. Sci.*, 45, 339, 1955.
46. FINDLAY, J.D. and BEAKLEY, W.R.: The effect of environmental temperature and humidity on the skin temperature of Ayrshire Calves - *J. Agric. Sci.*, 45, 353, 1955.
47. FINDLAY, J.D. and BEAKLEY, W.R.: The effect of environmental temperature and humidity on the scrotal temperature of Ayrshire Calves - *J. Agric. Sci.*, 45, 365, 1955.
48. FINDLAY, J.D. and BEAKLEY, W.R.: The effect of environmental temperature and humidity on the ear temperature of Ayrshire Calves - *J. Agric. Sci.*, 45, 375, 1955.
49. FINDLAY, J.D. and BEAKLEY, W.R.: The effect of environmental temperature and humidity on the respiration rate of Ayrshire Calves - *J. Agric. Sci.*, 45, 452, 1955.
50. FINDLAY, J.D. and BEAKLEY, W.R.: The effect of environmental temperature and humidity on the heart rate of Ayrshire Calves - *J. Agric. Sci.*, 45, 461, 1955.
51. FINDLAY, J.D. : The respiratory activity of Calves subjected to thermal stress - *J. Physiol.*, 136, 300, 1957.
52. FRY, F.E.J. : Effects of the environment on animal activity - *U. of T. Studies Biol. Ser. no. 55*, *Pubn. Ont. Fish. Res. Lab*, 68, 1-62, 1947.
53. FRY, F.E.J. a.o. : Cruising speed of goldfish in relation to water temperature - *Journ. Fish. Res. Bd. Canada*, 7, 4, 169-175, 1948.
54. FRY, F.E.J. : Some environment relations of the speckled trout (*Salvelinus fontinalis*) - *Proc. N.E. Atlantic Fish. Conf.*, 1-29, 1951.
55. HAUBOLD, H. : Fütterungs- und wetterbedingte Schwankungen des Jahreszeitlichen Vitamin-A-Gehaltes in den MilCHFetten - *Milchwissenschaft*, 10, 340-348, 375, 382, 1955.
56. HAUBOLD, H. : Veterinarian Bioclimatology - Areal report: Symposium Int. Soc. Bioclim. and Biomet., Paris 29 - 31 August 1956.
57. JANOWSKI, T. : Investigations on the Microclimate of the Rabbit's dwelling - *Med. Wet.*, 12, 1955.
58. JANOWSKI, T. : Studies on the hygienic conditions of cattle-breeding environments along the Vistula - *Rocz. Nauk. Rol.*, 70, 4, 1956.
59. JANOWSKI, T. : Catathermometric Studies - *Med. Wet.*, 9, 1956.
60. JANOWSKI, T. and RYLKO, A.: Tables of Physiological Saturation Deficit according to Szymkiewicz for 39° - *Manuscript*, Cracow, 1958.
61. JOHNSON, B.C., MITCHELL, H.H. and HAMILTON, T.S.: The riboflavin requirement of growing pigs at two environmental temperatures - *J. Anim. Sci.*, 8, 626, 1949.
62. JOHNSON, B.C., MITCHELL, H.H., HAMILTON, T.S. and HAINES, W.T.: The riboflavin requirement of the growing pig at two environmental temperatures - *J. Nutrition* 41, 317-337, 1950.
63. JOHNSON, H.D. and BRODY, S.: A comparative study of the effects of environmental temperature on various physiological responses of rabbits and cattle - *Thesis*, 1955.
64. KESNER, B. : Methods of investigation of the influence of the basic factors of the atmospheric surrounding on the exchange of gases applied to calves and cattle - *Annals of Czechoslovak Acad. of Agric. Sciences, Ser. A*, 5, 1954.
65. KESNER, B. : The results of microclimatical research of different types of pigsties - *Annals of the Czechoslovak Academy of Agric. Sciences, the series Veterinary*, 6, 1955.
66. KESNER, B. : Carbon dioxide in the air of stables - *Annals of the Czechoslov. Acad. of Agric. Sci.*, the series Animal Production, 1, 1956.
67. KESNER, B. : Studies in the climate of stables. An outline of the methods of measurement of microclimatic elements in stables and an example of microclimatic research - *Ann. of the Czechoslov. Acad. of Agric. Sci.*, the Series Animal Production, 3, 1956.

68. KESNER, B. : Study of barn and stable climate: Temperature - Annals of the Czechoslovak Acad. of Agric. Sci., the Series Animal Production, 5, 1956.
69. KESNER, B. : Study of climatic conditions in barns: Humidity conditions - Annals of the Czechoslov. Acad. of Agric. Sci., the Series Animal production, 7, 1956.
70. KOLLER, R. : Klima und Lebensäusserung unserer Haustiere. I. Allgemeines - Fortpflanzung, Zuchthygiene und Haustierbesamung, 5, 12, 154-159, 1955.
71. KOLLER, R. : Klima und Lebensäusserungen unserer Haustiere. II. Der Einfluss des Lichtes auf die Haustiere, mit besonderer Berücksichtigung ihrer Fortpflanzung - Fortpflanzung, Zuchthygiene und Haustierbesamung, 6, 8 und 10, 1956.
72. KRUEGER, A.P. and SMITH, R.: The effects of air ions on the living mammalian trachea - The Jnl. of Gen. Physiology, 42, 1, 69-82, 1958.
73. KRUEGER, L. : Einfluss der Luftmassen auf den Fettgehalt der Milch - Areal report: Symposium Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956.
74. LEE, D.H.K. : Reactions of Domestic Fowls to hot atmospheres - Proc. Roy. Soc. Qld., 53, 105, 1941.
75. LEE, D.H.K. and ROBINSON, K.W.: Reactions of the rabbit, pig, cat, dog and sheep to hot atmospheres - Proc. Roy. Soc. Qld., 53, 129-200, 1941.
76. LEE, D.H.K. : Experiments in Poultry Housing for Hot Weather - Red. Comb. Bull., 6, 163, 1942.
77. LEE, D.H.K., ROBINSON, K.W., YEATES, N.T.M. and SCOTT, M.I.R.: Poultry husbandry in hot climates, experimental enquiries - Poult. Sci., 24, 195, 1945.
78. LEE, D.H.K. and ROBINSON, K.W.: Animal behaviour and heat regulation in hot atmospheres - Univ. Qld. Papers, Dept. Physiol., 1, 9, 1946.
79. LEE, D.H.K. : Reactions of dairy cattle over a range of controlled temperature and humidity - Feder. Proc., 6, 150, 1947.
80. LEE, D.H.K. and ROBINSON, K.W.: The effect of the nutritional plane upon the reactions of animals to heat - Proc. Aust. N.Z. Assoc. Adv. Sci., 1946 and J. Anim. Sci., 6, 182, 1947.
81. LEE, D.H.K. : Reactions of Jersey Cows in Milk to hot atmospheres - J. Dairy Res., 15, 219, 1948.
82. LEE, D.H.K. : Reactions of Jersey Calves to hot atmospheres - J. Dairy Res., 15, 227, 1948.
83. LEE, D.H.K. a.o. : Assessment of the adaptability of livestock to climatic stress - J. Anim. Sci., 7, 391, 1948.
84. LEE, D.H.K. : Direct relationships between climate and animal welfare - March 1949 British Commonw. Sci. Off. Conference. Specialist Conference in Agriculture, Australia 1949. Paper for presentation to Session A: Review - Climatic Factors.
85. LEE, D.H.K. : Improving Livestock under tropical and sub-tropical conditions - FAO Development Paper 3, Washington D.C., 1950.
86. LEE, D.H.K. and PHILLIPS R.W. (ed.): Report on the Inter-American meeting on Livestock Production, Rpt. FAO, 1950.
87. LEE, D.H.K. a.o. : The effect of the dietary Plane upon the reactions of two breeds of sheep during short exposures to hot environments - Aust. J. Agric. Res., 1, 2, 217-230, 1950.
88. LEE, D.H.K. : Studies of heat regulation in the sheep with special reference to the merino - Aust. J. Agr. Res. 1, 2, 200-216, 1950.
89. LEE, D.H.K. and RIEK, R.F.: The time factor in response of Dairy cattle to hot conditions - Mimeographed for the 43rd Meeting of the Am. Soc. of Animal Production, 23 November 1951.
90. LEE, D.H.K. : Report to the Government of India on studies in animal climatology - FAO of UN, ETAP FAO, Rpt. 53, 1952.
91. LEE, D.H.K., FOHRMAN, M.M., MCDOWELL, R.E.: Progress report of the Red Sindhi Crossbreeding Experiment at Beltsville Md. - Dept. of Agr. Bureau of Dairy Industry Inf., 128, 1953.
92. LEE, D.H.K. : Manual of field studies on the heat tolerance of domestic animals - FAO Development Paper no. 38 (Agriculture), 1953.
93. LEE, D.H.K. : Respiratory activity as an index of heat tolerance in Jersey and Sindhi X Jersey (F_1) crossbred cows - J. Anim. Sci., 12, 573, 1953.
94. LEE, D.H.K. and PHILLIPS, R.W. (ed.): Report of the second Inter-American meeting on Livestock Production - FAO Development Paper no. 33 (Agriculture), 1953.
95. LEE, D.H.K., MACDOWELL, R.E. and FOHRMAN, M.H.: The relationships of Surface Area to heat tolerance in Jerseys and Sindhi Jersey (F_1) Crossbred Cows - J. Anim. Sci., 12, 4, 747-756, 1953.
96. LEE, D.H.K., MACDOWELL, R.E., MCMULLAN, H.W., FOHRMAN, M.H. and SWETT, W.W.: Body weights, body measurements, and surface area of Jersey and Sindhi-Jersey (F_1) crossbred females - J. Dairy Sci., 37, 1420-1428, 1954.

97. LEROY, P. : Croissance disharmonique de la crête du Coq domestique maintenu à l'obscurité - Séances de l'Acad. des Sciences, 243, 210-212, 1956.
98. LEROY, P. : Croissance de la crête du coq domestique maintenu à l'obscurité - Les Ann. d'Endocr., 19, 1, 55-68, 1958.
99. MACDONALD, M.A. and BELL, J.M.: Effects of low fluctuating temperatures on farm animals. I. Influence of ambient air temperature on the respiration rate, heart rate, and rectal temperature of lactating Holstein-Friesian cows - Can. Jnl. of Animal Sci., 38, 10-22, 1958.
100. MACDONALD, M.A. and BELL, J.M.: Effects of low fluctuating temperatures on farm animals. II. Influence of ambient air temperature on water intake of lactating Holstein-Friesian cows - Can. J. of Animal Sci., 38, 23-32, 1958.
101. MACLEOD, J. : The nature, epidemiology and control of sheep Myiasis in Britain - J. Comp. Path. Therap., 50, 10-32, 1937.
102. MACLEOD, J. : The experimental production of cutaneous Myiasis of sheep - Parasitology, 29, 526-529, 1937.
103. MACLEOD, J. : The ticks of domestic animals in Britain - Emp. J. Exp. Agric. 7, 97-110, 1939.
104. McDONALD, I.W. : Facilities for studies in climate physiology of sheep and cattle at the Sheep Biology Lab., Prospect, N.S.W., Australia - Australia-UNESCO Symp. on Arid Zone Climatology, Canberra. Papers from Australia and New Zealand, 30, 6, 1956.
105. McFARLANE, W.V., BROOKS, C.M. and PINKSTON, J.A.: Effects of variations in temperature on excitability, conduction and refractory period of the mammalian heart in situ - Am. J. Physiol., 181, 1952.
106. McFARLANE, W.V., MORRIS, R.J. and HOWARD, B.: Water economy of tropical Merino sheep - Nature, 178, 304, 1956.
107. PRIMAULT, B. : Du climat des étables ouvertes - Le Sillon romand. 57ème année, Lausanne, no. 39, 1955.
108. PRIMAULT, B. : De l'influence des variations de la pression atmosphérique sur l'apparition de la fièvre aphteuse - Schw. Archiv.f. Tierheilkunde, 97, 9, 1955.
109. PRIMAULT, B. : Des critères d'acclimatation du bétail en zone subtropicale - Miscelanea Geofisica à Luanda, Angola, 13-24, 1956.
110. PRIMAULT, B. : Influence of weather on the incidence and development of infectious diseases of livestock - Commission for Agricultural Met., 2nd session, Warsaw, 1958. World Met. Org., Rep. No. CAgM-II/Doc. 30 (2-11-1958).
111. ROBINSON, K.W. and KLEMM, G.H.: A study of heat tolerance of grade Australian Illawarra Shorthorn cows during early lactation - Austr. J. Agric. Res., 4, 224-234, 1953.
112. ROBINSON, K.W. : The role of water for heat dissipation by a Jersey cow and a corriedale ewe - Austr. J. Agr. Res. 5, 568-577, 1954.
113. ROBINSON, K.W. and KNAPP, B.J.: The heat tolerance of two breeds of calves from 1 to 12 months of age - Austr. J. Agric. Res., 6, 350, 364, 1955.
114. SCHEIN, M.W. : Dairy-cattle breeding studies at the Iberia livestock experiment station - US Dept. of Agric., Agr. Res. Adm., Bureau of Dairy Ind., Washington 25, D.C., Sept. 1952.
115. SCHEIN, M.W. a.o. : Dairy cattle research at the Iberia livestock experiment station - United States Dept. of Agr., April 1955.
116. SCHEIN, M.W. : Report of research and activities conducted at the Iberia Livestock Dairy Experimental Station Jeanerette, Louisiana from Jan. 1, 1954 - Dec. 31, 1954. - US Dept. of Agr., 1955.
117. SCHEIN, M.W., MACDOWELL, R.E., LEE, D.H.K. and HYDE, C.E.: Heat tolerances of jersey and sindhi-jersey crossbreds in Louisiana and Maryland - J. of Dairy Science, 40, 11, 1405-1415, 1957.
118. TRUMBLE, H.C. : The relation of wet-season variability to fat-lamb production in S. Australia - Trans. Roy. Soc. of S. Aust., 69, 16-21, 1945.

ZOOLOGICAL BIOCLIMATOLOGY

Section D: World literature

ENTOMOLOGICAL BIOCLIMATOLOGY*

1. ASHTON, P.S. : Light intensity measurements in rain forest near Santarem, Brazil - J. Ecology, 46, 65-70, 1958.
2. BAR-ZEEV, M. : The effect of temperature on the growth rate and survival of the immature stages of *Aedes Aegypti* L. - Bull. ent. Res., 49, 157 - 163, 1958.
3. BELOZEROV, V.N. : The influence of air humidity on the mite *Ornithonyssus bacoti* Hirst 1913 (Parasitiformes, Liponyssidae) - Entom. Obozrenie, Moskva, 37, 47-63, 1958. (In Russian).
4. BIEL, E.R. : New Jersey climate and its influence on life - Proc. 43rd Ann. Meeting N.J., Mosq. Ext. Ass., 79-85, 1956.
5. BUCKENS, F. : Climatologie, technique et habitation tropicale - Fol. scient. Afr. contr., 2, 11-13, 1956.
6. BURCHFIELD, H.P. and GOENAGA, A. : Equipment for producing simulated rain for measuring the tenacity of spray deposits to foliage - Contr. Boyce Thompson Inst., 19, 133-140, 1957.
7. BUNNING, E. : Ueber den Temperatureinfluss auf die endogene Tagesrhythmik, besonders bei *Periplaneta americana* - Biol. Zbl., 77, 141-152, 1958.
8. CREGG, J.B. : The action of climate on the larva, prepupae and pupae of certain blowflies - pp. 387-389 IN - XIV Int. Congr. Zool., Proc., Copenhagen, Aug. 1953, 1956.
9. DIGBY, P.S. : Flight activity in the blowfly *Calliphora erythrocephala* in relation to light and radiant heat, with special reference to adaptation - J. exp. Biol., 35, 1-19, 1958.
0. DORE, W.G. : A simple chemical light meter - Ecology, 39, 151-152, 1958
1. DUPORTE, E.M. : Some modern trends in entomology - Ann. Soc. ent. Québec, 1, 51-52, 1956.
2. EL-ZIADY, S. : A probable effect of the moonlight on the vertical distribution of Diptera - Bull. Soc. ent. Egypte, 41, 655-662, 1957.
3. GRIFFITHS, J.S. : Climatic zones of East Africa - E. Afr. agric. J., 23, 179-185, 1958.
4. HARKER, J.E. : Diurnal rhythms in the animal kingdom - Biol. Rev., 33, 1-52, 1958.
5. HENSON, W.R. : A temperature gradient apparatus for small insects - Canad. Ent., 90, 54-59, 1958.
6. JOHNSON, C.G. : The distribution of insects in the air and the empirical relation of density to height - J. anim. Ecol., 26, 477-494, 1957.
7. NICOLAS, J.P. : Les climats africains en biogéographie humaine - Bull. Inst. français Afr. noire (B), 20, 7-67, 1958.
8. PAVLOV, S. : Die Bewegungsorientierung der photopositiven Insekten auf Farben - Acta Fac. Rer. nat. Universit. Comenianae, Bratislava (Zool.), 2, 155-162, 1957.
9. PLATT, R.B., LOVE, G.J. and WILLIAMS, E.L. : A positive correlation between relative humidity and the distribution and abundance of *Aedes* - Ecology, 39, 167-169, 1958.

20. SALT, R.W. : Natural occurrence of glycerol in insects and its relation to their ability to survive freezing - Canadian Ent., 89, 491-494, 1957.
21. SMITH, W.W. and LOVE, G.J.: Effects of drought on the composition of rural mosquito populations as reflected by light trap catches - Mosquito News, 16, 279 - 281, 1956.
22. SNOW, W.E. : Stratification of arthropods in a wet stump cavity - Ecology, 39, 83 - 88, 1958.
23. TAKASU, K. : Microclimatic study IV Microclimatic characteristics of some specified surface conditions - Ber. Ohara Inst. (Biol.), 11, 22-74, 1957.
24. WANG, H.H., and CHANG, C.: The outdoor nocturnal biting activity of ANOPHELES HYRCANUS var. SINENSIS in relation to the factors of temperature, humidity, light and rain - Acta ent. Sinica, 7, 473-480, 1957 (In Chinese).
25. WELLINGTON, W.G. : The synoptic approach to studies of insects and climate - Ann. Rev. Ent., 2, 143-162, 1957.
26. WILLIAMS, C.B., and EL-ZIADY, S.: On the relative distribution of insects at 5 and 30 feet - Bull. Soc. ent. Egypte, 41, 663-675, 1957.
27. WOLSKA, H. : Preliminary investigations on the thermic preferendum of some insects and spiders encountered in snow - Polia biol., Krakow, 5, 195-208, 1957. (In Polish).

PART IV

HUMAN BIOCLIMATOLOGY

(1959)

Section A : Physiological bioclimatology

1. General physiological bioclimatology
2. Geographical bioclimatology
3. Ethnological bioclimatology
4. Acclimatisation bioclimatology

HUMAN BIOCLIMATOLOGY

Section A4 : Physiological bioclimatology (Acclimatisation-bioclimatology)

A CONTRIBUTION TO THE PHYSIOLOGY OF DAMP COLD
AN ASPECT OF TEMPERATE CLIMATE

by

Dr. E.T. Renbourn (Great Britain)*

PART I THE CHILLS OF A DAMP COLD CLIMATE
AN HISTORICAL SURVEY

by

Dr. E.T. Renbourn (Great Britain)

1. EARLY PERIODS TO THE END
OF THE 17th CENTURY

In earliest times it was widely accepted that the soul, the life principle and the intellect were, in some way, linked with air and breathing. It is hence not surprising that the early Greek philosophers took it for granted that atmospheric air, with its vapours and contents, flowed freely into and out of the body not only through the lungs but through invisible pores of the skin and internal organs. Such rhythmic breathing, known as insensible perspiration (Greek, "adelos diapnoe" - to breathe through invisibly) was believed to be brought about by the alternate contraction and relaxation of the heart and blood vessels. Great importance was attached to this pulsatile process, because not only did it allow the whole body to be thoroughly "aired" but the insensible excretory vapours (and "fuliginous" or sooty fumes produced in the burning of the "vital flame" of the heart) could be passed out of the body. (Ref. no. 1).

The Greeks believed that the human frame was composed of four primitive elements - Air, Earth, Fire and Water - which had the dual properties of hot and moist, cold and dry, hot and dry, and cold and moist respectively. These elements were, somehow, combined to produce the four bodily humours - Blood (hot and moist), Black Bile (cold and dry), Yellow Bile (hot and dry) and Phlegm (cold and moist). It is clear that such physical properties, elements and humours were not such as to be easily measured or even ascertained by the senses, and contradictory statements abound in the early literature.

The nature, constitution or temperament of people was, so it was said, due partly to the mixture of the four primary properties or humours they were born with, but also to the climate or weather in which they lived. In damp countries, said Hippocrates, the inhabitants have a humid, gross, flabby and squat habit of body, and are liable to rheumatism of the joints, sciatica and gout. It was this excess humidity of the body (partly derived from the moisture of the surrounding air) which, he said, produced the poorly visible joints and veins, gave roughness to the voice and made these people less prolific than their neighbours. In any case, he added, this was to be expected, for did not the men wear breeches and spend most of the time on horseback? Children, it seems, were of a hot and moist constitution, but this became colder and drier as age advanced. (Ref. no. 2).

Aristotle showed in the "Meteorologica" that he was conversant with the simpler phenomena of climate and weather and the nature of clouds, rain and dew. His pupil, Theophrastus, laid down the rules of weather lore in the "Book of Signs". Theophrastus believed, as did Hippocrates, that moisture clogged the intellect of man. "Other living creatures are inferior in understanding", he said, "for they breathe air that comes from the earth and take moister nourishment. The same principle explains why children lack understanding; for they are extremely moist and, in consequence, air cannot make its way throughout the body but it is set apart in the breast leaving them sluggish and witless" (Ref. no. 3). On the matter of the most healthful air, Galen

*) Director Clothing and Stores Experimental Establishment. Directorate of Physiological and Biological Research, Ministry of Supply, Farnborough, Hants., Great Britain.

noted that "pure air is that which is not defiled by exhalations from pools and marshes or from a pit giving off harmful vapours. And harmful too is that which is contaminated from the putrefaction of animals and vegetables. But the diversity of air which arises from heat, cold, dryness and moisture is not the same to all people "but.. the best is that most opposite to the predominant quality" (Ref. no. 4). Thus, for the sanguine (hot and moist) individual a cold and dry climate was, apparently, the best; but cold and damp was clearly more fitted for those of a bilious or hot and dry constitution - who were easily moved to anger and to violence. These various ideas on weather, climate and on the temperament of man, were to continue, often without modification, through medieval periods, the renaissance and later centuries, to almost modern times.

Many Greeks and later Roman physicians of the so called Methodic Schools ascribed fevers, inflammations and other diseases, to blocking of the pores of the skin - itself arising, so said Galen, from the insensible perspiration being too much increased or too thick, to the pores being unduly contracted by unwanted cold, or clogged by an air too damp or too heavy. Thus it followed that an atmosphere both cold and damp was particularly dangerous to health. For, when the excretory vapours could not get out of the skin or lungs, the stagnating humours of the body became greatly increased in amount (plethora) and could be directed to other places, such as the brain, the bowels, the lungs or kidneys, and there give rise to disease.

John Cais, describing the "Ephemera Brittanica" or "English Sweat" (1552) (Ref. no. 5), said that this arose from "takeinge infection from evill brethynges places, landynge muddy and rotte groundes .. burieng dede bodyes .. dong Hilles". He advised, "drying the moiste aier with fyres ... in houses .. or in that side of cities and houses that lieth towards the enfection".

It must have always been common observation that water cooled more than air at the same heat and that clouds hid the warmth of the sun. "That air is thick", said Robert Burton, "which is kept longer before it be warmed by the sun beams, and once heated like an oven will keep itself from cold". (Reference no. 6).

The old Greek and Roman ideas on the nature and purpose of the insensible perspiration of lungs and skin were resuscitated by Sanctorius at the beginning of the 17th century. (Ref. 1 and 7). By weighing the "matter of perspiration" (insensible perspiration - but for centuries called, plainly, "perspiration") under various conditions, he showed, apparently, that it was far more important than all the other "servile excretions" put together. Violent passions, a cold damp or foggy air, and the cold dews of the dawn were, he said, the common causes of closing up the skin pores, and, hence, inevitably led to fevers, rheumatism, inflammations and to most other ills of the flesh. He stressed that a damp air, with its contained contagious particles, was also able to get into the body through the pores of the skin, particularly when these were first dilated by hot weather or by violent exertion. "The heat of the sun", said Francis Bacon, "is much milder and gentler in degree than that of a fire. It is much more moist in quality, especially as it is transmitted through the air". (Ref. no. 8). The warmth of clothing, he suspected, was due to the air it contained, for, he said, "there is doubt as to whether the warmth of wool, skin, feathers and the like, is derived from slight inherent heat since they are animal excretions .. or merely from the confinement and separation of air. For all air appears to possess a certain degree of warmth when separated from the external atmosphere .. Heat, with regard to the human senses", he continued, "is various and relative, so that lukewarm appears hot if the hand be ice cold, and cold if the hand be hot". (ref. no. 8). This matter was, however, well known to the early Greeks, many of whom insisted that, in any case, the senses could be fallacious. Robert Boyle (1665) (ref. no. 9) appreciated convection as a source of heat loss; and, a wind, he said, "by its motion is able not only to drive away the air contiguous to the hands and face and the warm steams of the body which tempered its coldness, but to pierce deeper than the calm air is wont into the pores of the body". "Cold naturally stops the pores", added Nicholas Culpeper, "congeals the humours, stops sweate, staies motion, and by this so stops the power of the spirits that it consumes the strength of the head and bowels". (ref. no. 10).

2. THE CHILL OF DAMP COLD. 18th CENTURY

In his book on the so called "Non Naturals" (1707) (ref. no. 11), Jeremiah Wainwright gave some thirty pages to the subject of the physics of the air and its effects on the body. "The Fens of Cambridgeshire", he wrote, "fill the Air with Vapours, whereby its Elasticity is weaken'd, the Fibres becoming relax'd and the Pores of the Skin obstructed. Now, if Coldness be added to the Moisture of the Air it will with so much more certainly produce these Distempers.... for Cold contracts and binds up, makes the Blood more compact, so that its Motion becomes slower and the Viscidity greater". Not only was damp cold a question of weather and climate but also of the night time; for, even in warm weather, cold and so called treacherous dews were often present before dawn. Wainwright ascribed consumption to "a fenny, heavy and smoaky air", but

it was to take almost two centuries to realise that the cold "Vapours" of the fens, and the "Distempers" they produced, arose from the malarial ("mal-aria") mosquito and the parasite transmitted in its bite.

John Arbuthnot (1733) elaborated even more on the effects of the atmosphere. "Moist air", he stressed, "relaxeth the human Frame by weakening the Spring of the Air, and Moisture helps Air to insinuate itself into the Pores of human Bodies and to increase its Dimensions; and perhaps a due quantity of it is necessary for a right State of the Fluids and Solids ... It is very possible that human Bodies absorb Water from moist Air as fixt, dry, alkaline Salts do .. and many sudden changes may happen by absorbing the outward Air and all its Qualities and Contents. Nothing accounts so clearly for epidemical Diseases seizing human Creatures.." (ref. no. 12).

Senac (1749), the French physician, wrote more clearly on the effects of a damp cold climate. "When air is both humid and cold", he wrote, "it produces a more intense impression than when dry, for water is of a greater density and hence robs the body of more heat. It may, besides, be impregnated with various corpuscles which can, for example, seize on the lungs". (ref. no. 13). Another Frenchman, Sauvages, in his "Dissertation on Air" (1753), added: "Most vapours which are harmful to man are burning or stinking and extinguish the activity of the electric fluid, for... moisture destroys electricity". (Ref. no 14). Such ideas on damp air and electricity were to be met with periodically in the literature. Reminiscing on the campaigns of 1742 - 1750, John Pringle, Surgeon-General to the British Army, noted that "the evenings growing cool, night fogs, occasioning a stoppage of perspiration (viz. insensible perspiration), brought on the dysentery" (Ref. no. 15). Naval surgeons of the time, although aware of the real cause of scurvy, nevertheless insisted that an important contributory factor was the obstructed "perspiration" arising from the effects of "noxious" land breezes and the "damp cold air from the sea". John Chandler (1761) (Ref. no. 16) enjoined, most stringently (as did Pringle, Count de Sax and others), the use of waistcoats or flannel belts (Ref. no. 17) for maintaining the flow of insensible perspiration in the face of the dews and chills of the night. He realised that there were other factors than air concerned in health and comfort, for a changing barometer could also stop the flow of cutaneous vapours. Of itself, a falling barometer had, he said. "a considerable effect on the organ of feeling, and changes the state of our bodies from easy to disagreeable sensations.... in a manner similar to that when we retire from the sun's rays into the shade". (Ref. no. 16). "It is incumbent upon the learned", wrote Tissot in 1768, "to choose as much as possible a temperate place and dry air. These are beneficial to the lungs, encourage the circulation and strengthen the fibres. Cold and dry air is tolerable, but a cold and damp air is very dangerous; it increased the disorders incident to Men of Letters, relaxes, obstructs the insensible perspiration, and produces catarrhs, rheumatisms and palsies". (Ref. no. 18). The moisture of the air was assumed to produce its effects by either mechanically obstructing the exit of cutaneous vapours through the skin pores or by stimulating the so called "absorbent" vessels (or lymphatics) of the skin. Some believed that moisture somehow "debilitated" the cutaneous blood vessels, thus impairing the natural force which expelled the insensible perspiration through their walls.

Even at this period there was already controversy as to whether wool was the best material to protect the body from cold and damp. Colombier, French army surgeon, was in doubt for, said he, wool "easily retains miasms floating in the air, dries with difficulty and finally imbibes sweat which then corrupts and gives out putrid and dangerous odours". (Ref. no. 19). Gilbert Blane (1785) argued, further, that wool was to be regarded as a sort of filter "to separate off the impurities of the air before it comes in contact with the surface of the body, and, therefore, any unreasonable change of apparel may be very unsafe to health". (Ref. no. 20).

Edward Rigby (1785) put forward the idea, expressed by Senac, that dry air is a good thermal insulator. Elaborating on the matter, he noted: "moist air is likewise a better conductor of heat than when dry, because water, though of the same temperature of the air, is well known to carry it off more quickly than air will do. If, therefore, these two causes (viz. cold and damp) unite, as in the case of moist and windy weather, we may easily understand why the heat of animals should be carried off more quickly and the animal experience a greater sense of cold than when the air is still and dry though the thermometer should, in both cases, stand at the same point". (Ref. no. 21). About this period, Count Rumford apparently proved the matter by his experiments on heated metal cylinders (covered with various cloths) allowed to cool under a variety of conditions. (Ref. no. 22). He also demonstrated the truth of the matter vexing Francis Bacon - that the warmth of natural materials is due, in great part, to the air they contain.

Writing at the end of the century in his popular "Family Medicine", Willich pointed out that "cold renders bodies more compact, particularly the solid parts of the animal structure such as muscles, nerves and bones; they become more elastic in winter. Damp or moist air suddenly relaxes and enervates, occasions a slowing of the fluids which gives rise to obstructions and impedes both the circulation and secretion of the humours by choking the insensible perspiration". If, said he, "the moisture of the air increases, we experience an uncomfortable torpor or ennui; with the loss of energy we lose our gaiety and the mind is depressed as well as the body. Damp

places and districts are always unwholesome but more particularly so in cold weather..... (Ref. no. 23). Like Senac and others, Willich believed the effects of weather were in some way related to electricity. This, he said, "supplies the fibres with their natural tone, stimulates the vessels to a more vigorous action and increases the serenity of mind. This principle does not exist in the atmosphere of all places in equal quantity .. nor is it so at different seasons or at all times of the day. All men do not inhale this electric matter in equal quantities, and the remarkable differences in temperament are thus produced". (Ref. no. 23). Such an argument can be found even in modern books of medical climatology; nevertheless, much of it is reminiscent of Galen and Hippocrates. In the same year, Hufeland, discussing means of prolonging life, gave his opinion on weather and climate. "Air which contains a fine moisture", he noted, "is best for attaining great age .. for moist air, by being already saturated, has less attraction over bodies, that is to say, consumes them less. Besides, in a moist air, there is almost a greater uniformity of temperature Lastly it keeps the organs longer pliable and youthful. A striking proof of this is afforded by Islands ... which have at all times been, and still are, the cradles of old age". (Ref. no. 24). Here we have an argument in favour of a damp environment, but its logic is, again, reminiscent of earlier periods. Most of the remarks made during the 18th century on damp cold were, in great part, of a subjective nature. Physicians were, however, aware of the known meteorological instruments and occasionally made use of them with fair success. Louis Cotte, in his "Treatise on Meteorology" (Ref. no. 25), gave a chapter on Medical Climatology and discussed the importance of air pressure, humidity, temperature and wind on health and disease; but there was little outstanding to note. John Quincy, writing on hygrometers in his "Medical Dictionary" of 1794, said: "E x far the earliest notices of changes of this kind may be made use of by a physician, the skilful alone can judge". (Ref. no. 26). It is doubtful whether such measurements were often made during damp cold weather.

3. PHYSIOLOGY OF THE DAMP COLD PHENOMENON EARLY 19th CENTURY

Bichat (1802), the French anatomist, was amazed that animals could, apparently, take up so much atmospheric humidity into the body and, he added, "during a fog of twenty four hours duration, thrushes or red breasts are reported to become so fat that they are unable to fly from the sportsman". (Ref. no. 27). William Buchan, in his long popular "Domestic Medicine" (1807), continued the old ideas. "Very cold air obstructs the perspiration, constricts the solids and condenses the fluids. It occasions rheumatisms, coughs and catarrh, with diseases of the throat and breast. Air that is moist destroys the elasticity or springs of solids, induces a phlegmatic and lax disposition and disposes the body to agues or intermittent fevers". (Ref. no. 28). Such diseases, said he, were frequent in Holland, the Fens of Cambridgeshire and the Hundreds of Essex, but their real cause has already been suggested.

When, in 1824, William Edwards, French physiologist (Ref. no. 29), showed experimentally that, in man and animals, the amount of insensible perspiration was much greater in a dry than damp atmosphere of the same heat, he appeared to have confirmed the old belief that damp air clogged the "exhalant vessels" and skin pores. He also demonstrated, apparently, that a cold damp air did not, however, chill the body more than a dry one at the same temperature. This he explained by the fact that although damp air was a better thermal conductor, it simultaneously inhibited insensible perspiration and, thus, decreased both evaporation and heat loss from the body. This confirmation of the long accepted belief in the chilling effect of damp cold was, nevertheless, to pass almost unnoticed.

Describing his experiences in the disastrous Russian Campaign of 1812, Moricheau Beaupré (1826) (Ref. no. 30), army surgeon, wrote on the effect of cold on the French soldiers. He divided cold into "real" and "sensible", the former representing that of the environment, and the latter the sensations (which could be modified by the "nervous power") felt by the body. "Real" cold was, he said, of a dry or damp nature. In dry cold the atmospheric temperature fell to freezing point or below, the air was contracted and its water vapour greatly diminished. Hence, he argued, its oxygen must be greater; and since, furthermore, the insensible perspiration of the lungs was increased (that of the skin being decreased), respiration was enhanced and the whole body stimulated to greater activity. The real danger of dry cold to the campaigning soldier lay in the absence of good food, the insufficient rest and the lack of available warm shelter. "Dry cold", he continued, "is a sign of a durable or little variable atmospheric state; no evaporation from the ground occurs, the heavens are serene, the air light, cheery, pure and clear; this condition is salutary to man. Humid cold is characterised by a less depression of temperature but the air holds much aqueous vapour. The humid state is noxious and is the most dangerous enemy of the animal economy; it produces effects altogether opposed to the former. Of this we are convinced by comparison of an inhabitant of a humid country with one of a dry cold; the one

is dull, bloated and phlegmatic, the other robust, nimble and active. Those of a sanguine or bilious nature are less chilled than those of a nervous or lymphatic temperament". Beaupré was well aware that the difference between dry and damp cold was one of weather and climate and not simply of air humidity alone; but he failed to stress the importance of barometric pressure, the sun and wind. There was, in fact, little objective basis for any of his arguments, many of which can be traced to Sanctorius, Galen and Hippocrates.

The existing ideas on the effect of a humid air were bedevilled by John Stark who, in 1833 (Ref. no. 31), believed he had proved that dark colours (and clothing) were more capable of absorbing odours, and hence (so others reasoned) humid air and its miasms. Andrew Combe, (1837), (Ref. no. 32) renowned hygienist of his time, went back a century or more in some of his arguments. "A moist air diminishes evaporation from the skin", he said, "partially shutting up the natural outlet of the superfluous heat of the body; and as it, at the same time, checks the exit of waste matters which ought to be thrown out, and which are known to be injurious to the system as an active poison, the hurtful consequences of such weather and climate, and the fevers, dysenteries and colds to which it gives rise, are partially accounted by it... A moist state of the atmosphere is also favourable to absorption, and hence if noxious effluvia are at the time floating in air, they are more easily received into the system. It is on this account that night air is so unwholesome, particularly in malaria districts which are loaded with moisture and miasmata or marsh poison; for where the air is dry as well as hot, free evaporation takes place and absorption is null".

Having reviewed the earlier literature, William Carpenter (1855) (Ref. no. 33) came to the conclusion that considerable amounts of atmospheric vapour must be absorbed into the human body. He went even further than Bichat and instanced the case of two jockeys who, within an hour or so of a race, became overweight by several pounds under conditions where such vapour absorption could be the only possible explanation. Such was the credulity of the physiologist a century ago! Much was known, he continued, of the adverse effect of damp air at high temperatures, but "accurate experiments are yet wanting to determine the influence of humidity of cold air..... The depressing influence of cold with moisture, the existence of which is a matter of ordinary experience, is probably exerted directly upon the nervous system". Carpenter was well aware of the work of Edwards but doubted the accuracy of the latter's experiments on the effects of damp cold.

Ronald Martin (1856) (Ref. no. 34) had his own ideas as to the casual factors of agues and intermittent fever. "Electricity", he noted, "may be concerned with the generation of malaria, whatever its nature. A low state of electric tension causes obstruction of the nervous power and produces enervation of the capillary system and an inverted condition of the bowels. Humid air, at an equal or superior temperature, produces a peculiar sensation of cold which differs not only in its intensity but in its nature. It is more profoundly felt, and seems to penetrate the whole system, and predisposes particularly to paleness and shivering. In dry air, on the contrary, a sensation is experienced which is called sharp cold... and an increase in activity is experienced... Damp cold must tend to produce, in individuals whose power of developing heat is rather feeble, the series of actions which constitute the accession of an intermittent fever, particularly if exposed to that action during sleep. The chief effect of moist air is exerted on evaporation from lungs and skin".

In 1858, Coulier (Ref. no. 35), professor of Military Chemistry at Vale de Grace in France, repeated Rumford's work on heat loss from fabric coated heated cylinders. The results led him to conclude that there were two levels of water equilibria in garment fabrics when worn; that arising from water vapour (from skin within and humid air from without) condensed in the fibre - "l'eau hygrometrique" - with liberation of latent heat, and that due to deposition of liquid water in the fabric interstices (from rain or sweat) with displacement of air - "l'eau d'interposition". This appears to be the beginning of interest in what is now called "sorption heat" of textile fibres, a property which may, itself, shed some light on the physiology of the damp cold phenomenon. Moore (1862) (Ref. no. 36), like Blane before him, argued that clothes could act as a filter of miasmas, for, he said, "there are facts tending to prove that the body well wrapped in flannel, or clad in warm clothing, so as to prevent a check to perspiration and chill from the night air, is more capable of resisting malaria and inflammation of the liver and kidney". Edmund Parkes (1866) (Ref. no. 37), Professor of Hygiene at the Army Medical School in England, referred to the earlier work of Lehmann, on birds and rabbits, which suggested that more carbonic acid was exhaled from the lungs in a moist than in a dry atmosphere; but he was not prepared to say whether this was physiologically good or harmful. As for malaria, since this, he said, entered the body through the lungs or stomach, he could not agree with Moore as to the particular value of flannel in its prevention.

4. PHYSIOLOGY OF THE DAMP COLD PHENOMENON LATE 19th CENTURY

John Parkin, writing in 1882 (Ref. no. 38) on the relationship of climate and weather to phthisis, was naturally concerned with the effects of cold damp conditions. Although his arguments were not always consistent, he also concluded that cold damp was not necessarily more chilling than cold dry weather and, therefore, not particularly harmful. "Its is well known", he said, "that a damp or moist air generally feels colder than a dry one - temperature and all other circumstances being the same. The reason is that a damp air is a better conductor of heat than a dry one, particularly when accompanied by a strong wind. But no such effect is produced when there is calm or with a fog which has a tendency to check radiation from the surface. Hence it is that a moist atmosphere is frequently close and muggy, or, in other words, warmer than a dry one. On the other hand, a cold dry air may allow a much greater radiation of heat from the surface than a cold humid one, especially when accompanied by wind. Fogs and moisture may, therefore, prove beneficial, rather than otherwise, under certain circumstances and in cold latitudes. These impressions are confirmed by the fact that a dry and warm climate, so far from proving beneficial or acting as a preventative of phthisis, would seem to produce the opposite effect. This is the case of the South of France, or of Provence and Languedoc". (Ref. no. 38). Giving mountain resorts as an example, Parkin stressed the remarkable heating effect of a winter's sun under freezing conditions. He criticised other students of medical climatology (amongst whom he should, perhaps, be included) who "ascribed effects to causes recognisable to their senses, rather than to those deduced from strict analysis of all the phenomena". Parkin's suggestions concerning the effects of moist air on the radiation from the human body (based no doubt on the work of Fyndall), on the warmth of sunlight and the effect of wind, are aspects of the cold damp phenomenon to which sufficient thought has not yet been given. In spite of this illumination thrown on the subject, Thomson (Ref. no. 39), in the same year, went back half a century or more in his arguments, for, he said, "damp and moisture is one of the most prolific and most generally acknowledged source of disease in whatever way it is applied to the body; whether in the atmosphere, or the bed, it is alike apt to be productive of bad consequence and often of the most serious character". When combined with decaying vegetable matter, and more especially when increased by heat, fever and agues are the result; when cold and damp unite their depressing influences, colds of every kind, influenza, scrofula and consumption, rheumatism and neuralgia ensue... The chief evils to be dreaded are the cold winds which come off the water laden with vapour and which are liable to produce croup in children and catarrhal affection in the predisposed". (Ref. no. 39). Gustav Jäger (Ref. no. 40) (the originator of "sanitary" Jaeger wool) insisted that wool must be worn from head to toe, for not only did it absorb harmful damp effluvia of the air, but also a volatile "noxious" agent liberated in the insensible excretions of the body (especially after severe and unpleasant emotions!). Following Stark he insisted that dark coloured wool was far the best for removing such harmful matters (but light colours could be acceptable in the summer).

On the subject of damp cold, scientific enlightenment and return to the humoral theories of Galen and Hippocrates alternated to the very end of the 19th century. Even as late as 1897, Quain's "Dictionary of Medicine" (Ref. no. 41) written and revised for the expert, refused to accept the existence of "le microbe du paludisme" of Laveran as the cause of malaria. As for the vector mosquito, this was not even mentioned. On the other hand it was still seriously entertained that this disease was "the product of moisture ... and of vegetable decomposition". Before the beginning of the new century, Rubner and von Lewschen (Ref. no. 42), using a careful technique, confirmed the finding of Edwards that a high air humidity lessened evaporative cooling of the body by decreasing the amount of insensible perspiration.

5. CONCLUSION

The fear of cold and damp probably arose with the ancient Greek concept of insensible perspiration and the belief that atmospheric air, with its water vapour and effluvia, could pass into the body through the pores of the skin. For centuries, so called logical arguments, subjective and personal impressions and co-incidental factors, were interpreted as representing objective truths. Because he used a balance in his experiments, Sanctorius gave a stamp of authority to the dogma that obstruction to the "matter of perspiration", by cold or damp, was the cause of the discomforts and most of the diseases of the flesh. Such dogma was to remain till almost the end of the last century. There is no doubt, however, that malnutrition, bad hygiene, infection and other non climatic factors have played their role in the colds, the pneumonias, consumption, rheumatism, scurvy, dysentery and other diseases, which for centuries were ascribed to the humid cold of the atmosphere.

In the past, damp cold and its effects was mainly the concern of physicians who wrote lengthily on the matter but rarely measured or even adequately defined it. Not all however, were afraid of the dire consequences. Hufeland was convinced that one lived longer in a damp air, Edwards failed to prove its peculiar chilling qualities and Parkin advised consumptives to remain in the cold damp of England.

The expression damp cold is used nowadays more often by the layman, but it is still assumed by the more discerning that damp air is a better heat conductor and convector than dry air. However, modern workers have shown that there is, in fact, little difference in thermal insulation of moist and dry air and that convective heat loss in such atmospheres is almost the same. Little scientific work has yet been done on the matter of the cold damp phenomenon and this receives but scant attention in the texts of physiological or medical climatology.

More than two centuries ago it was suspected that the effects of a temperate winter involves a problem much wider than that of the physical properties of humid air (indoors or without) or even its effects on clothing and the skin. The solution to the problem may, perhaps, lie in the change of heat or of the water balance of the body resulting from the complex interacting factors of climate or weather. Psychological factors may play a part. Here lies work for the future. The first step consists, however, in defining what is meant by the loose expressions - damp cold and dry cold.

6. REFERENCES

1. RENBOURN, E.T. : The Natural History of Insensible Perspiration. A forgotten Doctrine of Health and Disease - Med. Hist. (in press).
2. HIPPOCRATES : The Genuine Works of Hippocrates - Trans. Adams, F., London, 1939.
3. STRATTON, G.E. : Theophrastus and the Greek Physiological Psychology before Aristotle. London, 1917.
4. GALEN : De Sanitate Tuenda. Galen's Hygiene - Trans. Green, R.M., Springfield, 1951.
5. CAUIS, J. : A Boke or Counsell against the Sweatinge Sicknesse etc, London, 1552.
6. BURTON, R. : The Anatomy of Melancholy - Dell, F. and Jordon-Smith, P., New York, 1938.
7. SANCTORIUS, S. : Ars de Statica medicina - 1614 Trans. Quincy, J. 2nd Ed., London, 1720.
8. BACON, F. : Novum Organum - 1620 Trans. Fowler, T., Oxford, 1878.
9. BOYLE, R. : New Experiments touching Cold etc., - London, 1665.
10. CULPEPER, N. : Culpeper's Last Legacy etc. - London, 1665.
11. WAINEWRIGHT, J. : A mechanical Account of the Non-Naturals etc. - London, 1707.
12. ARBUTHNOT, J. : An Essay Concerning the Effect on Air on Human Bodies - London, 1733.
13. SENAC, J.B. : Traité de la Structure du Coeur de son Action et de ses Maladies - Paris, 1752.
14. SAUVAGES, F.B. de : Dissertation comment l'Air, suivant ses différentes Qualités, agit sur le Corps Humain. - Bordeaux, 1753.
15. PRINGLE, J. : Observations on the Diseases of the Army - London, 1752.
16. CHANDLER, J. : A Treatise on the Disease called a Cold - London, 1761.
17. RENBOURN, E.T. : The History of the Flannel Binder and Cholera Belt - Med. Hist. 1, 211-225, 1957.
18. TISSOT, S.A. : Essai sur les Maladies des Gens du Monde - Lausanne, 1770.
19. COLOMBIER, J.C. : Code de Médecine Militaire pour le Service de Terre - Paris, 1772.
20. BLANE, G. : Observations on Diseases incident to Seamen - London, 1785.
21. RIGBY, E. : An Essay on the Theory of the Production of Animal Heat - London, 1785.
22. RENBOURN, E.T. : The Physiology of Textiles and Clothing. A Historical Note - J. Roy. Army Med. Corps, 99, 203-213. 1952-3.

23. WILlich, A.F.M. : Lectures on Diet and Regimen etc. - London, 1798.
24. HUFELAND, C.M. : Art de Prolonger la Vie Humaine 2nd Ed.- Lausanne, 1798.
25. COTTE, L. : Traité de Météorologie - Paris, 1774.
26. QUINCY, J. : Lexicon Physico-Medicum or a New Medical Dictionary - London , 1794.
27. BICHAT, M.F.X. : Anatomie Générale Appliqué à la Physiologie et à la Médecine Paris, 1802.
28. BUCHAN, W. : Buchan's Domestic Medicine Modernized, etc. - London, 1807.
29. EDWARDS, W.F. : Sur l'influence des Agens physiques sur la Vie - Paris, 1824.
30. BEAUPRÉ, M. : A Treatise on the Effects and Properties of Cold, with a Sketch, Historical and Medical of the Russian Campaign - Trans. Clendenning, J., Edinburgh, 1826.
31. STARK, J. : On the Influence of Colour on Heat and Odours - Phil.Trans. Roy Soc. London, 123,285-312, 1833.
32. COMBE, A. : The Principles of Physiology applied to the preservation of Health - Edinburgh, 1834.
33. CARPENTER, W.B. : Principles of Human Physiology, etc. 5th Ed. - London, 1855.
34. MARTIN, R. : The influence of Tropical Climates on European Constitutions London, 1856.
35. COULIER, Dr. : Experiences sur les Etoffes qui servent à confectionner les Vêtements militaires - J. de la Physiol. de l'homme et des Animaux 1, 122-138, 1858.
36. MOORE, W.J. : Health in the Tropics - London, 1862.
37. PARKES, E.A. : A Manual of Practical Hygiene, etc. 2nd Ed. - London, 1866.
38. PARKIN, J. : Climate.. Its Influence in the Production and Prevention of Phthisis - London, 1882.
39. THOMSON, S. : A Dictionary of Domestic Medicine. 24th Ed. - London, 1882.
40. JAEGER, G. : Die Normale Kleidung als Gesundheitsschutz - Stuttgart, 1880.
41. QUAIN, R. : A Dictionary of Medicine, New Ed. - London, 1894.
42. RUBNER, M., LEWSCHW, V. : Ueber den Einfluss der Feuchtigkeitsschwankungen unbewegter J auf den Menschen während körperlicher Ruhe - 29, 1-55, 1897.

PART II AN INVESTIGATION ON CLOTHED MEN UNDER INDOOR CONDITIONS

by

Dr. E.T. Renbourn, A.M. Reid and D.R. Gilling (Great Britain)

With statistical design and analysis by J. Draper

1. INTRODUCTION

The expression damp, humid or moist cold is used in a variety of ways. It may refer to the environment of habitation or to the climate and weather conditions arising during the winter in maritime temperature zones. The adjectives may refer to a high relative humidity, to the presence of precipitation - rain, sleet and damp snow - or to presence of wet conditions underfoot (wet cold).

Dry cold, on the other hand, has similarly not been adequately described. Meteorologists and climatic physiologists normally use the term to mean freezing weather conditions with frost or snow on the ground. However, owing to variability of weather, a wet cold day may be followed by a dry cold night. Even under freezing conditions, snow may change from being dry and powdery to a clinging dust which melts into clothing when either the ambient temperature rises or with warm conditions in a tent etc. In the same way, vapour saturated air may be breathed into a sleeping bag and condense; and, with heavy exertion, sweat will accumulate in the clothing from within, to evaporate (or wick), condense and freeze in the outer layers. Although under freezing conditions the atmospheric air contains very little moisture (at most about 4 mm Hg partial pressure) the relative humidity out of doors is itself very high, but usually very low in habitation, particularly if this is warm by American standards. Relative humidity under freezing conditions is a rather meaningless term - humidity is better expressed as absolute humidity in terms of vapour pressure.

"Dry cold" is also used popularly for a hypothetical weather condition corresponding in temperature to "damp cold" weather (sun, cloud, wind, barometric pressure, etc. usually being unspecified) but with low humidity. In point of fact, at temperatures between 32°F and 48°F, a relative humidity below about 40% does not often occur in nature. The greater chilling effect of damp cold is generally explained on the assumption that dry air is a better thermal insulator than damp air. The historical development of this idea has already been outlined. As pointed out recently by Guillemin (ref. no. 7) and by Burton (ref. no. 4), there is, in fact, no great difference in thermal insulation between dry and very humid air. Guillemin (ref. no. 7) also points out that convective cooling by dry and humid air at moderate speeds of flow (proportional to square root of heat conductivity x specific heat x density) is also not very different. It must, hence, be accepted that under still air or at moderate wind speeds, the purely physical effect of dry or damp cold air will not be very different. Woodcock (ref. no. 15) has recently given an excellent review of the theoretical relationships, near freezing point, of vapour and liquid water in air, through clothing layers and at the skin level.

Although there is no great factual evidence on the matter, what is popularly called "damp cold" weather may be associated with the interacting variables of a weather front - of a falling barometer and changing air temperature (say between 35-50°F), of generalised clouds and increased electrical state of the atmosphere, the presence of gusty winds and the likelihood of rain. "Dry cold" weather, on the other hand, may occur with a rising barometer, a stable and lower temperature, clear skies, little wind and bright sunlight. The feeling of "dry cold", in suitably dressed individuals, is related peculiarly to the extremities - hands and feet. That many people feel colder "all over", miserable and less active under so called cold humid weather, is undoubtedly true.

Apart from air movement and temperature, the independent physiological effects of any one meteorological condition are insufficiently understood. It has been assumed by some (ref. nos. 9, 11) that the factors constituting a weather front may, in some way yet unknown, have an effect on the higher nervous centres (with resulting changes in pulse rate, blood pressure, blood cells etc.); and a state of tissue water retention could, conceivably, explain many of the clinical phenomena (aching corns, rheumatic pains, headache, generalised skin discomfort) associated with the body acting as a "cosmic resonator" (ref. no. 11). Bright colours and a clear sky produce, in some individuals, a "feeling" of warmth. Perhaps, in the absence of sun, the presence of cloud and the thought of impending rain may, in a similar way, induce a "feeling" of cold or of damp. Much has been written on the effects of weather and climate on man, but how far some of the above phenomena are primarily psychological is not known.

Even under freezing conditions, solar radiation has an appreciable warming effect - both physiological and psychological. Clouds absorb the earth's long waves even more efficiently than the

solar short infra red radiation, but the effect of cloud cover on radiation heat loss from the clothed body (a matter considered some seventy years ago) is, however, likely to be negligible. Although, at low temperature, relative humidity of the air is not an important physical factor in convective heat loss, winds occur more commonly under so called "damp cold" weather conditions. Heat loss will also occur by rain impinging on the clothing, and, if absorbed, impairing its thermal insulation. It is now generally recognised that, under physical conditions, textile materials absorb water vapour with the liberation of appreciable quantities of heat (ref. nos. 1, 10); but little work has yet been done on the value of such "sorption" heat to the individual wearing wool in the form of garments.

A somewhat neglected factor in the problem of thermal comfort in cold weather, is that of shelter. In America, Canada and certain parts of Europe, a large proportion of the population live in winter under much warmer and drier indoor conditions (about 75-80°F and 25-30% R.H.) than those in England (about 60-65°F and a higher humidity), even with similar weather conditions. In the former countries, individuals put on warm dry clothing in the mornings over the already warm skin. Such conditioning may, perhaps, make them temporarily immune to the effects of cold weather whether damp or dry.

This report is concerned with the effects, under indoor conditions, of moderately low and high humidity at a low air temperature (36°F) and the modifying influence of warm and cool housing environments.

2. EXPERIMENTAL METHOD

EXPERIMENTAL CONDITIONS

Groups of men were conditioned to indoor environments corresponding, more or less, to those of an average "American" and "British" house during winter months, and then exposed to two indoor cold environments of the same temperature, air movement and radiation characteristics, but of different humidity. By choice, a temperature of 60°F would have been adopted for the "British" and 75°F for "American" conditions. However, owing to a sudden mild spell of weather during the period of the trial, and the difficulty experienced in maintaining a temperature of 60°F in the simple hot chamber available, it was decided to adopt a temperature of 65°F. In order to maintain a reasonable difference between the conditioning temperature, the "American" environment was raised to 80°F. The mean relative humidities of the two environments were "American" 37% and "British" 58%. The cold environment was produced in a chamber whose entry was within a few feet of that of the hot chamber. An average air temperature of 36°F was maintained. Technical difficulty was experienced in maintaining a constant relative humidity in the type of chamber in use, and the average conditions that could be maintained were 50% R.H. ("moderate humidity") and 85% ("high humidity"). Whilst the levels and constancy of humidity were not that desired, it is considered that the difference between the relative humidity does correspond to that likely to occur in nature at a temperature of 36°F and was sufficiently wide to serve as basis for preliminary investigation. In all cases the air movement was minimal (15 ft/min.). The effect of slight variability in air movement in different parts of the chambers was minimised by men occupying the same positions during all exposures. Wall temperature in the chambers was at air-temperature. Since it is possible that weather conditions to which men are exposed may conceivably affect the experimental results, the conditions out-of-doors, in the building housing the climatic chambers, and in the conditioning and cold environments, are given in Table 1 below:

TABLE 1 - ENVIRONMENTAL CONDITIONS

ENVIRONMENT		A.M.		P.M.	
		Temp. °F	R.H. %	Temp. °F	R.H. %
OUTSIDE:	Mean	50	78	57	69
	Range	42 - 59	52 - 100	52 - 67	33 - 100
IN BUILDINGS:	Mean	57	59	63	50
	Range	53 - 64	42 - 83	56 - 67	30 - 69
CONDITIONING ENVIRONMENT:					
(A) "American":	Mean	79.4	37	79.4	41
	Range	75 - 81	35 - 42	78 - 80	38 - 44
(B) "British":	Mean	65.7	58	66	58
	Range	64 - 69	55 - 66	65 - 70	53 - 70
COLD ENVIRONMENTS:					
(X) "Dry":	Mean	36.3	47	36.8	53
	Range	35.3 - 37.0	42 - 55	35.5 - 38.3	43 - 58
(Y) "Damp":	Mean	36.3	87	37.1	85
	Range	35.5 - 37.2	75 - 94	36.8 - 37.7	79 - 91

DESIGN AND ANALYSIS

The basic design of experiment consisted of a 4 x 4 Latin Square (see below) which allowed different orders of conditions to be balanced. On each morning, for 4 mornings of a week (Tuesday to Friday), a group of 4 subjects was used for the measurement of body temperature, clothing humidity and temperature gradients (three men only), and of subjective data. This was done for a total of 4 weeks (total 16 men). In the afternoons, another group of 4 men was used for nude body weight loss, respiratory ventilation volume measurement and subjective data. Over a 3 week period this gave a total of 12 men (with grand total of 28 men for subjective data).

DESIGNED BLOCK

Week	Day		Tuesday	Wednesday	Thursday	Friday	ENVIRONMENTS A = "American" B = "British" X = Moderate RH cold Y = High RH cold
	Subject						
1 AM PM	1 - 4 1a - 4a	BX	AX	BY	AY		
2 AM PM	5 - 8 5a - 8a	AX	AY	BX	BY		
3 AM PM	9 - 12 9a - 12a	AY	BY	AX	BX		
4 AM PM	13 - 16	BY	BX	AY	AX		

Groups of men were available for a period consisting of one preliminary day and four experimental days - the shortest period within which the subjects could experience all the environmental combinations. The object of the short period was to minimise the possible effects of outside weather and living conditions on the men. The first day of each week was used as an indoctrination period, during which men were exposed to the environmental combination of the first experimental day. Measurements were not taken, but subjects were able to become acquainted with the apparatus and procedures for the objective and subjective measurements.

The design of the experiment was such that the effects of the two conditioning environments upon the objective measurements obtained in the two cold exposures, and the effects of the latter alone, could be separately analysed. Means of AX, AY, BX, BY are all directly comparable.

Overall means for comparison of the two conditioning $\left\{ A = \frac{AX + AY}{2}, B = \frac{BX + BY}{2} \right\}$ and the two cold environments $\left\{ X = \frac{AX + BX}{2}, Y = \frac{AY + BY}{2} \right\}$ are given in the tables and figures below.

Strictly speaking only the differences between A and B, and between X and Y means have a physical meaning and no valid comparison can be made for example between A and X means.

Separate χ^2 tests were carried out on the subjective data obtained at the beginning, middle and end of the sessions in the cold environments. Morning and afternoon data were summated as there was insufficient information for separate analyses. Data from the three periods during a session were not combined, as three opinions from one subject, during one session, cannot be considered as independent. Such combinations would treat the data as if 3 x 28 i.e. 84 subjects had been used - an obviously fallacious assumption. Subjective data taken after men left the cold chamber were also treated by χ^2 analysis.

EXPERIMENTAL PROCEDURE

Fit young men of age 19-24 years were used as experimental subjects. The following clothing was issued to the man (as it had been shown that when wearing them they could remain in the cold chamber for two hours without shivering or severe discomfort): woollen jacket and trousers, woollen shirt, woollen vest and short woollen pants, woollen gloves and socks (total weight of garments 7½ lbs); heavy boots. No headgear was worn.

In order to simulate, with the facilities available, the changes occurring with sleeping indoors, the clothing was exposed overnight to the conditioning environment for the next day. On arrival at the laboratory in the mornings subjects visited the toilet, walked into the conditioning environment, removed their own clothing, and were equipped with harness for rectal and skin temperatures. The conditioned clothing, with incorporated elements for temperature and humidity gradients, was then put on. Men then sat quietly on chairs on which they remained for the whole experimental session (conditioning and cold exposure). This minimised physiological

changes (viz. Body temperature etc.) due to movement and variation in posture. The maximum experimental period (sitting) that could be permitted in the conditioning environment was 45 minutes. Subjects were then carried into the cold environment, at 6 minutes intervals, where they remained seated for a period of 102 minutes (again the longest period available).

In the afternoons, men entered the conditioning environment as above, were weighed nude before dressing in the conditioned clothing, and then sat down. Respiratory ventilation volumes were measured halfway through conditioning and cold periods. After being carried back to the conditioning environment, subjects undressed and were re-weighed. Otherwise the procedure was as for the mornings. During the experimental periods men were allowed to read or converse, but no other activities (smoking etc.) were allowed.

MEASUREMENTS

OBJECTIVE

BODY TEMPERATURE

Skin temperature was measured with thermocouples (soldered to copper gauze buttons and strapped on to the skin so as to leave the junction freely exposed) at 5 points (chest, shoulder, upper arm, thigh and calf) under the clothing (see fig. 1). Because of the interest in the face as a possible "trigger" factor in cold phenomena, cheek temperature was also measured with a thermocouple applicator. Rectal temperature was measured 10 cm along a flexible thermocouple element kept in position by a simple harness. Skin temperature is conventionally measured "near" the skin (on top of hairs etc.), only a few selected sites can be investigated at a time, and area "weighting" may not be very accurate particularly when rapid changes of internal temperature are occurring. It was, hence, considered of no particular value, in the present study, to estimate average skin temperature of the "whole body". For similar reasons (viz. local variation etc. in rectal temperature, ref. no.14), "mean body temperature", "heat storage" and "tissue insulation" were not used in the comparisons.

TEMPERATURE AND RELATIVE HUMIDITY GRADIENTS BETWEEN CLOTHING LAYERS (see fig. 2 and 3)

Humidity sensing elements of lithium chloride impregnated polyvinyl alcohol films were used, with which a thermocouple was incorporated (ref. no.12). The chest was considered a suitable point for investigation. An element was attached to the skin (about 1/10" from the skin) 2" below the left nipple, a second attached to the outer surface of the vest and another to outer surface of the shirt. The three positions were slightly staggered in order not to distort the gradient artificially. Readings were taken every 6 minutes in the cold environment.

RESPIRATORY VENTILATION VOLUME

This was measured with the Wolff "Integrating Motor Pneumotachograph" (IMP). Such measurement was used as an index of metabolic rate, and in the context of the present experiment (same men, same activity) was considered of sufficient accuracy for comparative purposes (ref. no.5). Sampling periods of 15 minutes were used during the middle of the acclimatising and cold sessions.

NUDE WEIGHT LOSS

Subjects were weighed nude (with the usual precautions) on a man balance, of accuracy ± 5 gm, on first entering the conditioning environment and as soon as possible on return from the cold environment.

SUBJECTIVE

The object of the trial and the purpose of the subjective phase was explained to the subjects on the first day of the trial, and at no time were they aware of the environments to which they were to be exposed. No questions were asked in the conditioning environment. Immediately on entering the cold environment, they were presented with the following questionnaire on a card:

- FACE - Does your face feel DRY OR MOIST ? WARM, COOL OR COLD ?
- BODY - Does your body feel COMFORTABLE OR COLD? If comfortable, WARM OR COOL ?
If cold, COLD OR VERY COLD ?
- FEET - Do your feet feel COMFORTABLE OR COLD ? If comfortable, WARM OR COOL ?
If cold, COLD, VERY COLD OR PAINFULLY COLD ?

In order not to be influenced by each other's results, subjects indicated their answer by a pointer. The whole questionnaire was presented again midway through the session and after its termination. Immediately on leaving the chamber, each subject was asked - "How did you feel in the cold chamber compared with yesterday ?"

3. RESULTS

OBJECTIVE DATA

All measurements for which variance analyses were made, showed, as expected a significant man to man variation - an individual type of response to temperature and to humidity exposure. Some of the man to man effects may, however, be due to other factors, viz. different positions in the chamber (see para.13). For none of the objective measurements was there a significant interaction between conditioning and cold environments i.e. the difference between the two conditioning environments could be considered the same for both "cold damp" and "cold dry" experimental environments.

EFFECT OF CONDITIONING ENVIRONMENTS ON COLD EXPOSURE

RECTAL AND SKIN TEMPERATURE

The mean initial and final rectal and skin temperatures are given in Table 2 below and the means taken every 24 minutes in Figures 4a and 5.

TABLE 2 - EFFECT OF CONDITIONING ENVIRONMENTS ON COLD EXPOSURE
(32 experiments per environment)

Part of the Body	Initial Body Temperature °F				Final Body Temperature °F			
	"British" Envir.	"American" Envir.	S.E. Diff.	Sig.	"British" Envir.	"American" Envir.	S.E. Diff.	Sig.
Rectal Temp.	99.1	99.2	0.40	Not Sig.	98.7	98.9	0.18	Not Sig.
Skin Temp.:								
Cheek	83.1	87.0	0.99	P < 1 %	70.7	71.0	1.22	" "
Chest	93.8	95.5	0.47	P < 1 %	91.9	91.4	0.41	" "
Shoulder	91.2	92.8	0.36	P < 0.1%	89.1	88.8	0.61	" "
Upper Arm	88.7	91.2	0.38	P < 0.1%	83.8	84.4	0.33	" "
Thigh	87.0	89.6	0.39	P < 0.1%	81.5	82.1	0.54	" "
Calf	81.0	83.1	1.34	Not Sig.	73.5	73.8	0.76	" "

Although there was no significant difference either in initial or final rectal temperature between the effects of the conditioning environments, it is to be noted that in both cases there was a mean rise of about half a degree between 6 and 30 minutes after entry into the cold, followed by a fall of about three quarters of a degree at 102 minutes. Differences between conditioning environments, either in rise or fall are, however, not significant. This point will be taken up later. Of the initial skin temperatures, all except that of the calf demonstrated significantly higher temperature for the warmer conditioning environment ("American"). As might be expected, the cheek (the only exposed skin site) showed the greatest mean difference (3.9°F). None of the skin temperatures taken after 102 minutes showed any significant differences - the warming effect on the skin had worn off.

TEMPERATURE GRADIENTS BETWEEN CLOTHING LAYERS

Owing to a number of "missing points" (due to difficulties in instrumentation) it was not possible to do a variance analysis of temperature gradients, but the means are shown in Figure 6a. It is seen that between each layer, considerable higher initial temperatures were obtained after conditioning at 80°F ("American") than after 65°F ("British"). Since the outer layers of clothing are in close contact with ambient temperature and temperature regulation occurs at the skin surface, it is to be expected, as was the case, the difference between the effects of the two conditioning environments, should be greater between shirt and battle dress tunic (7.3°F) and between vest and shirt (7.7°F) than between skin and vest (3.3°F). In contrast with the skin temperatures, it is seen that the difference in gradients, although reduced, persists throughout the period in the cold environment. It is not possible to attach levels of significance to these differences but, in comparison to the final skin temperature difference on the chest of 0.5°F, that between vest and shirt was 4.7°F and that between shirt and battledress 3.1°F. Since the conditioning environment differed in both temperature and humidity, it is not possible to separate these effects. Data from the humidity elements will be dealt with below.

RESPIRATORY VENTILATION VOLUME

The mean difference between the conditioning environments was "American" - "British" = 0.2 litres per man per minute, (S.E. = 0.62) not significant. One can assume that significant differences in the corresponding metabolic rates would not be present.

BODY WEIGHT LOSS

The mean weight loss per individual was 52 gm higher in the warmer environment, and, with S.E. = 9.8, this is highly significant ($P < 0.1\%$). The difference is due mainly to greater insensible perspiration (and, probably to light sweating) in the warmer environment. This result was expected.

EFFECT OF COLD ENVIRONMENTS

RECTAL AND SKIN TEMPERATURES

The means are shown in Figures 4b and 7 and the means of initial and final temperatures are given in Table 3 below:

TABLE 3 - EFFECT OF COLD ENVIRONMENTS
(32 experiments per environment)

Part of the Body	Initial Body Temperature °F				Final Body Temperature °F			
	High R.H.	Moderate R.H.	S.E. Diff.	Sig.	High R.H.	Moderate R.H.	S.E. Diff.	Sig.
Rectal Temp.	99.1	99.1	0.18	Not Sig.	99.0	98.7	0.18	Not Sig.
Skin Temp.:								
Cheek	85.2	84.9	0.98	" "	70.3	71.4	1.22	" "
Chest	95.2	94.1	0.47	$P < 5\%$	92.5	90.8	0.41	$P < 0.1\%$
Shoulder	92.5	91.5	0.36	$P \approx 1\%$	89.3	88.6	0.32	Not Sig.
Upper Arm	90.1	89.7	0.53	Not Sig.	84.5	83.7	0.95	" "
Thigh	88.8	87.8	1.28	" "	82.0	81.6	0.52	" "
Calf	83.0	81.1	1.34	" "	73.2	74.1	0.76	" "

As in the case of the effect of the conditioning environments, there was no significant difference in initial or final rectal temperature between "cold damp" and "cold dry" environments. There was no significant difference between environments in rise or fall of temperature.

It is seen from figure 7 that skin temperatures of chest and shoulder had almost reached steady values at 102 minutes, whereas those of cheek and limbs had not. Of the initial skin temperatures, those of chest and shoulder were respectively 1.1°F ($P < 5\%$) and 1.0°F ($P < 1\%$) higher for the cold damp environment, but other sites showed no significant differences. Only the chest showed a significantly higher (1.7°F, $P < 0.1\%$) final skin temperature in the cold damp. However, the drop in chest skin temperature during exposure to "dry" cold was 3.3°F with a drop of 2.7°F in damp cold, a difference which is not significant. From the physiological point of view we therefore, treat these results (and those of shoulder temperature) with some reserve. One would have expected the cheek to show the greatest difference between cold environments, since this was the only part of the skin exposed (i.e. nude skin).

TEMPERATURE AND RELATIVE HUMIDITY GRADIENTS

Figure 6b shows that, in contrast with the marked difference associated with the conditioning environments, differences in clothing temperature gradients between cold environments were negligible and clearly not significant.

Earlier work (Ref. no.12) had indicated that the humidity elements used in this study showed promise for use near the skin and between clothing layers. In the present trial, however, their performance was disappointing. While it was not possible to draw quantitative conclusions from the data, the results gave the following strong indications:

- under the same conditions there were marked differences between individuals;
- no consistent differences were observed between either the conditioning or cold environments;
- there was a marked fall in relative humidity at all positions, variable in rate and extent, on changing from the conditioning to the cold environments. This fall was contrary to first expectation, since the cooling of air in the layers of clothing, after entering the cold environments, would increase the relative humidity due to the water vapour already present.

This makes the assumption of still air conditions, but both the direction of the change in humidity, and its rapidity of onset, suggest a much brisker gaseous exchange between the surroundings and the clothing layers than was anticipated. Thus, the average vapour pressure of both conditioning environments was approximately 10 mm Hg, which (neglecting for the moment evaporation from the skin), on permeating to the layer between the skin and vest at a temperature of say 90°F, would give rise to a relative humidity of approximately 28%. In the cold environments, on the other hand, even at the high humidity, the vapour pressure was only about 4.6 mm, which at 90°F represents a relative humidity of 13%. This expected difference, based on the above figures, is of the same order as that actually found in the experiment.

While therefore, it was disappointing not to be able to obtain precise data on vapour pressure gradients under cold dry and damp conditions, the results indicate the presence of phenomena of real interest and the continuing need of a reliable method of measurement.

RESPIRATORY VENTILATION VOLUME

The mean differences in respiration ventilation volumes between high and moderate humidity was 0.8 litres per man per minute. With S.E. = 0.62 this is not significant. One may again assume no significant differences in corresponding metabolic rates.

BODY WEIGHT LOSS

The mean body weight loss per person between high and moderate humidity was 7 gm, and with S.E. = 9.8 this is not significant. It can be assumed that, in the cold, there is no significant difference in insensible perspiration loss between the moderate and high humidity. This is in contrast with the results obtained from the warm and cool conditioning environments.

SUBJECTIVE DATA

Replies to the questionnaire presented immediately on entering the cold environment showed no significant differences (χ^2 test) between any of the four combinations of conditioning and cold environments for face or feet. For the face, scores ranged from "warm" to "cold", but for the feet all were "warm" or "cool". Replies for the body were all "warm" or "cool" but there was a significantly greater proportion of "cool" replies ($P < 5\%$) in changing from the "British" conditioning to the moderate humidity than for the other three combinations, between which there was no difference. This is directly opposed to the popular impression of greater chilling in the damp. It is to be noted that the greater sensation of chilling in the cold at the moderate humidity, was present only when changing from the "British" and not from the "American" conditioning environment. The inference here is that in the latter case, any effect of humidity in the cold environment was masked by the reservoir of warmth acquired from the higher temperature of the "American" environment. There was no significant difference between any of the combinations of environment from the replies to the questionnaire put half way through the cold session. The range of replies was, however, extended to include some of "cold" and "very cold" for the body, and "cold" for the feet. With the final replies at the end of the cold session, there was no significant difference between any of the combinations of environment for the face or the body. With the feet, however, there were no differences between damp or dry cold, but in both of these the subjects were warmer after the "American" than after the "British" conditioning environment ($P < 2.5\%$). The replies for the feet extended from "warm" to "very cold", but none of the subjects was painfully cold.

It is worthy of note that no differences between any of the environments were found in sensation of the face. It was again thought that this, the only exposed site, might have been sensitive to difference in humidity immediately the subjects entered the chamber. In fact, however, the replies were very consistent throughout the session - 83 "dry", 28 "moist" at the beginning, and 85 "dry" and 27 "moist" at the end. It may be concluded that under the relative humidities and conditions of experiment, the subjects were not able to discriminate air of moderate or high humidity at a temperature of 36°F, the majority of replies for the face being "dry" for all conditions. It appeared, from the questionnaires taken after leaving the cold environments, that the opinions of thermal comfort, compared to that of the previous day, were more influenced by the conditioning than the cold environments.

4. DISCUSSION

In the last four years, three papers have been published dealing with the cooling power of damp and dry air at temperatures below 60°F. To facilitate discussion, details of these are given below together with those of the present study.

TABLE 4 - ANALYSIS OF INVESTIGATIONS ON COLD DAMP PHENOMENON

	A u t h o r s			
	Guillemin 1954 (Ref. no. 7)	Burton et al 1956 (Ref. no. 4)	Iampietro et al 1958 (Ref. no. 8)	Renbourn et al 1958
Cooled object	Nude and clothed metal cylinder	nude men	Nude men	Clothed men
Treatment before cold exposure	-	Unspecified	"Comfortable" Temp. 60 min.	"British" and "American" Envir. 45 min.
Cold Envir. Temp.	40°F and 50°F	48°F and 58°F	50°F and 60°F	36°F
R.H.	15% and 85%	34% and 80%	30% and 95%	50% and 85%
V.P. mm Hg.		10.0	12.6	2.75 4.6
Wind	1 and 4 mph	0	1 and 10 mph	0
Time	40 min.	100 min.	120 min.	102 min.
Posture and Activity	-	Horizontal	Unspecified	Sitting
Measurements:				
Temperature;				
Surface	+	+	+	+
Deep	-	+	+	+
Clothing	-	0	0	+
(R.H.)	-	0	0	(+)
Metabolism	-	+	+	+
Nude wt. loss	-	0	0	+
Subjective	-	+	+	+
Total duration Experiment for any subject	-	Up to 3 months	8 days	4 days
-, not applicable; 0, not done; +, done; (+), done, unsuccessful.				

Guillemin (Ref. no. 7) concluded that "for an inanimate test object simulating the thermal properties of a nude or clothed sedentary man, in windy or calm air, there is no significant difference in chilling effect between a dry cold and a moist cold environment". This agrees with the known physical properties of damp cold and dry cold air.

The other three papers are of a physiological nature and can be suitably dealt with together. Both Burton et al (Ref. no. 4) and Iampietro et al (Ref. no. 8) were concerned with the effect of humidity on nude men at temperatures of about 50° and 60°F; the present study deals with clothed men at a temperature near freezing point.

Design of experiment may be of some moment in laboratory experiments dealing with environmental factors, for, since men are ordinarily exposed during most of the day (and night) to uncontrolled indoor and variable outdoor weather conditions, this may, conceivably, modify their response during the shorter exposure to another environment, albeit in a controlled climatic chamber. The longer the total duration of such an experiment, the greater the probability of such extraneous source of variation. Such variation is minimised if groups of men (rather than individuals) are exposed together over the same short period of time. The size of a climatic chamber and the number of available men may, however, be critical factors. It has been shown that, after lying down in a constant environment, it may require $\frac{3}{4}$ - 3 hours before an apparent steady state of rectal temperature is reached (Ref. no. 14). For this reason, as long as possible a period of equilibration should be allowed to minimise possible sources of error outlined above. Although gentle movement has little obvious effect on rectal temperature, a change from horizontal to the vertical may cause the internal temperature to rise within 5 minutes and to continue to rise (by 0.5 to 1.0°F) for half an hour or more (Ref. no. 14). Furthermore, diurnal and other physiological variations of temperature may occur independently of an experiment (Ref. no. 14). For these various reasons, in investigations where change of internal temperature is likely to be small, it may not be possible to draw valid conclusions from the results (even if highly significant statistically) unless adequate control of experiment be used.

Although 45 minutes of conditioning to the "American" and "British" indoor environments produced no significant difference between them in rectal temperature on subsequent cold exposure,

there was significantly higher skin temperature, at all points, in men who had been in the warmer environment. Such men also showed higher temperatures between clothing at chest level. After 102 minutes in the cold, when the skin effect had worn off, the between-clothing temperatures continued higher. These results suggest that people sleeping indoors at what corresponds to dry summer heat, and putting on warm dry clothing in the morning, will be less affected by outside cold air (whether of high or low humidity) than those coming from a cool ("British") habitation or shelter. It is not certain, however, that the pre-warmed individual retains a physiological advantage after a few hours cold exposure. Our results correspond to those derived from a motionless individual under still air conditions. Under variable activities out of doors, the period of greater warmth may be appreciably less than the two or so hours suggested by the experiments. Individuals habituated to a "hot-house" (by British standards) in the winter, appear to be those who particularly complain of being "chilled to the marrow" when living in a cool house and exposed to temperate winter conditions. It may well be that our American and Canadian colleagues, who live in towns (and have heated vehicles etc.) do not ordinarily feel the cold because they are not often, or for long, exposed to it.

In all the three above physiological studies, exposure to cold was associated by a variable rise followed by a fall of rectal temperature. In the case of Iampietro et al and of ourselves, no significant difference between rise or fall of temperature was due to humidity itself. However, at an ambient temperature of 48°F, Burton et al found no change at high humidity but a rise of rectal temperature at low humidity of 0.36°F after 50 minutes (P=15%), and of 0.56°F (P=7%) at 100 minutes exposure. This greater rise (but hardly significant) in dry cold was ascribed to vasoconstriction (as calculated) and the small increase in metabolic heat production.

Earlier work in this laboratory was not able to confirm a consistent "paradoxical deep temperature phenomenon" on hot or cold exposure (viz. early fall on entry into the heat followed by a rise, and vice versa), and one of us has unpublished data showing that mental "apprehension" may be associated with a fall in skin and a rise in oral temperature (of mean about 0.5°F). There seems to be some evidence for such a mental state being more associated with cold than with warm exposure in a climatic chamber. In the present experiments the cold chamber contained far more scientific "hardware", with periodic irritating noises due to thermocouple switching gear. For these various reasons, we prefer, at present, to treat the mechanism of the rise of rectal temperature in the cold with some reservation. It is of interest that there appears to be little "awareness" of an appreciable rise or fall of internal temperature. This phenomenon has been noted before (Ref. no.14).

Burton et al found no difference in skin temperature between high and low humidity in the cold except in the case of the chest which was 2.5°F higher (P=4%) at low humidity. Buettner (ref. no. 3) has shown that when atmospheric vapour pressure is above 21 mm Hg, water vapour passes into the skin ("negative" insensible perspiration). This may conceivably be associated with liberation of "sorption" heat. These facts were used by Burton et al (Ref. no.4) to support the following argument; "the paradox of a greater sensation of cold and metabolic response at low humidity, though skin temperatures and heat loss were independent of humidity, could be due to the uptake of moisture in the skin at high humidity.." However, the temperature of vapour saturated air at a partial pressure of 21 mm Hg is about 73°F. With the available data, there is as yet no clear evidence that under subjectively "cold" conditions (as used in the three investigations under consideration) a high relative humidity will, per se, diminish evaporative cooling because of vapour absorption by the skin. There is some controversy in the literature as to the effect of high humidity on insensible perspiration loss by clothed men under warm air conditions (70°F - 80°F) (Ref. nos. 2, 16) but the general consensus of opinion is that it is increased. Insensible perspiration appears to decrease with cold conditions but this is less marked in clothed men (Ref. no.10a). The effect of relative humidity at low temperature is, however, not known. Iampietro et al found no difference in skin temperature between high and low humidity, but mean skin temperatures of whole body and not individual skin temperatures were examined. In our own case, no significant differences were found except at chest and shoulder on early exposure, with a highly significant difference at the chest at the end of the exposure, the higher temperatures being found at the higher humidity. This result (which we nevertheless treat with reserve, see Results) is in contrast to the findings from the warm conditioning environment, where, early in exposure, most skin points (particularly that of the face) showed significantly higher temperatures. A higher skin temperature in damp cold is, in any case, not likely to be explained by a decreased evaporative cooling of the skin, for, in our data, no significant difference was found between insensible perspiration loss in the cold at high or the moderate humidity. The negligible effect of humidity on skin temperature in general fits in with the finding of no subjective difference in chilling.

If, at 40°F, the "sorption" heat of wool is associated with the marked rise of temperature (8-9°F) observed by Fourt (Ref. no. 6) (using a fabric assembly of wool serge and wool underwear), it might be anticipated that some of this would be noted in resting men, wearing three layers of wool, even with the smaller humidity change used in the present experiment. However, no difference in clothing temperature gradients between moderate and high humidity was found at

chest level (the only skin site to show any consistent change). It is unfortunate that gradients at other sites could not be measured, and it is regretted that we were not able to obtain data on vapour pressure at the skin level and between clothing. Even under resting conditions, convection currents pass out from the body, chimney ventilation passes up from the chest and bellows ventilation of the trunk (due to respiration) occurs some twenty times a minute. This may, perhaps, partly explain our lack of evidence for "sorption" heat of clothing under physiological conditions. Out of doors, the air movement arising from marching and that of wind is to be considered. Our present data are few and preliminary, and more detailed work on the physiological problem of "sorption" heat is to be carried out in this laboratory.

As anticipated, Iampietro et al found a marked cooling action of wind (presumably streamlined in a wind tunnel) on the nude body, but with no significant effect arising from the relative humidity of the moving air. Early work in our laboratory (on the comparison of wind resistant and wind permeable garments) (Ref. no.13) showed no differences in skin temperature arising from turbulent air in a climatic chamber at 10°F or with winds out of doors at 40°F-60°F, yet with highly significant differences at 60°F in streamlined air, of similar speeds, in a wind tunnel. Since natural winds are usually turbulent, results obtained in a wind tunnel may somewhat exaggerate their physiological importance in the clothed man under temperate conditions.

The only differences found by Iampietro et al between damp and dry cold was a greater metabolism for nude men (3/4 experiments) in the latter environment, but there was no associated subjective difference. Our own results, however, show no such significant metabolic difference for clothed men.

It may be concluded that for clothed (and probably nude) men, under conditions of moderate cold, few physiological differences of practical importance are present due to the air humidity itself. The myth of chills arising from cold damp air and the fear of "the cold dews of the night" can, perhaps, after many centuries, be laid down; and to this can be added the old but still prevalent belief that damp air is a better heat conductor and convector than dry air.

Problems of body heat regulation will always require controlled laboratory experiments for their elucidation. But it is to be remembered that a hot chamber is not the equivalent of a hot climate, either biologically, physiologically, psychologically or even meteorologically; the same is fully true for a cold chamber. If differences are found between results derived from physical models in a conditioned cubicle, from nude men resting in a climatic chamber and from active people out of doors, it must be conceded that such differences may, in part, be due to the fact that phenomena of increasing complexities are being compared.

5. REFERENCES

1. BAXTER, S., CASSIE, A.B.D. : Rate of Propagation of Temperature Change through Textiles in humid Atmospheres. Part V. The Comparison of Different Textile Fibres - Trans. Faraday Soc., 37, 517-525, 1941.
2. BENEDICT, F.G., CARPENTER, T.M.: The Metabolism and Energy Transformation of Healthy Men during Rest - Carnegie Institute, Washington, 1910.
3. BUETTNER, K. : Diffusion of Water and Water Vapour through Human Skin - J. Appl. Physiol., 6, 229-241, 1953-4.
4. BURTON, A.C., SNYDER, R.A., LEACH, W.G.: Damp Cold versus Dry Cold. Specific Effects of Humidity in Heat Exchange of Unclothed Men - J. Appl. Physiol., 8, 269-278, 1955-6.
5. DRAPER, J., EDWARDS, R.G., HARDY, R.W.: Method of Estimating the respiratory cost of a Task by use of Minute-Volume Determinations - J. Appl. Physiol., 6, 297-305, 1953-4.
6. FOURT, L. : Temperature rise in Fabric Layers due to Heat of Regain under Cold-Moist Conditions - H.Q. Quartermaster Res. & Dev. Command, Textile Clothing & Footwear Division, Report No. 214, 1958.
7. GUILLEMIN, V. : Relative Chilling Effects of Moist Cold and Dry Cold Climates - J. Aviation Med., 25, 651-659, 1954.
8. IAMPIETRO, P.F., BASS, D.E., BUSKIRK, E.R.: Heat Exchanges of Nude Men in the Cold. Effect of Humidity, Temperature and Wind Speed - J. Appl. Physiol., 12, 351-356, 1958.
9. MILLS, A.L. : Climate makes the Man - New York, 1942.
10. NEIBACH, J.A., HERRINGTON, L.P.: Note on Hygroscopic Properties of Clothing in Relation to Human Heat Loss - Science, 95, 387-388, 1942.
- 10a. NEWBURGH, L.H., HARRIS, M. : C.A.M. Rep. Ann. Arbor, 1945
11. PETERSON, W.F. : The Patient and the Weather - Ann Arbor, 1, 111, 1936.

12. PHIPPS, L.W., RENBOURN, E.T., TAYLOR, P.F.: The Measurement of Relative Humidity and Vapour Pressure near the Skin - J. Physiol., 127, Proc. 46-48, 1955.
13. RENBOURN, E.T., PHIPPS, L.W., STOKES, A.W.: Unpublished Ministry of Supply Report, England, 1950.
14. RENBOURN, E.T., TAYLOR, P.F.: Unpublished Ministry of Supply Report, England, 1955.
15. WOODCOCK, A.H. : A Theoretical Interpretation of the Sensation of Damp Cold experienced by a Clothed-Man - Quartermaster Climatic Res. Lab. Environmental Protection Section, Report no. 199, America, 1953.
16. YAGLOU, C.P. : Physical and Physiological Principles of Air Conditioning - J. Amer. Med. Assoc., 108, 1708-1713, 1937.

ACKNOWLEDGMENTS

Our thanks are due to Mr. W.J.A. Humphries for the design and construction of automatic thermo-couple selector apparatus, and for the untiring and careful work carried out by Mr. C.A. Prigmore and Mr. G.W. Hicks over an arduous trial period of several weeks.

S U M M A R Y

Clothed men at rest were exposed to two conditioning "indoor" environments ("British", cool moist; "American", warm dry) for 45 minutes, and then to a cold environment (36°F) at moderate and high relative humidity (50% and 85%). The following measurements were taken: Temperature - skin, rectum and clothing gradients; nude body weight change and metabolism; subjective data. An attempt was made to measure clothing humidity gradients. The results showed that warm conditioning kept the clothing and skin at a higher temperature for a matter of $1\frac{1}{2}$ hours or more during cold exposure. The physiological value of such conditioning may be less marked under field conditions. No physiological objective or subjective differences of practical importance were found between the effects of still air at 36°F at moderate or high humidity, except for the finding of a slightly warmer skin of the chest at the high humidity. These results agree fairly well with those of other workers for nude men.

From the data obtained there is no clear evidence that "sorption" heat of wool is an important factor in keeping the body warm at resting conditions. Under field conditions, "sorption" heat might be of still less value. There is no basis for the supposition that moist air is either a better thermal conductor or a more efficient heat convector than dry air. The physiological difference between "dry cold" and "damp cold" is not due to properties of humid air, but to the associated weather or climatic differences (solar radiation, cloud cover, wind, barometric pressure, etc.) and, to some extent, to the mode of living indoors. Data from physical models and from physiological experiments in climatic chambers cannot be directly transposed into the realities of weather and climate out of doors.




Fig. 1 CLOSE-UP OF HUMIDITY ELEMENT
The outer threads are for protecting
the film



Fig. 2 HUMIDITY ELEMENT EXPOSED




Fig. 3 THERMOCOUPLE HARNESS
Arrows show position of gauze button,
thermocouples and wire leading to
rectal element.

FIG. 4.




FIG. 4. RECTAL TEMPERATURES.

- (a) EFFECT OF CONDITIONING ENVIRONMENTS ON COLD EXPOSURE.
(b) EFFECT OF COLD ENVIRONMENTS.
(32 EXPERIMENTS PER ENVIRONMENT).




FIG. 5 SKIN TEMPERATURES.
EFFECT OF CONDITIONING ENVIRONMENT ON COLD EXPOSURE.
(32 EXPERIMENTS PER ENVIRONMENT)




FIG. 6. TEMPERATURES BETWEEN CLOTHING LAYERS.
(a) EFFECT OF CONDITIONING ENVIRONMENTS ON COLD EXPOSURE.
(b) EFFECT OF COLD ENVIRONMENTS.
(24 EXPERIMENTS PER ENVIRONMENT)




FIG. 7. SKIN TEMPERATURES.
EFFECT OF COLD ENVIRONMENTS
(32 EXPERIMENTS PER ENVIRONMENT)

Section B : Social bioclimatology

1. Social bioclimatology (general)
2. Psychological bioclimatology (including
aestheto-bioclimatology)
3. Archeological bioclimatology

HUMAN BIOCLIMATOLOGY

Section B 2: Psychological Bioclimatology

PARTICIPATION OF SOME METEOROLOGICAL FACTORS
ON THE DETERMINATION OF EMOTIONAL REACTIVITY

by

J. Wellnerová and D. Svorad (Czechoslovakia)*

I. INTRODUCTION

It is widely accepted that the weather is in close connection to the behaviour of man, particularly to human emotionality (8, 2). Emotionality can, to a certain degree, be considered to be correlated with vegetative regulations at their highest levels in the hypothalamus (5). There were found two constitutional types of reactivity in rats (13; 15, 16, 17, 18). One of them is characterized by a vegetative, emotional and generally functional lability (rats functionally unstable or emotionally reactive, 19), the other functionally stable, emotionally unreactive. This finding has been affirmed substantially later (1). With regard to the mentioned connection between vegetative functions and emotionality on the one hand and human emotionality and meteorological influences on the other hand, we tried to find out the changes of emotionality occurring under the influence of temperature and air humidity on emotionally reactive (ER) and emotionally nonreactive (ENR) rats.

II. METHOD

Experiments were carried out on white male rats of the Wistar strain aged approximately 4 months. There were 9 - 11 animals in one experimental group.

Emotionality was tested in several ways: by Hall's tests (9, 10), by the modified open-field test (4) and by the timidity test (3). All these tests are based on the presumption that the exposure of the animal to a strange, fear-provoking situation can evoke the emotion of fear expressed by defecation, running (ambulation) or disability of hungry animals to take food.

HALL'S DEFECTION TEST OF EMOTIONALITY: The animal is exposed to a sound and light stimulus (2 bells, 2 buzzers, a 40 W bulb) for 2 minutes on four successive days in an arena 50 cm in diameter and height. Emotional defecation, expressed by the number of faecal boluses, was taken as the criterion.

HALL'S AMBULATION TEST OF EMOTIONALITY: The conditions are the same as in the previous test. The number of metres run by the animal on the floor of the arena, which is divided into equal squares, during an exposure lasting for 2 minutes, is counted.

THE MODIFIED OPEN-FIELD TEST: Food is placed in the middle of the arena. The time which the hungry animal spends on food during a 10 minutes exposure is taken down (maximum score 600 sec.).

TIMIDITY TEST: The cage with the hungry animal is placed at the end of a 12.5 cm wide and 125 cm long wooden runner, resting on two supports 90 cm high. At the beginning of the test the cage is placed in such a position which enables the animal to crawl into the runner. The time taken by the animal to reach the food at the far end of the runner is taken down. This time is the timidity measure of the animal. If the animal does not leave the cage for 35 minutes, the test is finished and the animal has an arbitrary score of 2000 sec. If the animal puts its two front paws outside the cage but does not leave it, the score is 1800 sec.

The animals were kept in surroundings at a temperature of 18° C and 53% relative air humidity. Emotionality tests were carried out under conditions of variable temperature (21, 27 and 33° C) and air humidity (50, 70 and 90%). Exposure of animals to a certain temperature and air humidity was always limited to the duration of the test.

*. Institute of Physiology of the Czechoslovak Academy of Sciences, Prague, Czechoslovakia.

III. RESULTS

1. EMOTIONAL DEFECTION: Fig. 1 shows that difference in the emotional score between ER and ENR rats is maintained by using different air temperature and air humidity. A temperature of 33°C at air humidities of 50% and also of 70% and 90% leads to an increased emotional score in ENR rats which at the higher humidities is at the level of the emotional score in ER rats. At temperatures of 21° C and 27° C, in conditions of variable air humidity there is a positive correlation $r = 0.69 \pm 0.08$ between the emotional characteristic of the animals (i.e. ER and ENR) and their defecation reactivity. It is further possible to deduct from the results of these experiments that an increase of temperature and air humidity is not followed by any significant changes in the defecational score in ER rats. Compared with the defecation score at 21° C this score is increased significantly in ENR rats at a temperature of 33° C and corresponding air humidities of 50% ($P < 0.001$), 70% ($P < 0.05$) and 90% ($P < 0.01$).

2. EMOTIONAL AMBULATION: Hall (10) found that there is an indirect connection between emotional defecation and emotional change of locomotor activity in the open-field test. The results of our experiments with emotional ambulation (Fig. 2) in ER and ENR rats show that a statistically significant difference in these two types of rats is maintained in ambulatory activity in several combinations of temperature and air humidity, but this was not significant at 50% air humidity and 33° C temperature, and at 90% air humidity and 27° C and 33° C. The figure shows further that increased temperature and air humidity raises also the emotional ambulation of ER and ENR rats. This increase is statistically insignificant, except for a significant increase at the level of $P < 0.05$ between 70% and 90% of air humidity at 33° C of air temperature in ER rats.

3. THE REACTIVITY IN THE MODIFIED OPEN-FIELD TEST: Contrary to the previous test situations, the influence of increased air temperature and air humidity on the emotional behaviour makes itself felt significantly in the open-field test. Emotional reactivity (or fear-reaction) of ENR rats increases significantly the more, the higher the temperature and humidity of air. In Fig. 3 the difference between the score of 50% and 90% of air humidity at 21° C is significant at the level of $P < 0.003$, at 33° C at the level of $P < 0.001$. Between the values of 21° C and 33° C of air temperature and at 50% air humidity there is no significant difference, but the difference is significant at the level of $P < 0.05$ at 90% air humidity.

4. THE RESULTS OF THE TIMIDITY TEST IN ENR RATS (Fig. 4): It is shown that air temperature and air humidity are decisive factors for emotional reactivity. The higher the temperature and humidity of air, the higher also the score of this test. The difference of score between 50% and 90% is significant at a temperature of 21° C at the level of $P < 0.001$ and at 33° C at the level of $P < 0.02$; on the other hand there is no significant difference between the score at 21° C and 33° C either at 50% or at 90% of air humidity.

IV. DISCUSSION

Emotional reactivity is a variable function determined by a number of factors (6). In an experimental situation where only air humidity and air temperature vary, reactivity changes significantly. Therefore it can be concluded that temperature and air humidity are also determining factors for emotional behaviour. The exposure of the animal to varying temperature and air humidity was in our experiments carried out over a relatively short time. It is, therefore, possible to presume that the accomplished results are only to a certain extent expressing the absolute values of temperature and air humidity. They could be the results of a gradient of change of these two meteorological factors (i.e. the transition from temperature and air humidity, in which the animals were kept, to the temperature and air humidity of the experimental situation). This sudden change of temperature and air humidity leads to an increased emotional unsteadiness of the animal which is not expressed equally in all tests of emotionality used. It is likely that each of the tests expresses a certain, reciprocally different component of emotionality. This opinion is confirmed by substantially similar results in the defecation and ambulation test which have a common test situation and differ only in the registration of a different (defecative or ambulatory) activity of the animal.

The presented results show further that the reactivity to air temperature and humidity is also an individually variable function. It is known (11, 17) that there are significant differences in the constitutional disposition of man to perceive meteorological changes. But these differences can disappear at a high temperature and air humidity as shown by the results of these experiments.

We presume that the facts about a different reactivity to meteorological influence on rats, which differ by their constitution, can bring new aspects to some psychopathological syndromes of human behaviour which can be studied on the experimental model of the so called animal hypnosis (14, 17). This reaction, called by Kretschmer (12) the phylogenetic precedent of human hysteria and other emotional disorders, is noticed in those very rats described as ER. By proving the reactivity of these rats to some meteorological influences, the way to a better understanding of the connection between human emotionality (or its disturbances) and some meteorological factors can be facilitated.

V. REFERENCES

1. BAYARD, J. : The Duration of Tonic Immobility in Guinea Pigs - J.comp.physiol. Psychol., 50, 130, 1957.
2. BERG, H. : Solar-terrestrische Beziehungen in Meteorologie und Biologie. (Probleme der kosmischen Physik, Bd.30) - Leipzig, 1957.
3. BINDRA, D. : What Makes Rats Hoard ? - J. comp.physiol.Psychol., 42,397,1949.
4. BINDRA, D., and THOMPSON, W.R.: An Evolution of Defecation and Urination as Measures of Fearfulness - J. comp. physiol.Psychol., 46,43, 1953.
5. BIRKMAYER, W. : Psychosomatische Medizin und neurovegetatives System - Acta neuroveg., 17, 131, 1957.
6. BROADHURST, P.L. : Determinants of Emotionality in the Rat. I. Situational Factors - Brit.J. Psychol., 48, 1, 1957.
7. CURRY, M. : Klimaunterschiede und wetterbedingte Reaktionstypen - Med.meteorolog.Hft., 5,39, 1951.
8. DUNBAR, F. : Emotions and Bodily Changes, 4.Ed. , New York, 1954.
9. HALL, C.S. : Emotional Behavior in the Rat. I. Defecation and Urination as Measures of Individual Differences in Emotionality - J. comp. Psychol., 18, 385, 1934.
10. HALL, C.S. : Emotional Behavior in the Rat. III. The Relationship between Emotionality and Ambulatory Activity - J.comp. Psychol., 22,345, 1936.
11. HOEHNE, K. : Konstitutions- und Reaktionstypen in der Bioklimatik - Med. meteorol.Hft., 5, 66, 1951.
12. KRETSCHMER, E. : Hysterie, Reflex und Instinkt, 6.Aufl., Stuttgart, 1958.
13. SVORAD, D. : To the Question of the Test of Seizure Susceptibility. 1st Congress of Czechoslovak Neurologists (Summary of papers in Czech) - Jeseník-Lázně, 1956.
14. SVORAD, D. : "Animal Hypnosis" (Totstellreflex) as Experimental Model for Psychiatry - A.M.A. Arch.Neurol.Psychiat., Chicago, 77,533,1957.
15. SVORAD, D. : Typology of Rats According to Inhibition Susceptibility. (Inhibition Susceptibility II) - Physiol.bohemoslov., 7, 348, 1958.
16. SVORAD, D. : The Question of "Inhibition Susceptibility". (Inhibition Susceptibility I.) - Physiol.bohemoslov., 7,341, 1958.
17. SVORAD, D. : Changes in Topographical Distribution of Glycogen in the Brain during Animal Hypnosis - Nature, 181, 775, 1958.
18. SVORAD, D. : Ueber die Beziehungen zwischen tagesperiodischen Veränderungen im Gehirnglykogen und der vegetativen Reaktivität bei Ratten - Acta neuroveg., 20, 33, 1959.
19. SVORAD, D., and WELLNEROVA, J.: Inhibition Susceptibility in the Light of so called Diencephalic Tests (In Czech) - Čs.fysiol., 7, 561, 1958.

SUMMARY

Rats were exposed to conditions of various temperature (21, 27 and 33° C) and air humidity (50%, 70% and 90%) and their emotional reactivity was estimated.

Increased temperature and air humidity increases more or less expressively the emotional reactivity. It was further stated that interindividual differences of emotional behaviour are maintained in conditions of lower levels of variously warm and humid air.




Fig. 1 Influence of various temperature and air humidity on emotional defecation. (Defecation score is expressed by an average amount of boluses during one test exposure).




Fig. 2 Ambulation under conditions of various temperature and air humidity. (Ambulation score: average amount of meters during one test exposure).




Fig. 3 Reactivity in the modified open-field test under conditions of various temperature and air humidity. (Score: average amount of minutes and its standard error).




Fig. 4 Reactivity in the timidity test. (Score: average amount of minutes)

HUMAN BIOCLIMATOLOGY
Section B 2: Psychological Bioclimatology

TEMPERATURE AND AIR HUMIDITY AS FACTORS INFLUENCING
SLEEP AND WAKEFULNESS

by

D. Svorad and J. Wellnerová (Czechoslovakia)*

I. INTRODUCTION

The rhythm of sleep and wakefulness is one of the basic biological factors. Although it is doubtful whether it is of a predominantly exogenic or endogenic origin (2, 13, 15), it is known that changes of environment, introduced gradually, do not disturb it greatly. On the other hand, the aperiodic influence of meteorological factors of a certain gradient can lead to its deep changes, expressed by insomnia (1, 4, 7).

It seemed useful to verify these empiric data by experiment because in that way it is possible to expose the organism to isolated meteorological factors and define quantitatively their biological significance (11). There are only limited possibilities of doing such an experiment on man, and it is more convenient to use animals.

The aim of this work was to find out the effect of temperature and air humidity on the rhythm of sleep and wakefulness, in certain experimental situations.

II. METHOD

Experiments were carried out on male rats of the Wistar strain at the approximate age of four months. There were 8 - 10 animals in one experimental group.

The rhythm of sleep and wakefulness was registered by an electronic actograph, i.e. a photo-electric circuit with amplifier brought into action by the spontaneous activity of the animal. The spontaneous activity was registered by an electric counter. The criteria of sleep and wakefulness depended on the frequency of spontaneous activity on the counter.

Insomnia was evoked by a system of relay circuits switching on a battery of acoustic, optical and vibrating stimuli at certain time intervals. By combining single stimuli with an electric shock into the paws of the animal, it is possible to elaborate a conditioned emotion of fear (6, 15) which, being continuously activated, does not allow the animal to fall asleep.

Air humidity was measured by a psychrometer. Further methodic data are given below.

III. RESULTS

1. CHANGES IN THE DURATION OF SLEEP AND WAKEFULNESS AT DIFFERENT TEMPERATURES AND AIR HUMIDITIES.

For 24 hours 4 groups of rats were exposed to different temperature and air humidity conditions based on the combination of two different kinds of temperature and air humidity. At the same time, the spontaneous activity of the animals was measured as the index of sleep and wakefulness. Fig. 1 shows that both at 21°C and at 27°C an increase of wakefulness, or decrease of sleep was present at a higher air humidity, compared with the values at lower air humidity. At the temperature of 21°C this difference was significant at the level of $P < 0.001$ and at 27°C at the level of $P < 0.01$.

The influence of temperature and air humidity on insomnia was followed in a further series of experiments. In all these experiments insomnia was evoked in the above mentioned manner for 24 hours at a temperature of 18°C and 53% air humidity. In the first part of the experiments animals were exposed to conditions of different temperature and air humidity for one hour before

* Institute of Physiology of the Czechoslovak Academy of Sciences, Prague, Czechoslovakia.

evoking insomnia and after that the influence of the exposure on insomnia was followed. In the second part of these experiments an exposure lasting one hour was carried out during insomnia. In both cases only temperature and air humidity of the exposure varied. The influence of the exposure on insomnia was measured by the sleep debt. Sleep debt (10) expresses the disturbance of the usual ratio of duration of wakefulness and sleep in the cycle of sleep and wakefulness of the given individual. It can occur if the usual duration of the phase of wakefulness is lengthened. In man it evokes a feeling of muscle lassitude and of being "below par". Individuals, accumulating such a deficiency of rest, may feel that they could return to "normal" by prolonging the duration of sleep in one single night (for example considerably prolonged sleep on Sunday mornings). Sleep debt in our experiments is not expressed by the length of sleep but by the lower intensity and frequency of spontaneous activity. In all our experiments we measured the sleep debt by one hour recording of spontaneous activity of the animal at a temperature of 18° C and 53% of air humidity. Since there is a rather great diurnal variation of spontaneous activity in rats, records were always taken between 7 and 8 p.m. The intact animal (i.e. one with an undisturbed rhythm of sleep and wakefulness) has an actographic value of 834 ± 196 at that time of day. After 24 hours of insomnia, ending always at 7 p.m., spontaneous activity was lower (i.e. sleep debt higher) than the one measured on an intact animal. Its values, as shown in Fig. 2 and 3, were always influenced by the temperature and air humidity of the exposure. These findings were also significant.

2. INFLUENCING SLEEP DEBT BY EXPOSURE TO VARYING TEMPERATURES AND HUMIDITIES BEFORE INSOMNIA.

In the first place, the animals were exposed to conditions of various temperature and air humidity (fig. 2) on a single occasion. The exposure lasted one hour. After that the animals were kept awake for 24 hours at a temperature of 18° C and 53% air humidity. After the end of insomnia their "need for sleep" was measured by sleep debt (in values of spontaneous activity at a temperature of 18° C and 53% air humidity). Fig. 2 gives the results of these experiments. It shows that sleep debt increased at the same ratio as the temperature and air humidity of the exposure. The increase of air humidity from 70 to 90% led to a statistically significant lowering of spontaneous activity (i.e. to an increase of sleep debt) at all temperatures used in the experiment. Likewise there was a significant difference between the height of sleep debt at 50 and 70% of air humidity at the temperature of 27 and also of 33° C. On the other hand, the increase of air temperature, e.g. from 27 to 33° C, had effect only at 90% air humidity ($P < 0.01$).

3. INFLUENCING OF SLEEP DEBT BY THE EXPOSURE APPLIED DURING INSOMNIA.

Insomnia, lasting for 24 hours, was evoked in the animals. After 12 hours of insomnia the animals were exposed to various degrees of temperature and air humidity for a single time. This exposure lasted for one hour (Fig. 3). In a further experimental variant, four exposures instead of one were carried out during insomnia (at approximately 5 hours intervals, each exposure lasting again 1 hour). The results of both these experiments (fig. 3) did not differ greatly and were practically the same as the results of the preceding exposure (Fig. 2). It follows that the higher the air humidity at a given temperature, the lower the spontaneous activity of the animal, i.e. the higher sleep debt.

IV. DISCUSSION

Results of the submitted experiments show the significant influence of both air temperature and air humidity on the rhythm of sleep and wakefulness.

Both studied stimuli differed basically from others influencing the organisms. In the first place, there is their continuous influence on the organism. Whereas other outside influences stimulate various receptors at periodic or aperiodic intervals, air humidity and air temperature act continuously on the organism. Thus they are continuous signals. The organism, aided by its regulative and compensative systems, working on the principle of an automatic control with feedback (3) can keep the homeostasis (9, 18). It can adapt itself to changes of its environment only to a certain extent. If the possibilities of its adaptability are overstepped, in our case the thermoregulative ones, a general reaction, i.e. a general escape reaction, sets in. To bring about an escape reaction, brain activity, not compatible with the state of sleep, has to be assumed. Therefore it is possible to presume that a drastic and continuous stimulus which does not allow the organism to protect itself by means of intelligent activity (clothes and ways of living, 8), will result in the disturbance of the normal rhythm of sleep and wakefulness. This is confirmed by the results of the submitted experiment.

The peculiarity of warm and humid air as a stimulus is given by the fact that these stimuli have an extraordinarily wide receptor field, presented by the whole surface of the body, and that the organism, therefore, has no possibilities to exclude the reception of these stimuli as is the case with stimuli attacking other exteroceptors and distance receptors. The result is a lasting

afflux of impulses which disturb the regulative mechanisms of sleep and wakefulness (12, 16).

An exposure lasting one hour, irrespective to the time of its application, is a stressor (14) which cannot be compensated by the organism. This explains the practically identical effect of the exposure preceeding insomnia and the exposure applied during insomnia on the one hand, and a single and repeated exposure on the other hand.

V. REFERENCES

1. AMELUNG, W., BECKER, F., BENDER, J. and PFEIFFER, C.A.: Störungen des vegetativen Nervensystems und Wettergeschehen (Neue Arbeitsmethoden meteoropathologischer Forschung) - Arch. physik. Therapie, 2, 181, 1950.
2. ASCHOFF, J. : Zeitgeber der tierischen Tagesperiodik - Naturwiss., 41, 49, 1954.
3. ASHBY, W.R. : An Introduction to Cybernetics - London, 1957.
4. BECKER, F., CATEL, W., KLEMM, E., STRAUPE, G., and KALKBRENNER, A.: Experimentelle Beiträge über die Beziehungen zwischen neuzeitlicher Bioklimatik und Medizin - Aerzt. Forsch., 3, 436, 1949.
5. BERG, H. : Exogener oder endogener 24-Stunden Rhythmus ? - Grenzgeb. Med., 2, 386, 1949.
6. BRADY, J.V. : The Effect of Electro-Convulsive Shock on a Conditioned Emotional Response: The Permanence of the Effect - J. comp. physiol. Psychol., 44, 507, 1951.
7. de RUDDER, B. : Grundriss einer Meteorobiologie des Menschen - Berlin, Göttingen, Heidelberg, 1952.
8. DIRNAGL, K. : Die Wetterempfindlichkeit als Zivilisationsschaden ? In Zivilisationsschäden und deren Physiko-balneo-klimato-therapie - Leipzig, 1957.
9. DRISCHEL, H. : Die vegetative Regulation als theoretisches Problem einer selbstständigen Regelung - Acta neuroveg., 6, 317, 1953.
10. KLEITMAN, N. : Sleep and Wakefulness - Chicago, 1939.
11. LANDSBERG, H.E. : Trends in Climatology - Science, 128, 749, 1958.
12. MAGOUN, H.W. : Ascending Reticular Activating System in the Brain Stem - Arch. Neurol. Psychiat. (Chicago), 67, 145, 1952.
13. MENZEL, W. : Endogene Änderungen des Tagesrhythmus bei Gesunden und Kranken - Med.-meteorol., 5, 28, 1951.
14. SELYE, H. : Physiology and Pathology of Exposure to Stress. A Treatise Based on the Concepts of the General-Adaptation-Syndrome and the Diseases of Adaptation - Montreal, 1950.
15. STEIN, L. : Reserpine and the Learning of Fear - Science, 124, 1082, 1956.
16. SVORAD, D. : Reticular Activating System of Brain Stem and "Animal Hypnosis" - Science, 125, 156, 1957.
17. SVORAD, D. and KOHOUT, L. : Method of Evoking Sleeplessness by Conditioned Emotional Reaction - Physiol. bohemoslov., in print.
18. WAGNER, R. : Probleme und Beispiele biologischer Regelung - Stuttgart, 1954.

SUMMARY

In experiments on rats we studied the influence of temperature (21, 27 and 33° C) and air humidity (50, 70 and 90 %) on the duration of sleep and insomnia.

The sleep of an animal, exposed to variously warm and humid air for 24 hours, is shortened the more, the higher the humidity and temperature of air. After insomnia lasting 24 hours and preceded by an exposure to variously warm and humid air for one hour, sleep debt is increased at the rate at which air humidity and temperature of the exposure is raised. Similar changes of the animal's sleep depth occur also in such cases where the exposure of the animal is carried out during insomnia.




Fig. 1 Spontaneous activity (wakefulness) of the rat in conditions of various temperature and air humidity. Spontaneous activity is expressed by values obtained from the electric counter. (Average value for one animal and its standard error).




Fig. 2 Sleep debt after a single exposure of the animal to conditions of variable temperature and air humidity. (A one hour exposure carried out before the 24 hours insomnia). Sleep debt expressed in values obtained from the counter of spontaneous activity (mean and its standard error).




Fig. 3 Sleep debt after a single and repeated exposure of the animal carried out during the 24 hours insomnia. White columns: a one hour exposure during insomnia. Black columns: four one hour exposures during insomnia.

Section C : Pathological bioclimatology

1. General pathological bioclimatology
2. Meteorological pathology
3. Climatological pathology
4. Air pollution pathology
 - a. Pollution with organic particles (pollen, fungi, etc.)
 - b. Pollution with inorganic particles (dust, etc.)
 - c. Chemical pollution
5. Geographical climatopathology
6. Climatotherapy
 - a. General Climatotherapy and Therapeutic Climates
 - b. Thalassotherapy (climatological-)
 - c. Heliotherapy
 - d. Thermotherapy
 - e. Aerosol therapy (and Iono therapy)
 - f. Socio-climatotherapy
 - g. Climatic health resorts
 - h. Other therapeutic methods

HUMAN BIOCLIMATOLOGY
Section C 6 e: Ionisation Therapy

A STUDY OF BIOCHEMICAL AND HAEMATOLOGICAL CHANGES
UNDER THE APPLICATION OF IONIZED AIR

by

Ludvík Erban (Czechoslovakia) *

I. INTRODUCTION

During the last decade a number of papers have been published in the medical literature dealing with the influence of ionized air on the organism. By ionized air we mean an atmosphere which contains electrically charged particles, consisting of single atoms, their groups (ionized molecules) or of bigger aggregations of different origin.

The first to call attention to the effects of ionized air, was the prominent Russian hygienist J.P. Skvortsov who studied, as early as about 1900, the influence of atmospherical electricity on human organisms. Later on, the subject was studied by Dessauer, Rajčevski, Sokolov, Strassburger, Happel, Frey, Michalovicz, Vasiljev, Tchijevsky, Busigina, Minch, Hicks, Hansell and others. Most of these papers concerned the examination of subjective sensations of test persons, the changes of blood pressure, respiration frequency, basal metabolism, chronaximetry, and, in some cases, also changes in the blood picture and sedimentation rate. In animals, changes of the total body weight, the weight of organs and growth of hair were investigated histologically, as well as changes in the thyroid and adrenal glands. The results obtained by different authors in applying positively and negatively ionized air, are very inconsistent. The experiments were carried out under different conditions. Different doses of ions were applied with different lengths of exposures and the ionized air contained a mixture of a smaller or greater predominance of positive or negative ions. Nevertheless most of the authors agree in the finding that negatively ionized air causes comfort, tranquilization, euphoria, and reduces blood pressure, while positively ionized air has opposite effects. These findings have promoted the use of ionized air for therapeutic purposes especially in the treatment of hypertension.

Although clinical, histological and physiological methods were used in the investigation of the influence of air ions, modern biochemical methods have not yet found sufficient application in this field. For this reason, as well as for their relatively high degree of objectivity, we have directed our attention particularly to these methods. Unlike most of the quoted authors, our studies were not limited to the influence of short time exposures but especially to the chronic effects of regular doses repeated over a prolonged period of time.

II. EXPERIMENTAL

The group of test persons consisted of men aged between 25 to 45 years. For technical reasons it was not possible to follow any group of control persons. The test persons inhaled artificially ionized air which was obtained by irradiating a current of atmospherical air by a radioactive substance placed in an apparatus. The device consists of a cylinder (about 8 cm. diameter) in which a thin plate of an alpha emitting radioactive substance is inserted. A constant current of clean air was introduced by means of a ventilator placed outside the experimental room. The ionization of air fluctuated round 4 million positive and negative ions in 1 ml. In the first series of experiments most of the negative, and in the second series most of the positive ions were eliminated by an electrostatic field, so that the test persons were inhaling predominantly positive ions in the first series of experiments and negative ions in the second series, the number of undesirable ions not exceeding 3 per cent of the total number.

The exposures lasted 1 hour and were applied to each person continuously 3 times a week, regularly for a period of 8 weeks in each of the both series. The second series was carried out with practically the same group of persons as the first one but about 3 months later.

* Institute of Hygiene, Prague, Director: Dr. K. Symon.

The following values were observed:

A. before and after each exposure: blood pressure
temperature
pulse

- B. once a week: 1. content of Na^+ , K^+ and Ca^{2+} in blood by a flame photometer.
2. total content of cholesterol colorimetrically on the principle of the Liebmenn-Burchardt reaction - the green colouring with acetanhydride and sulphuric acid - and particularly it's free as well as it's ester bound part with column chromatography.
3. total content of protein refractometrically.
4. content of separate protein fractions with paper electrophoresis of our own modification, using slips of paper freely stretched in the air, closed in a humid chamber.
5. content of the 17-ketosteroids in urine with the common Robbie Gibson method.

In addition, blood pressure and sedimentation rate controls were carried out weekly, for orientation.

III. RESULTS

The results obtained were statistically evaluated for each value under investigation so that the average values of parallel data, obtained in all test persons, were fixed and the amount of statistical dispersion was determined:

$$\sigma = \sqrt{\frac{\sum A^2}{n - 1}}$$

Further, with regard to blood pressure, also changes between the value before the exposure and after it were computed and evaluated in the same way. The average values and the marked limits of the dispersion were put into graphs, dependent on time.

IV. DISCUSSION

The first series of experiments with positive ions has shown that the blood pressure continuously rises throughout the whole experiment and this finding indicates a cumulation of effects. The rise is more significant in the systolic pressure that increased during the whole 8 weeks' experiment by 18 mm Hg (Graph 1).

Graph 2 shows the change of blood pressure during the exposure. This change is a kind of measure of sensibility of the blood pressure response to the applied dose of aeroions. It appears that the response of the organism becomes more expressive with repeated applications of positive aeroions. During the first ten days of the experiment the systolic pressure showed a slight decline, as the test persons mostly fell asleep after the beginning of the experiment. Although they used to fall asleep during the application of air ions after the tenth day too, the increase of their systolic as well as diastolic pressure above the values before the experiments appeared quite distinctly.

Graph 3 shows the changes of the total cholesterol and its fractions. The decrease of the total content of cholesterol is due substantially to the decline of the free cholesterol.

The total content of proteins in the serum is not influenced at all. On the contrary, the protein fractions show more striking shifts (Graph 4). While a decrease of albumins can be observed, signs of an increase appear in alpha, beta and gamma globulin.

In the course of inhaling positively ionized air, the content of the 17-ketosteroids in urine increases (Graph 5).

The blood picture did not show any changes and neither did the sedimentation rate.

The second series of experiments with negative ions that - according to literary statements - influence the organism favourably, has not shown any perceptible influence on the blood pressure (Graph 1), perhaps only an insignificant decline of the diastolic pressure.

As to the changes of the blood pressure during the exposure to negative ions (Graph 6), the systolic pressure shows a slight decrease which is however constant during the whole experiment. This is evidently the simple influence of tranquility or sleep. The diastolic pressure shows signs of falling, but the changes are within the limits of the dispersion so that it is possible that they are a matter of occasional fluctuation of results.

When negatively ionized air was applied, no changes either in the total content of cholesterol or in the content of its fractions were observed (Graph 7).

The total content of proteins in the serum did not change, not even when negative ions were

inhaled. Yet the protein fractions showed changes (Graph 8) which are opposite to those occurring under positive ionization. The albumins increased and globulins decreased.

The values of the 17-ketosteroids show an increase similar to those in the positive ionization but this increase is more moderate here (Graph 9).

The blood picture does not show any significant changes. It is only possible to observe an insignificant sign of a raise in the number of erythrocytes and a fall of the haemoglobin, but both these effects lie fully within the limits of the statistical dispersion. Also the mineral metabolism is not significantly affected, neither by positive nor negative air ions.

V. CONCLUSION

On the whole, the findings indicate an evident influence of the chronic administration of air ions. Especially positive ions show such an influence, where the whole picture of changes conspicuously reminds us of the complex of changes observed during the administration of ACTH.

Negative ions show much less definite effects. As the experiments were carried out on healthy individuals we had no occasion to ascertain if negative aeroions favourably influence a sick organism, as quoted in the literature. It is possible that the insignificance of the effects of negative ions in our experiments were due partly to the fact that the apparatus we had to our disposition did not emit ions of exclusively one sign. An admixture of a small number of positive ions could interfere with the effect of negative ions if it existed.

The existence of a chronic influence of aeroions under longterm administration, the signs of which have been ascertained in the above described experiments, will be subject of a further, more detailed, experimental study.



VI. SUMMARY

This paper deals with the study of the influence of artificially ionized air on human organisms. An attempt has been made to estimate objectively this influence by exact methods, especially by biochemical ones. A group of test persons inhaled positively ionized air during a period of two months at regular intervals (3 times a week during one hour) and they inhaled negatively ionized air during the second part of the experiment. The positive ionization caused an increase in the blood pressure. In blood proteins the albumins decreased and globulins increased. The total and the free cholesterol fell and the 17-ketosteroids increased. The negative ionization did not affect the value of the blood pressure and its effect was observed in the increase of albumins and the decrease of globulins. The other tests did not show any synonymous changes.

VII. REFERENCES

1. AYSOY, S. : Comment doit vivre l'homme a Ankara, "station à champ électrique positif" - Rev. Path. comp. 48, 584, 1946.
2. BRYCHTA, J. : Hygienicko-klimatický průzkum krajského města Hradec Králové - Sborník prací VLA 5, 51, 1953.
3. BUSIGINA, M.V., MINCH, A.A. : Vlijanije otricatel'no ionizovannovo vozducha na chododovuju recepciju koži - Gigijena 12, 5, 1956.
4. COUVIER, G. : La pression atmosphérique et le champ électrique en climatologie infantile - Presse therm. clim. 88, 145, 1951.
5. EDSTRÖM, G. : Studies in natural and artificial atmospheric electric ions - Lund, Sweden, 1935.
6. ENGHOFF, H., BÁRÁNY, E. : Zur Frage des Einflusses unipolar beladener Luft auf die Atmung - Acta med. scand. LXXXV, 184, 1935.
7. FREY, W. : Atmung und Luftionisation - Schweiz. med. Wschr. 80, 1387, 1950.
8. FREY, W. : Ionisation der Luft in geschlossenen Räumen - Schweiz.med.Wschr. 82, 994, 1952.
9. GROBSTEIN A KERSANOV : Lečeniye genuinnoj ozeny aeroioizaciej i prjamym prelivanyem krovi - Vest. Oto-rino-laring, 1, 51, 1948.
10. HOŘEJŠÍ, J. : Základy chemického vyšetřování v lékařství - Praha 1953.
11. HOŘEJŠÍ, J. : Bílkoviny krevní plasmy - Praha, 1956.
12. KANDRAČ, M.J. : Steroidní diagnostika - Bratislava, 1955.

13. KJUNCEL : Ingalcionnaja terapija ionizirovannym vozduhom - Vest. Otorino-laring, 2, 72, 1950.
14. MICHALOVICZ, M. : Význam stavu ionisace vzduchu ve fyziologii a patologii pediatrické - Pediat. pol. 83, 1954.
15. REISS, O. : Die Inhalationstherapie mit Luftionen - Arch. phys. Ther., 2, 386, 1950.
16. REITER, R. : Einige Ergebnisse einer halbjährigen luftelektrischbioklimatologischer Untersuchungen - Arch. phys. Ther. 3, 216, 1951.
17. ROHRER, E. : Ein Beitrag zur Frage der biologischen Wirkung ionisierter Luft - Schweiz. med. Wschr. 82, 1063, 1952.
18. SCHRÖRER : Ueber biologische Wirkungen ionisierter Luft - Schweiz.med.Wschr. 82, 350, 1952.
19. STEINKOPFF, Th. : Luftioneneinatmung - Physikalische Therapie, 119, 1952.
20. TCHIJEVSKY, A.L. : Die Wege des Eindringens von Luftionen in den Organismus und die physiologische Wirkung von Luftionen - Acta med. scand. LXXXIII, 219, 1934.
21. VASILJEV, L.L. : Teorija i praktika lečenijsa ionizirovannym vozduhom - Leningrad 1951.
22. YAGLOU, BENJAMIN, BRANDT : The influence of respiration and transpiration on ionic content of air occupied rooms - J. industr. Hyg. XV, 8, 1933.
23. YAGLOU, BRANDT, BENJAMIN : Observation on a group of subject before during and after exposure to ionized air - J. industr. Hyg. XV, 341, 1933.




Graph 1.

above: changes of the blood pressure during the application of positively ionized air measured at the end of each inhalation.


below: changes of the blood pressure during the application of negatively ionized air measured at the end of each inhalation.

(Averages with marked dispersion)




Graph 2.

Sensibility of the blood pressure to aeroions. Differences of the blood pressure before and after the inhalation of positively ionized air with marked dispersion.



Graph 3.


Changes of the total cholesterol and its fractions in dependence on the duration of the positively ionized air inhalation.
(Averages with marked dispersion)



Graph 4.

Electrophoretic analysis of blood proteins on the filtration paper. Changes in the protein spectrum in test persons inhaling positively ionized air during 2 months.


(Averages with marked dispersion)



Graph 5.


Increase of the 17-ketosteroids in urine in the course of the inhalation of positively ionized air.

(Averages with marked dispersion)




Graph 6.

Sensibility of the blood pressure to negative aeroions. Differences between the blood pressures after the inhalation of artificially ionized air with marked dispersion.




Graph 7.

Cholesterol and its fractions in dependence on the duration of the negatively ionized air application.
(Averages with marked dispersion)



Graph 8.

Changes in the protein spectrum ascertained by means of electrophoretic analysis of the blood proteins on the filtration paper in test persons inhaling negatively ionized air during 2 months
(Averages with marked dispersion)



Graph 9.

Values of the 17-ketosteroids in the course of the inhalation of negatively ionized air.
(Averages with marked dispersion)

Section D : Urban bioclimatology

- 1. General urban bioclimatology**
- 2. Architectural bioclimatology**
- 3. Sanatorium bioclimatology**

Section E : World literature

HUMAN BIOCLIMATOLOGY

Section E: World literature *

GENERAL PHYSIOLOGICAL BIOCLIMATOLOGY

1. ALVAREZ, A.M. : Studies of the relationship between atmospheric phenomena and human physiology - Puerto Rico Journ. of Publ. Health and Trop. Med., 9, 2, 1933.
2. ALVAREZ, A.M. : Puntos aparentes de contacto entre la marcha diaria de los componentes magneticos terrestres, la de ciertos elementos solares, la presion diastolica humana y los computos totales leucocitarios - Puerto Rico J. Publ. Health and Trop. Med., 10, 3, 374, 1935.
3. AMELUNG, W. : Luftkörper und Blutdruck - (Zusammen mit Prof. F. Bauer). Zeitschrift d. angew. Bäder- und Klimaheilkunde, 2, 1955.
4. AVANZO, G. D' : Influenza del Clima tropico-equatoriale sulla Glicemia, Polso, Pressione arteriosa, e Temperatura corporea durante la navigazione transoceanica - La Medicina del Lavoro, 6, 7, 392, 1954.
5. BERNSTEIN, S.H. : Effects of housing on incidence and spread of common respiratory diseases among Air Force recruits - School of Aviation Medicine USAF, Report 57-81. June 1957.
6. BERRY, L.J., BEUZEVILLE FERRO, C. and KRUMDIECK BOIT, C.: Metabolic studies of Guinea Pigs native to the High Andes and to the Peruvian coastal plain - School of Aviat. Med., USAF, Randolph AFB, Texas, Sept. 1957.
7. BONTKE, E. and BEHMANN, F.W.: Steuerbare Hypothermie durch intravasale Kühlung beim Hund - Ber. ges. Physiol. u. Pharmakol., 172, 124, 1955.
8. BREUER, K.H. : Beitrag zur Frage meteorischer Einflüsse auf die Leukocythenzahl - Diss. med. Kiel, 1950.
9. BROWN Jr., E.B. : Breath holding at altitude and breath holding as a demonstration of Anoxia - Navy Research Report X-431 (Ac-228-f) October 1944.
10. BROWN, E.B., CLARK, R.T. and HOFFMAN, C.E.: Blood oxygen saturations and duration of consciousness in anoxia at high altitudes - The American Journ. of Physiology, 145, 5, 685, 1946.
11. BROWN Jr., E.B., SMEDAL, H.A. and HOFFMAN, C.E.: Incidence of Benda pain in a short exposure to simulated altitudes of 26.000, 28.000 and 30.000 feet - Journal of Aviation Medicine, 17, 67, 1946.
12. BROWN Jr., E.B. : Diseases associated with low capillary resistance - Am. Heart J., 34, 2, 241-248, 1947.
13. BROWN Jr., E.B., CAMOBELL, G.S., JOHNSON, M.N., HEMINGWAY, A. and VISSCHER, M.B.: Charges in response to inhalation of CO₂ before and after 24 hours of hyperventilation in Man - J. of Applied Physiol., I, 333, 1948.
14. BROWN Jr., E.B., HEMINGWAY, A., CAMPBELL, G.S., GOLLAN, F., and ELAM, J.O.: Chronic Hyperventilation in normal human subjects - Fed. Proc., 7, 52, 1948.
15. BROWN, E.B., CAMPBELL, S., ELAM, J.O., GOLLAN, F., HEMINGWAY, A. and VISSCHER, M.B.: Electrolyte changes with chronic passive hyperventilation in man - Jour. of Applied Physiol., 1, 12, 1949.
16. BROWN, E.B., HEMINGWAY, A., and VISSCHER, M.B.: Arterial blood pH and pCO₂ changes in response to CO₂ inhalation after 24 hours of passive hyperventilation - J. of Applied Physiol., 11, 2, 1950.
17. BROWN Jr., E.B. : Changes in brain pH response to CO₂ after prolonged hypoxic hyperventilation - J. Appl. Physiol., 2, 549, 1950.
18. BROWN Jr., E.B., VISSCHER, M.B. and LIFSON, N.: Experimental physiology - Minneapolis, Burgess Publishing Comp., 1950.
19. BROWN Jr., E.B. and MILLER, F.A.: Ventricular fibrillation following a rapid fall in alveolar carbon dioxide concentration - Am. J. Physiol., 169, 56, 1952.
20. BROWN Jr., E.B. and BILLINGS, H.H.: Splenic contraction and blood volume changes produced by high carbon dioxide breathing - Proc. of the XIX Intern. physiological congress, 19, 239, 1953.

* References received from members by Secretariat of I.S.B.B. since 1957.

21. BROWN Jr., E.B. : The physiological effects of hyperventilation - *Physiol.Reviews*, 33, 445, 1953.
22. BROWN Jr., E.B. : Plasma potassium changes produced by inhalation of 30% and 40% carbon dioxide in the dog - *Fed. Proc.*, 14, 21, 1955.
23. BUETTNER, K. : Radiation effects on man in space - *Bull. Amer. Meteorol. Soc.*, 32, 183, 1931.
24. BUETTNER, K. : *Physikalische Bioklimatologie* - Akadem. Verlagsges., Leipzig, 1938.
25. BUETTNER, K. : *Spektrale Empfindlichkeit und klimatische Verwendung der Kadmmzelle* (with E. Sutter) - *Strahlentherapie* 45, 445-456, 1932.
26. BUETTNER, K. : Wärme- und Strahlungshaushalt des Menschen und des Frigorimeters - *Strahlentherapie*, 48, 386, 1933.
27. BUETTNER, K. : Zur Physik der Bestimmung der menschlichen Energiebilanz im Zimmer und im Freien - *Verh.D.Ges.inn.Med. XLV. Kongress*, Wiesbaden, 369, 1933.
28. BUETTNER, K. : Die Wärmeübertragung durch Leitung und Konvektion, Verdunstung und Strahlung in Bioklimatologie und Meteorologie - *Veröff. Preuss. Meteor. Institut. (Habilit. Schrift)*, 1934.
29. BUETTNER, K. : Bedeutung und Messung der Oberflächenfeuchte für die Transpirationsanalyse - *Biolog. Zentralbl.*, 55, 356, 1935.
30. BUETTNER, K. : Feuchte und Verdunstung der menschlichen Haut - *Forsch. u. Fortschr.*, 12, 151, 1936.
31. BUETTNER, K. : Ueber die Wärmestrahlung und die Reflexionseigenschaften der menschlichen Haut - *Strahlentherapie*, 58, 345, 1937.
32. BUETTNER, K. : Erythembildung durch Sonnen- und Himmelstrahlung, *Strahlentherapie*, 61, 610, 1938.
33. BUETTNER, K. and KUHN, H.: *Climates of the German Theaters of Operations. in German Aviation Medicine* - Dept. of the Air Force, Washington D.C., 1950.
34. BUETTNER, K. : Physical aspects of human bioclimatology - *Compendium of Met. (Amer. Meteorol. Soc.)*, 1951.
35. BUETTNER, K. : Human bioclimatology and energy in arid and humid tropical zones - *Proc. conference on tropical housing, Univ. of Texas*, 1952.
36. BUETTNER, K. : *The aeropause* - Science, 1952.
37. BUETTNER, K., REISMANN, K., and TOPKA, E.Z.: Influence of solar activity, especially at high altitudes on the Takata-flocculation reaction in serum of normal persons - *Arch.Met.Geoph. und Biokl.*, 5 (B), 103-111, 1953.
38. BUETTNER, K. : The change of short- and long-wave albedo with zenith angle - *Bull. Am.Met. Soc.*, 35, 446, 1954.
39. BUETTNER, K. : Water reabsorption by soil and skin - *Lecture held on 31 Aug. 1956, Symp. on Bioclimatology at UNESCO H.Q., Paris.*
40. BUETTNER, K. and ODLAND, F.: Diffusion resistance of the skin barrier layer in vitro and in vivo - *Fed. Proc.*, March 1957.
41. BUETTNER, K. and GEORGE, F.: Physical factors of the skin barrier layer and water diffusion into human skin - *Fed. Proc.*, 16, 1, 1957.
42. BUETTNER, K. : Present knowledge on correlations between weather charges, sferics and air electric space charges, and human health and behavior - *Fed. Proc.*, 16, 2, 1957.
43. BUNGE, E. : Verlauf der Dunkeladaptation bei Sauerstoffmangel - *Arch.f. Augenhk.*, 110, 189, 1936.
44. BURCH, G.E. and WINSOR, T.: The role of the skin and corneal layer in edema formation - *The Journal of the American Medical Association*, 163-164, 1944.
45. BURCH, G.E. and WINSOR, T.: Diffusion of water through living and dead human skin with the identification of the diffusion inhibiting layer - *The Journal of Clinical Investigation*, 23, 937-938, 1944.
46. BURCH, G.E. and WINSOR, T.: Differential roles of layers of human epigastric skin on diffusion rate of water - *Archives of Internal Medicine*, 74, 428-436, 1944.
47. CARLSON, L.D. : A concise description of the demand oxygen system - *Air Surg. Bull.* 1, 14-17, 1944.
48. CARLSON, L.D. and WULFF, V.J.: Pulmonary ventilation of flyers at high altitudes- 1945.
49. CARLSON, L.D., LOVELACE, W.R. II and WULFF, V.J.: Oxygen consumption of aircrews - *Air.Surg. Bull.*, 2, 304-305, 1945.
50. CARLSON, L.D. and WULFF, V.J.: The application of basic physiologic data in the design of respiratory equipment - *J. Aero Science*, 14, 229-236, 1947.
51. CARLSON, L.D., MARTIN, A.W. and GATTONE, V.: An apparatus for the measurement of pulmonary function (abstract) - *Fed. Proc.*, 7, 18, 1948.
52. CARLSON, L.D., MARTIN, A.W. and BARK, R.S.: A method for studying reflex activity under varying conditions (abstract) - *Fed. Proc.*, 7, 18-19, 1948.
53. CARLSON, L.D., and WOLFE, W.A.: Studies of pulmonary capacity and mixing with the nitrogen meter - *J. Clin.Invest.*, 1568-1575, 1950.

54. CARLSON, L.D. : Respiratory exchange - Methods in Med. Res., 6, 60-73, 1954.
55. DANTI, M. : Blood-coagulation and external phenomena - Wetter und Leben, 7, 7, 1955.
56. DAUBERT, K. and ZIPP, H.: Ist die Pupille als Testorgan für biometeorologische Beobachtungen geeignet? - Münch.med.Wschr., 96, 11, 294-296, 1954.
57. DEGENHARDT, K.H. : Analysis of intra-uterine malformations of the vertebral column induced by oxygen deficiency in rabbits - X International Congress of Genetics, Proc., 2.
58. DEGENHARDT, K.H. : Tierexperimentelle Untersuchungen zur Aetiologie und Phänogenese (früh erworbener) axialer Fehlbildungen - Z. Mensch. Vererb.u. Konstitutionslehre, 34, 509-515, 1958.
59. DIRNAGL, K. und KANZ, E.: Wetterwirkungen auf die menschl. Physis u. Psyche als Ergebnis experimenteller Untersuchungen - Archiv.f. Hygiene und Infektionskrankheiten, 135, 117, 1951.
60. DIRNAGL, K. : Zur fortlaufenden Messung des Atemwiderstands - Z. Aerosolforschg.u. Therapie, 3, 475-484, 1953.
61. DIRNAGL, K. : Wirkung elektrischer Felder auf den Menschen - Münch. Med. Wschr. 96, 19, 564, 1954.
62. DIRNAGL, K. and STIEVE, F.E.: Experimentelle Untersuchungen über die Wirkungsweise der pneumatischen Kammern - Arch.Phys.Thor., 6, 2, 115, 1954.
63. DIRNAGL, K. : Bericht über Messungen am "Pari-Optimal-Vernebler" Modell Primus - Firmenschrift des Pari-Werkes, 1954.
64. DIRNAGL, K. and PICHMAYER, H.: Untersuchungen über den Einfluss der Atemtechnik auf die Resorption inhalierter Substanzen - Ztschr.f. Aerosol Forschg. u. Ther. 3,3, 240, 1954.
65. DIRNAGL, K., DREXEL, H. and QUENTIN, K.: Zur Resorption von Sulfat durch die Haut - Die Naturwissenschaften, 41, 16, 376, 1954.
66. DIRNAGL, K. and WELCKER, H.: Untersuchungen über den Einfluss der Tröpfchengrösse auf die Resorption inhalierter Nebel - Z. Aerosol.Forschg., 4, 4, 289 - 296, 1955.
67. DIRNAGL, K. and ESCHÉ, R.: Ultraschall Raumvernebler - Siemens Ztschr., 29, 9, 382, 1955.
68. DRISCHEL, H. : Die vegetative Regulation als theoretisches Problem einer selbsttätigen Regelung - Acta Neuroveget., 6, 317-346, 1953.
69. DRISCHEL, H. : Ueber Norm- und Fehleinstellungen vegetativer Regulationen - Aerzt. Forschg., 8, 1, 17-25, 1954.
70. DRISCHEL, H. : Organismische Reglertheorie - Vortrag gehalten am 5.7.54 auf der Tagung "Biophysik" der Deutschen Physikalischen Ges. in Berlin. Arbeitstagung Biophysik, VEB Deutscher Verlag der Wissenschaften, Berlin, 1955.
71. DRISCHEL, H. : Dynamik vegetativer Regelvorgänge - Regelungsvorgänge in der Biologie. Verlag München, 1956.
72. DRISCHEL, H. : Probleme der vegetativen Regulation (Eine organismische Reglertheorie) - Vortrag, gehalten am 7.3.56 im Aerztlichen Verein in Bremen.
73. DUGAL, L.P. and HEROUX, O.: Effets de l'acide ascorbique sur l'hypertension provoquée par desoxicorticostérone (DCA) - Rev.v.de Biol., 10, 123, 1951.
74. DUGAL, L.P. and HEROUX, O.: Effet de l'acide ascorbique sur l'hypertension expérimentale - Can.Jl. Med.Sc., 29, 4, 164, 1951.
75. DUGAL, L.P. and THERIEN, M.: Studies on the relations between ascorbic acid and adrenal function - Proc. Can. Physiol. Soc. Rev. Can. Biol., 2, 86, 1952.
76. DUGAL, L.P. and THERIEN, M.: Effects of ascorbic acid on the adrenal weight of normal and hypophysectomized rats - Science, 115, 598, 1952.
77. EFFENBERGER, E. : Einfluss des Wettergeschehens auf die Tonuslage des vegetativen Nervensystems - Med. Meteorol.Hefte, 3, 10-20, 1950.
78. GRUENEWALD, G., MUECHER, H. and UNGEHUEER, H.: Beiträge zur Analyse psychischer und psychologischer Wetterwirkung - Editio Cantor, Aulendorf/Württ.
79. HALHUBER, M.J. : Der Kreislaufgesunde und -kranke im Hochgebirgsklima - Nauheimer Fortbildungs-Lehrgänge, 20, 136-142, 1954.
80. HAUS, E. and JUNGSMANN, H.: Kreislaufreaktionen bei kurzdauerndem Höhenwechsel in den Alpen - Schw. Med. Wschr., 83, 1156, 1953.
81. HAUS, E. and JUNGSMANN, H.: Kreislaufreaktionen bei Bergbahnfahrten in den Alpen - Med. Klinik, 49, 775, 1954.
82. HAUS, E., BEYRER, K., HALHUBER, M. and OLBRIICH, E.: Vergleichende Kreislaufuntersuchungen bei Heissluftinhalation und Lichtbädern - Ztschr.Bäder- u. Klimaheilkunde, 1, 316, 1954.
83. HAUS, E. and JUNGSMANN, H.: Höhenreaktionen des Kreislaufs bei Gesunden und Kranken in den Alpen - Schw. Med. Wschr., 84, 1265, 1954.
84. HENDEL, W.D. : Studien über die Indikanausscheidung im Harn unter dem Einfluss klimatischer und balneärer Wirkungen - Diss. med., Kiel, 1941.

85. HEROUX, O : Influence de l'acide ascorbique sur la pression artérielle en fonction du cortex surrénalien - Laval Médical Extrait, 16, 7-100, 1951.
86. HERRMANN, R. and HEINKE, E.: Die Beziehungen zwischen den pH-Werten des Luftkondensats und denen der Hautoberfläche - Arch.f. Dermat., 191, 501, 1949.
87. HERRMANN, R. : Schwankungen des pH's der Haut und des Aerosols - Verhandl. der Balneol. Klimat. u. Rheumat., 7, 1950.
88. HERRMANN, R. and HEINKE, E.: Die Abhängigkeit der Aciditätsverhältnisse der Hautoberfläche von denen der Nebelkerne - Arch.f.Dermat., 190, 261, 1950.
89. HERRMANN, R. a.o. : Das Verhalten der Elektrolyte (Na.K. u. Ca.) im menschlichen Serum unter dem Einfluss des Dampfbaden - Arch.f.phys. Ther., 4, 423, 1952.
90. HERRMANN, R. and HEINKE, E.: Die Beeinflussung der K- und Ca-Konzentrationen im Serum durch künstliche Ozonisierung der Raumluft - Strahlenther., 87, 473, 1952.
91. HERRMANN, R. and HEINKE, E.: Die Beeinflussung der K- und Ca-Konzentrationen im Serum durch künstliche Ozonisierung der Raumluft - Heilbad und Kurort, 4, 41, 1952.
92. HERRMANN, R. and HEINKE, E.: Die Beeinflussung der K- und Ca-Konzentrationen im Serum durch künstliche Ozonisierung der Raumluft - Balneother. and Climatother. in Med. Sc. and Practice, Bonn, 146, 1953.
93. HERRMANN, R. and HEINKE, E.: Untersuchungen über die Beeinflussung der Haut- Aciditätsverhältnisse durch die Luft - Arch.f.Dermat., 193, 294, 1951.
94. HUDLETT, L. : Untersuchungen über die Abhängigkeit der Erythbildung von klimatischen Faktoren - Fundamenta Radiologica, 5, 15, 1939.
95. JUNGSMANN, H. and HAUS: Kreislaufreaktionen bei Bergbahnfahrten in den Alpen - Med. Klinik, 49, 19, 775-777. Sonderdruck.
96. JUNGSMANN, H. : Ueber die Bedeutung der Kreislaufeigenen Reflexe für die Epidurale Spinalanästhesie und den Anästhesiekollaps - Klin. Wschr., 31, 37/38, 896-900; Sonderdruck.
97. JUNGSMANN, H. : Das Ultraviolettstrahlenangebot und seine therapeutische Ausnutzungsmöglichkeit in den Heilklimaten Deutschlands - Aerztl. Abhlg. über d. Dt. Bäderwesen. Schriftenr. d. Dt. Bäderverb., 9.
98. JUNGSMANN, H. : Der Einfluss des Seeklimas auf die Funktion der Haut - Med. Met. Hefte, 3, 7, 1950.
99. JUNGSMANN, H. : Konstitution und Klima - Schriftenr. d. Dt. Bäderverb., 7-13, 1951.
100. JUNGSMANN, H. : Dermographismus und dermatographische Latenzzeit im Reifungsalter - Z. Kinderheilk., 71, 432, 1952.
101. JUNGSMANN, H. : Kreislaufumstellung im Hochgebirge und an der Nordsee - Schriftenr. d. Dt. Bäderverb., 8, 12, 1952.
102. JUNGSMANN, H. : Kreislaufwirkung von Bergfahrten - Z. Exper. Med., 119, 280, 1952.
103. JUNGSMANN, H. : Die Bedeutung des Reaktionstyps für Heilverfahren in versch. Klimaten - Aerztl. Abhlg. über das Dt. Bäderwesen, 5, 1952.
104. JUNGSMANN, H. : Wirkung spez. Klimate auf Konstitution und Kreislauf - Aerztl. Abhlg. über das Dt. Bäderwesen, 6, 1952.
105. JUNGSMANN, H. : Die Wirkung natürlicher Kurmittel auf den Kreislauf (Seebad - Bergfahrt) - Aerztl. Abhlg. über das Dt. Bäderwesen, 4, 1952.
106. JUNGSMANN, H. and ROHR, H.: Ueber die Form des Femoralispulses und ihre Veränderung unter dynamischer und mechanischer Beeinflussung - Pflügers Arch. Phys., 258, 38-46, 1953.
107. JUNGSMANN, H. and HAUS, E.: Kreislaufreaktionen bei kurzdauerndem Höhenwechsel in den Alpen - Schw. med. Wschr., 38, 1156, 1953.
108. JUNGSMANN, H. : Zur Differentialdiagnose der peripheren Durchblutungsstörungen ; Beitrag zum Problem der angiopathischen Reaktionslage - Klin. Wschr. 31, 149, 1953.
109. JUNGSMANN, H. : Kritisches zur Pulskurvenregistrierung und ihrer Auswertung - Z. experiment. Med. 122, 60-75, 1953.
110. JUNGSMANN, H. : Ueber den Einfluss der vegetativen Struktur des einzelnen Menschen auf den Erfolg klimatischer Behandlungen - Med. Met. Hefte, 8, 15, 1953.
111. JUNGSMANN, H. : Individuelle Klimatherapie. Kreislaufanalysen bei Erfolgen und Misserfolgen einer Kurbehandlung - Dtsch. Med. Wschr., 78, 98 & 134, 1953.
112. JUNGSMANN, H. : Ueber den Einfluss von Tageszeit, Nahrungsaufnahme und Nüchternheit auf den Kreislauf - Zs.f. Kreislaufforschung, 43, 3/4, 120, 1954.
113. JUNGSMANN, H. : Der Kreislaufgesunde und -kranke im Nordseeklima - Aerztl. Fortbildungskursus Bad Nauheim. Nauheimer Fortbildungslehrgänge, 20, 129, 1954.
114. JUNGSMANN, H., HALHUBER, M.J.: Physikalische Kreislaufanalysen im natürlichen Heissluft - Emanatorium des Gasteiner Thermalstollens - Wiener Med. Wschr., 104, 12, 248-251, 1954.
115. JUNGSMANN, H. : Ueber spezifische und unspezifische Wirkungen hydroklimatischer Kuren auf das neurovegetative System - Intern. Kongress f. Thalassotherapie, Balneol. und Klimatol., Opatija/Yougosl., 7, 13, 5, 1954.

116. JUNGSMANN, H. : Ueber die Wirkung des alpinen Höhenklimas auf den gesunden und kranken Menschen - Ztschr. f. angew. Bäder- und Klimaheilkunde, 1, 3, 1954. (Sonderdruck).
117. JUNGSMANN, H. and HAUS, E.: Höhenreaktion des Kreislaufs bei Gesunden und Kranken in den Alpen - Schw. Med. Wschr., 84, 1265, 1954.
118. JUNGSMANN, H. : Experimentelle Kreislaufuntersuchungen zur Medizin-Meteorologie - Ergebnisse der physikalisch-diätischen Therapie, 5, 166, 1955.
119. JUNGSMANN, E. : Kreislaufumstellungen im Nordseeklima - Med. Met. Hefte, 10, 9, 1955.
120. JUNGSMANN, H. : Akklimatisationsprobleme bei Gesunden und Kreislaufkranken - Arch. Physik. Therap., 7, 357-363, 1955.
121. JUNGSMANN, H. and PIRLET, K.: Kreislaufanalytische Befunde bei unterschiedlichen Reaktionsweisen - Aerztl. Forschung, 9, 1, 528-534, 1955.
122. JUNGSMANN, H. : Die Kreislaufanalyse als Mittel zur Beobachtung von Wetter und Klimawirkungen auf den Menschen - Med. Met. Hefte, 11, 1956.
123. JUSATZ, J.H. : Der Einfluss des Wetters und der Sonnentätigkeit auf die Reaktionszeit bei gesunden Menschen - Ärztliche Sachverständigen Zeitung, 49, 5, 21-22, 1943.
124. JUSATZ, J.H. : Ökologie des Menschen als Forschungsaufgabe - Petermanns geographische Mitteilungen, 7/8, 227, 1944.
125. KAMIYAMA, K. : On effect of environment condition on Vasomotor reflex - Papers in Meteorological and Geophysics, 7, 1, 1956.
126. KAMIYAMA, K. : On the variation of type of vasomotor reflex by mountain residence - The Bulletin of Tokyo medical and dental University, 4, 1, 23-29, 1957.
127. KANZ, E. : Experimenteller Beitrag zur kapillar-mikroskopischen Untersuchung der Wetterbeeinflussbarkeit des Organismus - Münch. Med. Wschr., 25, 1263, 1952.
128. KAYSER, P.H. : Hauttemperaturmessungen am Sylter Strand - Veröff. d. Bioklim. Forschungsstelle, 2, 1937.
129. KAYSER, P.H. : Die Einwirkung des Sylter Seeklimas auf die Hauttemperatur - Z. exp. Med., 101, 408, 1937.
130. KEETON, R.W. : The advantages of the physiological viewpoint in Medicine - Illinois Medical J., June 1939.
131. KIEPENHEUER, K.O. : Zur Beeinflussung des menschlichen Blutserums durch die Sonne - Die Naturwissenschaften, 37, 234, 1950.
132. KLAPPICH, H. : Die Oberflächenfeuchte der menschlichen Haut und ihre meteorophysiologischen Beziehungen - Z. exp. Med., 103, 363, 1938.
133. KULLENBERG, B. : Microclimate and Bioclimate (in Swedish) - Svensk Naturvetenskap, 1956.
134. LADELL, W.S.S. : Physiological classification of climate - Int. West African Congress, Ibadan, 1949.
135. LANG, H. : Einfluss seeklimatischer Reize auf die alveoläre Kohlensäurespannung - Diss. med. Kiel, 1939.
136. LANGEN, D. : Experimentelle Studien über die Erythembildung der Sonnen- und Himmelstrahlung - Strahlentherapie, 63, 142, 1938.
137. LEBLOND, C.P. and SUE, P.: Influence of hypophysis on the thyroid uptake and general distribution of radioactive Na I - Anat. Rec., 79, supp. 42, 1941.
138. LEBLOND, C.P., GROSS, J. and LAUGIER, H.: Effect of fasting on resistance to anoxia - J. of Av. Med., 14, 262, 1943.
139. LEBLOND, C.P. and LAUGIER, H.: Effet du jeûne et la réalimentation sur la résistance à l'anoxie - Rev. Can. de Biol., 2, 474, 1943.
140. LEE, D.H.K. : Physiological considerations in the development of tropical centres - FLOR 1.
141. LEE, D.H.K. : Physiological considerations of working conditions in coal mines - FLOM 1.
142. LEE, D.H.K. : Atmospheric conditions in radar room - AVF 42.
143. LEE, D.H.K. : Physiological climatology in America - For section on Physical Geography, Centennial Studies in Geography, 1952.
144. LEE, D.H.K. : Physiological Climatology as a field of study - Ann. Assoc. Am. Geograph., 43, 2, 127-137, 1953.
145. LEE, D.H.K. : Physiological Climatology in America - Physiological Climatology "American Geography-Inventory & Prospect", Chapter 22, 1954.
146. LEHMANN, G. and SZAKALL, A.: Weitere Untersuchungen über den Einfluss der Ultraviolettbestrahlung auf den Arbeitsstoffwechsel und die Arbeitsfähigkeit des Menschen - Arb. Physiol., 6, 84-89, 1932.
147. LEHMANN, G. and SZAKALL, A.: Der Einfluss der Ultraviolettbestrahlung auf den Arbeitsstoffwechsel und die Arbeitsfähigkeit des Menschen - Arb. Physiol., 5, 278-341, 1932; Strahlentherapie 48, 1933.
148. LEHMANN, G. : Die Funktion der menschlichen Nase als Staubfilter - Arb. Physiol. 7, 167-175, 1933.

149. LEHMANN, G. : Die Bedeutung der Filtereigenschaften der Nase für die Entstehung der Staublunge - Münch.Med. Wschr., 1166, 1933.
150. LEHMANN, G. : Die Bedeutung der Nase für die Entstehung der Staublunge - Jahreskurse für ärztl. Fortbildung, 24, 1933; Forschungen u. Fortschritte 10, 268, 1934.
151. LEHMANN, G. : The dust filtering efficiency of the human nose and its significance in the causation of silicosis - The Journal of ind. Hyg., 17, 2, 1935.
152. LEHMANN, G. : Arbeitsfähigkeit und Ultraviolettbestrahlung - Forsch. und Fortschr. 1936, desgl. in der engl. Ausg.
153. LEHMANN, G. : Die Filterung der Atemluft und ihre Bedeutung für Staubkrankheiten - Erg. d. Hygiene, 10, 1-87, 1937.
154. LEHMANN, G. : Die Filterung der Atemluft und ihre Bedeutung für Staubkrankheiten - Verl. Springer, Berlin, 1938.
155. LEHMANN, G. : Versuche über die Wirkung von Sauerstoffatmung bei normalem Druck auf die Leistungsfähigkeit - Luftfahrtmed., 6, 183-200, 1942.
156. LEHMANN, G. : Menschliche Leistung und Leistungsfähigkeit - Forschungen und Fortschritte, 18, 80, 1942.
157. LEHMANN, G. and SZAKALL, A.: Versuche über die Hebung der Leistungsfähigkeit durch Ultraviolettbestrahlung - Arb. physiol., 13, 101-113, 1944.
158. LEHMANN, G. : Ultraviolettbestrahlung und Höhenfestigkeit - Luftfahrtmed., 9, 37-43, 1944.
159. LEHMANN, G. : Adrenalinbestimmung und ihre Anwendung für die Frage der Bioklimatik - Medizin-meteorolog. Hefte, 4, 30-32, 1950.
160. LEHMANN, G. : Ultraviolettbestrahlung als Mittel zur Erhaltung der Arbeitsfähigkeit und zur Gesunderhaltung des arbeitenden Menschen - Zbl.f. Arb. Med. und Arb. Schutz, 1, 1-4, 1951.
161. LEHMANN, G. : Der Einfluss der Ultraviolettbestrahlung auf Arbeitsstoffwechsel und Arbeitsfähigkeit - Strahlentherapie, 83, 109-112, 1950.
162. LEHMANN, G. : Tagesrhythmik und Leistungsbereitschaft - Verhandlungen d. dritten Konferenz d. inter. Ges. f. biol. Rhythmusforschung Acta Med. Scand. 278, 1953.
163. LEHMANN, G. : Die Bedeutung einiger Wellenlängenbereiche für die leistungssteigernde Wirkung der UV-Bestrahlung - Strahlenther., 95, 447-453, 1954.
164. LEHMANN, G. : Die Steigerung der Leistungsfähigkeit durch monochromatische Ultraviolettbestrahlung - Proc. of the First International Photobiological Congress, Amsterdam 1954, - 335. (Ed.H.Veenman & Zn. Wageningen, Neth., 1956.)
165. LEHMANN, G. and DIECKMANN, D.: Die Wirkung mechanischer Schwingungen (0,5 bis 100 Hertz) auf den Menschen - Forschungsber. des Ministeriums für Wirtschaft und Verkehr des Landes Nordrhein - Westfalen, 362; Westd. Verl. Köln und Opladen, 1957.
166. LEISTNER, W. : Die hygienische und bioklimatische Bedeutung des Dampfdruckes in Innenräumen und die Behaglichkeits- und Schwülegränze - Med.- Meteor. Hefte, 6, 18-31, 1951.
167. LEISTNER, W. : Der Verlauf individueller Leukozytenkurven in Abhängigkeit vom Witterungsgeschehen - Medizin-Meteorologische Hefte, 8, 1953.
168. LEXOW, K.G. : Einfluss der Luftfeuchte und Jahreszeit auf die Perspiratio insensibilis nocturna - Diss.med. Kiel, 1957.
169. LOTZ, R.G.A. : Strahlung und Stoffwechsel. II. Die Wirksamkeit der Ultra-Violett - und Ultrarotstrahlung einschliesslich des sichtbaren Lichtes - Zeitschrift für Naturforschung, 3b, 349, 1948.
170. LOTZ, R.G.A. : Ein Gerät zur Umsatzbestimmung. Zugleich ein Beitrag zur Wärmeleitfähigkeitsmessung von Expirationsluft - Zeitschrift f.Naturforschung, 3b, 357, 1948.
171. LOTZ, R.G.A. : Der Einfluss der Strahlung auf den Gasstoffwechsel - Verhandlungen d. Deutschen Zoologischen Ges. 1948, 468, 1949.
172. LOTZ, R.G.A. : Strahlung und Stoffwechsel. III. Das "Strahlungsklima" zur Steuerung des Gasstoffwechsels von Säugetieren - Ztschr. f.Physiol.Bed., 32, 228, 1950.
173. LOTZ, R.G.A. : Neue Forschungen über die Wirkung einer kombinierten Ultra-Violett- und Infrarotbestrahlung - Archiv.Phys. Ther., 3, 357, 1951.
174. LOTZ, R.G.A. : Der Einfluss einer kombinierten UV- und UR-Strahlung auf den Gasstoffwechsel bei Warmblütern - Strahlenther., 86, 149, 1951.
175. LOTZ, R.G.A., AMELUNG, W. and SCHMIDT, L.: Kreislaufanalytische Untersuchungen zur Diagnostik und Therapie vegetativer Störungen - Deutsch.Archiv.Klin. Mediz., 201, 568, 1954.
176. LOTZ, R.G.A., AMELUNG, W. and SCHMIDT, L.: Experimentelle Untersuchungen und kritische Betrachtungen zur Frage der vegetativen Labilität - Arch. phys. Ther., 7, 221, 1955.

177. MCDONAGH, J.E.R. : The influence of climate on the protein in the blood - An Exhibit before the First International Congress for Bioclimatology in Paris, August 1956.
178. MILIN, R., CIGLAR, M. and JOVANOVIĆ, V.: Influence du climat maritime sur les glandes surrénales du lapin - Int. Congress of Hydroclim. and Thalassother., Opatija, 1, 83-90, 1954.
179. MILIN, R., CIGLAR, M. and JOVANOVIĆ, V.: Influence du climat maritime sur la glande thyroïde du lapin - Int. Congr. of Hydroclim. and Thalassother., Opatija, 1, 109-114, 1954.
180. MILIN, R. and CIGLAR, M.: Influence de l'insolation sur la glande thyroïde du lapin - Comptes Rendus de l'Assoc. des Anatomistes, 42nd meeting, 1035-1038, Paris, 25-30 July 1955.
181. MILIN, R. : Contribution à l'étude de l'influence de la lumière et de l'obscurité sur l'hypothalamus - Pathophysiology Diencephalica, Int. Symp., 159-164, Milan, May 1956.
182. MILIN, R. and CIGLAR, M.: Die Wirkung der Dunkelheit auf den Hypophysenhinterlappen - Verhandlungen d. Anatomischen Ges. auf d. 53. Versammlung, 189-195, Stockholm, 22-25 August, 1956.
183. MILIN, R. : Die Wirkung des Lichtes auf den Hypophysenhinterlappen der Ratte - Verhandlungen d. Anatomischen Ges. auf d. 54. Versammlung, 191-198, Freiburg/Br., 22-25 September 1957.
184. MONGE, C. : Biological Basis of human behaviour - Int. Symp. and Anthropol. Sciences, Wenner Gren Found. New York, in Anthropology Today - The Univ. Chicago Press, 127-144, 1953; Man, climate and changes of altitude - Meteorological Monographs, Oct. 1954.
185. MOURIQUAND, G. and CHARPENTIER, R.: Météoropathologie et dystrophies infantiles. Coup de chaleur et choc hyponutritique - La Presse Médicale, 36, 102, 1625-1629, 1928.
186. MUECHER, H., GRUENEWALD, G. and DISTLER, W.H.: Psychische und physiologische Wirkungen des Wetters - Arzneimittelforschung Beiheft 6, 1956. (Publ. Editio Cantor Aulendorf i. Württ.).
187. NOLL, P. : Beobachtungen über Hauttemperaturen bei verschiedenen Wind- und Sonnenexpositionen - Diss.med. Kiel, 1951.
188. PAYNE, R.B. : A preliminary study of the motivational properties of cold stress - Arctic Aeromed. Lab., Ladd AFB, April 1959.
189. PENNDORF-PENDL, E. : Versuche über die Wirkung negativ ionisierter Luft auf die Blutbakterizidie des Kaninchens - Bioklim. Beibl., 10, 2/3, 86-88, 1943.
190. OTTO, W. : Ueber Biorhythmen bei Neugeborenen - Zschr. f. Endokrinologie, 32, 3/4, 1955.
191. OTTO, W. and NOACK, H.: Jahreszeitliche Schwankung der Körpermasse bei 100.000 neugeborenen Berliner und Leipziger Kindern - Lecture given for the "Sektion Geburtshilfe und Säuglingsfürsorge" der Akademie der Wissenschaften zu Berlin am 30.11.1956 - Z. menschl. Vererb.- und Konstitutionslehre, 34, 250-260, 1957.
192. OTTO, W. and NOACK, H.: Statistische Untersuchungen zur Frage jahreszeitlicher Einflüsse in der Geburtshilfe unter besonderer Berücksichtigung der Neugeborenenlänge - Archiv. f. Gynäkologie, 188, 457-473, 1957.
193. PFLEIDERER, H. : Bedeutung der Haut im Lichte heilklimatischer Forschung - Z. wiss. Bäderkunde, 3, 1928.
194. PFLEIDERER, H. : Das Grundgesetz des Energiwechsels in der Biologie - Bemerkg. zu der gleichnamigen Arbeit von H. Bohnenkamp, 1931, Kon.Wschr., 896, 1745, 1932.
195. PFLEIDERER, H. : Die Abkühlungsgrösse, ihre heilklimatische Bedeutung und ihre Meszmethode - Strahlentherapie, 40, 562, 1931.
196. PFLEIDERER, H. and LESS, L.: Die klimatischen Ansprüche und die Atmungswege des menschlichen Körpers - Bioklim. Beibl., 1, 1935.
197. PFLEIDERER, H. : Bioklimatische Forschungsstätten an der Nordsee - Med. Klin. 24/25, 1936.
198. PFLEIDERER, H. : Die bio-klimatische Bedeutung der Strahlen - "Die Welt der Strahlen" herausgegeben von H. Woltereck, Quelle u. Meyer, 1937.
199. PFLEIDERER, H. : Kontaktwärme und Wärmestrahlung in "Erforschung und Praxis d. Wärmebehandlung der Medizin einschl. Diathermie und Kurzwellentherapie" - 2. Frankfurter Konferenz f. med.-naturwiss. Zusammenarbeit, Mai 1937, herausgegeben von Rajewski u. Lampert, Steinkopf, 1937.
200. PFLEIDERER, H. and WANKE, R.: Messung und Beurteilung von Hauttemperatur-Asymmetrien bei Gesunden und Hirnverletzten - Der Nervenarzt, 12, 553, 1939.
201. PFLEIDERER, H. : Cellophanteppchen für die Prüfung der Lichtempfindlichkeit der Haut - Strahlentherapie, 90, 572-578, 1953.

202. PFLEIDERER, H. : Zur Methodik der Erythem- und Pigmentmessung - Proc. of the First International Photobiological Congress, Amsterdam, 248-250, 1954.
203. PFLEIDERER, H. : Untersuchungen über eine neue U.V.-Messmethode. Vortrag 4. Hanauer UV-Tagung, Jan. 1956 - Strahlentherapie, 4, 101, 1956.
204. PIERY and MILHAUD : Action du climat marin sur le peau et les fonctions cutanées - Imp. D. van Keerberghen & Fils, Rue Piers 101, Bruxelles.
205. PIERY, CLUZET, PONTUS and MILHAUD: Effets de la raréfaction atmosphérique sur le coeur - Archiv. d'électricité Médicale, mars-avril 1935.
206. PIERY, M. and MILHAUD, M.: Glandes endocrines et stations climatiques - Presse thermique et climatique, 15 mai 1936.
207. PIERY, M. and MILHAUD, M.: Climats et affections de l'appareil urinaire - Gaz.Méd.de France, 46, 1, 1939.
208. PIERY, M. and MILHAUD, M.: Climat marin en peau - Jl. de Médecine de Lyon, 5 septembre 1941.
209. PIERY, M. and MILHAUD, M.: Le vent: son importance en climatologie médicale - Gz. Méd. de France, 53, 14, 401-404, 1946.
210. REITER, R. : Steuerung von Geräten durch Körperpotentiale - Grenzgebiete der Medizin (Urban und Schwarzenberg), 1, 1, 1948.
211. REITER, R. : Elektrische Feldänderungen als Vermittler meteorotroper und solarer Wirkungen (unter Berücksichtigung der Membranpermeabilität und des Verschiebungsstromes im biologischen Objekt - Grenzgebiete der Medizin (Urban und Schwarzenberg), 1, 233, 1948.
212. REITER, R. : Isotope Elemente in der Medizin - Grenzgebiete der Medizin (Urban und Schwarzenberg), 2, 286, 1949.
213. REITER, R. and HIRSCHBERGER, M.: Wettervorgänge und Sonneneruptionen beeinflussen den elektrischen Hautwiderstand - Deutsche Medizinische Wochenschrift, 78, 47, 1640, 1953.
214. REITER, R. : Schmerzempfindung und Wetter (Ein Nachtrag zur Arbeit von Brunnhölzl, Schedel und Ungeheuer in ders. Zeitschrift Serie B, 4, 317, 1953. - Archiv f. Meteorologie, Geophysik und Bioklimatologie, Wien, Serie B, 5, 1, 112, 1953.
215. REITER, R. : Die Bedeutung extrem langwelliger elektromagnetischer Strahlungen in der Bioklimatologie - Strahlentherapie (Urban und Schwarzenberg), 4, 89, 628, 1953.
216. RIEMERSCHMIDT, G. : Ultraviolettreaktion der Haut und langwellige Lichtstrahlung - Veröffentlich. Bioklim. Forschungsstelle, 1, 1937.
217. RIEMERSCHMIDT, G. : Zur Erythemwirkung der natürlichen Ultraviolettstrahlung - Fundamenta Radiologica, 2, 126, 1938.
218. RIEMERSCHMIDT, G. and SCHEUER, O.: Die Wirkung von Teilbädern auf die Hauttemperatur - Z. Exp. Med., 1939.
219. RIEMERSCHMIDT, G. : Ueber die Vergleichbarkeit von Messungen mit dem UV-Dosimeter - Fundamenta Radiologica, 1940.
220. RING, G.C. : Metabolism and body temperature of normal and adrenalectomized rats during exposure to cold - Am.J. Physiol., 122, 435-445, 1938.
221. RISSE : Einflüsse von Wetter und Jahreszeit auf die Magensaftsekretion des Menschen - Diss. Köln, 1939.
222. SAPIN- JALOUSTRE, J.: Le problème de la vision dans le blizzard - Annales d'Optique oculaire, 60, 2, 36-53, 1953.
223. SARGENT, F. and IPPOLITO, T.: Phases of physiological response to meteorological stimuli - 5th annual report of Bowdoin Scientific Station Bull, 7, 33, 1941.
224. SARGENT, F. : The meteorological environment (pp 13-15); Polar front and blood pH (p. 46); Note on the cold front and interdiurnal metabolic gradients (p. 137); Perception threshold (pp. 446-447) in W.F. Petersen: Man-Weather-Sun, C. Thomas, Springfield, Ill. USA, 1947.
225. SARGENT, F. : Some fundamental principles of metabolism - MD Magazine, 4, 296, 1949. Book review.
226. SARGENT, F. : Seasonal Variation in Metabolism. A critique of homeostasis - Arch. Meteor. Geophys. u. Bioklim., Serie B, 3, 289, 1951.
227. SARGENT, F. and CONZOLAZIO, C.F.: Stress and Ketone Body metabolism - Science, 113, 631. 1951.
228. SARGENT, F. : Climatic stress: A bioclimatological principle - Bull.Am.Met.Soc.32, 271, 1951. (Abstract).
229. SARGENT, F. and GOLDEN, R.: The passage of ascorbic acid across the membrane of the human erythrocyte - Arch. Biochem. Biophys., 39, 138, 1952.
230. SARGENT, F., MELTZER, E. and ANDERSEN, K.L.: Erythrocyte sedimentation rate increased by dietary stress - Fed. Proc. 13, 99, 1954.
231. SARGENT, F., JOHNSON, E., SARGENT, V.W. and EVANS, R.D.: Interrelations in man among nitrogen balance, Caloric intake and protein intake (abstract) - Fed.Ass. 13, 462, 1954.

32. SARGENT, F. : Season and the metabolism of fat and carbohydrate: A study of vestigial physiology (In recent studies in bioclimatology) - Meteorol. Monog., 2, 8, 68, 1954.
33. SARGENT, F. and JOHNSON, R.E.: The effects of diet on renal function in healthy men - The American Jnl. of clin. nutrition, 4, 5, 467-481, 1956.
34. SARGENT, F. and SLUTSKY, H.L.: The natural history of the eccrine miliarias: a study in Human Ecology - New England Jnl. of Medicine, 256, 401-408 and 451-459, 1957.
35. SCHIELDRUP PAULSEN, H.: Klimatische innvirkninger på menneskets organisme - Naturen, 15, 1955.
36. SCHULZE, R. : Die biologische Wirkungen der optischen Strahlung - Nat. Med. D 21, Biophysik I, 114.
37. SCHULZE, R. : Zur Wirkung des Sonnen-UV-Strahlens auf die menschl. Haut - Naturw., 33, 373, 1946.
38. SCHULZE, R. : Die biologisch wirksamen Komponenten des Strahlungsklimas - Naturw., 34, 238, 1947.
39. SCHULZE, R. : Ueber die Wirkung der Sonnenstrahlung auf die menschl. Haut - Ann. Met., 1, 12, 1948.
40. SCHULZE, R. : Gewöhnung und Umstimmung der menschl. Haut bei Bestrahlung mit Sonne und Ultra-Vitaluxlampe - Strahlenther., 86, 51, 1951.
41. SCHULZE, R. : Ueber die "sekundäre pigmentierung" der menschlichen Haut als Folgeerscheinung des Zelltodes nach UV-Bestrahlung - Proc. of the international congress of Physical Medicine, 1952.
42. SEITZ, K. : Untersuchungen über Atmungsperspiration und Expirationstemperatur unter kühlen Bedingungen - Diss. med. Kiel, 1949.
43. SELYE, H. and SCHENKER, V.: Adrenal cortical assay method under cold conditions - Proc. Soc. Exper. Biol. & Med., 39, 518-522, 1938.
44. TAKATA, M. : Solare Einflüsse auf den Ausfall der Flockungszahlreaktion - Archiv für physikalische Therapie, 7, 3, 1955.
45. TALBOTT, J.H. and MULLER, G.L.: The effect of high altitudes on the Cholesterol, Lecithin and Fatty Acids in the Plasma of healthy men - Arch. Int. Med., 47, 855, 1931.
46. TALBOTT, J.H. and DILL, D.B.: Clinical observations at high altitude - Am. J. Med. Sc., 192, 626, 1936.
47. TALBOTT, J.H. : Studies at high altitude. II. - Folia Haematologica, 55, 23, 1936.
48. TOSHIO TORII : 24-hour periodicity of adrenal cortex - Endocrinologia japonica (in English), 2, 79-86, 1955.
49. UNGEHEUER, H., SCHEDEL, F. and BRUNNHOLZL, K.: Wetter und Durchblutung - Münchener Medizinische Wschr., 94, 1952.
50. VERZAR, F. and VÖGTLI, W.: Die Sauerstoffsättigung des arteriellen Blutes in mittleren Höhen und ihre Bedeutung als Klimafaktor - Physiologisches Inst. der Univ. Basel. Benno Schwabe & Co., Basel, 1945.
51. VERZAR, F. and VÖGTLI, W.: Die Sauerstoffsättigung des Kapillarblutes im Höhenklima - Aus dem Physiologischen Inst. der Univ. Basel. Benno Schwabe & Co., 1945.
52. VERZAR, F. and VÖGTLI, W.: Die initialen Erythrozyten-Veränderungen im Hochgebirge - Aus dem Physiologischen Inst. der Univ. Basel, Benno Schwabe & Co., 1945.
53. VERZAR, F. : Die ersten morphologischen Erythrozyten-Veränderungen im Hochgebirge - Physiologisches Inst. der Univ. Basel, Benno Schwabe & Co., 41-49, 1945.
54. VERZAR, F. : Der Gehalt des Blutplasmas an Eiweiss, Rest-Stickstoff, Chlor und Wasser im Höhenklima (3450 m.ü.M.) - Physiologisches Institut der Universität Basel, Benno Schwabe & Co., 50-52, 1945.
55. VERZAR, F., SOMOGYI, J.C. and WIRZ, H.: Veränderung von Gesamtblut- und Plasmamenge im Hochgebirge - Physiologisches Inst. der Univ. Basel und dem Hochalpinen Forschungsinst. auf Jungfrau-Joch. Benno Schwabe & Co., 53-59, 1945.
56. VERZAR, F., MASON, M. and FITZGERALD, O.: Die Aenderung des Serumbilirubins und der Erythrozytenanzahl im Hochgebirge - Physiologisches Inst. der Univ. Basel und das Hochalpinen Forschungsinst. auf Jungfrau-Joch. Benno Schwabe & Co., 61-68, 1945.
57. VERZAR, F., VÖGTLI, W. and DOETSCH, R.: Ueber die angebliche Erythrozytenzunahme in mittleren Höhen - Physiologisches Institut der Univ. Basel. Benno Schwabe & Co., 69-75, 1945.
58. VERZAR, F. and VÖGTLI, W.: Veränderungen der Atmung in grossen Höhen und bei künstlichem Sauerstoffmangel. I. - Physiologisches Institut der Univ. Basel, Benno Schwabe & Co., 81-91, 1945.
59. VERZAR, F. : Atmungsveränderungen in grossen Höhen und bei künstlichem Sauerstoffmangel. II. - Physiologisches Inst. der Univ. Basel, Benno Schwabe & Co., 92-99, 1945.

260. VERZAR, F. and VÖGTLI, W.: Der Sauerstoffdruck im Gewebe beim Aufstieg in grosse Höhen nach Untersuchungen an Ratten - Höhenklima-Forschungen des Basler physiologischen Inst. II, 1948.
261. VERZAR, F. and VÖGTLI, W.: Die O_2 und CO_2 -Spannung im Gewebe beim Aufstieg von Menschen in grosse Höhen - Höhenklima-Forschungen des Basler Physiolog. Inst. II, 1948.
262. VERZAR, F. and VÖGTLI, W.: Die Aenderung des Sauerstoffdrucks im Harn beim Aufstieg in grosse Höhen als Mass des Sauerstoffdrucks im Gewebe - Höhenklima-Forschungen des Basler Physiologischen Inst. II, 1948.
263. VERZAR, F. : Bemerkungen über die Grösse innersekretorischer Organe beim Aufenthalt in grossen Höhen - Höhenklima-Forschungen des Basler Physiologischen Inst. II, 1948.
264. VERZAR, F. : Störungen der Wärmeregulation in grossen Höhen, untersucht mittelst Novocainwirkung - Höhenklima-Forschungen des Basler Physiologischen Inst. II, 1948.
265. VERZAR, F. : Hauttemperaturmessungen im Tiefland und auf Jungfraujoch (3450 m ü.M.) - Höhenklima-Forschungen des Basler Physiologischen Inst. II, 1948.
266. VERZAR, F. and VÖGTLI, W.: Die Urobilingen-Ausscheidung in grossen Höhen als Mass des Blutabbaues - Höhenklima-Forschungen des Basler Physiologischen Inst. II, 1948.
267. VERZAR, F., BECK, I. and SASS-KORTSAK, A.: Resistenzänderung der Erythrocyten bei der Abnahme ihrer Zahl beim Abstieg aus grossen Höhen - Höhenklima-Forschungen des Basler Physiologischen Inst. II., 1948.
268. VERZAR, F., and WANG, S.J.: Erythropoese in der Höhe von 1880 m bei normalen und thyreoidektomierten Kaninchen - Schw. med. Wschr., 78, 713, 1949.
269. VERZAR, F. and SAILER, E.: Schilddrüsenfunktion bei vermindertem atmosphärischem Druck - Verhandlungen des Schweizerischen Vereins der Physiologen und Pharmakologen. 37.Tagung in Zürich am 4. November 1950 - Helv. Physiol. Acta, 8, C74, 1950.
270. VERZAR, F. and SAILER, E.: Die Ausscheidung von Corticosteroiden in Ruhe, bei Muskelarbeit und in der Höhe - Verhandlungen des Schw. Vereins der Physiologen und Pharmakologen. 37.Tagung in Zürich am 4. November 1950., Helv. Physiol. Acta, 8, C72, 1950.
271. VERZAR, F., PEYSER, E. and SASS-KORTSAK, A.: Influence of O_2 -content of inspired air on total lung volume - The Am. Jnl. of Physiol., 163, 1, 111, 1950.
272. VERZAR, F., BUEHLMAN, A. AND WANG, S.I.: Die Erythrocytenzahl in mittleren Höhen - Schweiz. Mediz. Wschr., 81, 4, 80, 1951.
273. VERZAR, F., WANG, S.J. and WIRZ, H.: Die O_2 -Sättigung des arteriellen Blutes bei Mensch und Kaninchen auf 1800 m ü.M. und ihr Zusammenhang mit der Erythrocytenzunahme - Schweiz. med. Wschr., 81, 82, 1951.
274. VERZAR, F. : Die "dritte Form" der Atmungsregulation beim Aufstieg in grosse Höhen und bei der Hochgebirgsbevölkerung der Anden - Bull. der Schweiz. Akademie der medizinischen Wissenschaften, 7, 3/4, 201-211, 1951.
275. VERZAR, F. and FLUECKIGER, E.: Senkung und Restitution der Körpertemperatur bei niedrigem atmosphärischem Druck und der Einfluss von Thyreoidea, Hypophyse und Nebennierenrinde auf dieselbe - Helvetica, Physiologica et Pharmacologica Acta, 10, 3, 349, 1952.
276. VERZAR, F., SAILER, E. and VIDOVIC, V.: Changes in thyroid activity at low atmospheric pressures and at high altitudes, as tested with ^{131}I - The Jnl. of endocrinology, 8, 308, 1952.
277. VERZAR, F. : Die Zahl der Lymphocyten und eosinophilen Leukocyten in 1800 und 3450 m Höhe - Schweiz. Med. Wschr., 82, 13, 324, 1952.
278. VERZAR, F. and FLUECKIGER, E.: Einfluss von Thyreoidea, Nebennierenrinde und Hypophyse auf Senkung und Restitution der Körpertemperatur bei niedrigem atmosphärischem Druck - Helv. Physiol. Acta, 10, C15, 1952.
279. VERZAR, F., KEITH, J. and PARCHET, V.: Temperatur und Feuchtigkeit der Luft in den Atemwegen - Pflügers Archiv, 257, 400-416, 1953.
280. WHITEN, A.J. : Weather and nerves - Weather, 4, 8, 1949.
281. WHITTOW, G.C. and GLASER, E.M.: Evidence for a nonspecific mechanism of habituation - Jnl. Physiol., 122, 1953.
282. WILDFUEHR, G. and MUECKE: Ueber Generationsdauerexperimente mit verschiedenen Bakterienarten unter bestimmten meteorologischen Bedingungen - Arch.f.Hyg., 134, 1, 1, 1951.
283. WILDFUEHR, G. : Ueber die Einwirkung des biotropen Frontenfaktors auf die allgemeine Resistenz - Zschr. f.Immf., 108, 318, 1951.
284. WILDFUEHR, G. : Ueber die Einwirkung von Wetterfronten auf Leukozyten und Blutplättchen - Zschr. f.d. ges. Inn.Med.u.i. Grenzgebiete, 11/12, 342, 1951.

5. WINSOR, T. and BECKETT, J.C.: Biologic effects of ionized air in man - Am.J.Phys.Med., 37, 2, 83-89, 1958.
6. WORDEN, J.L. : The effect of air ion concentration and polarity on the carbon dioxide capacity of mammalian blood plasma - Presented before the Am. Physiol. Society at the 38th Meeting of the Fed.of American Soc. f. Experimental Biol., Atlantic City, 12 April 1954.
7. WORDEN, J.L. : The effect of unipolar ionized air on the Hydrogen ion concentration of Mammalian Blood - Presented before the Division of Biol.Chem. of the Am. Chemical Soc., 125th Meeting, Kansas City, Missouri, 25 March 1954.
8. ZITOWITZ, L. and HARDY, J.D.: Influence of Cold Exposure on thermal burns in the rat - J. Applied Physiol., 12, 2, 147-154, 1958.

ACCLIMATISATION BIOCLIMATOLOGY

- . BEHMANN, F.W. : Wärmebilanz bei intravasaler Kühlung unter besonderer Berücksichtigung des Kältezitterns - Ber. ges. Physiol. u. Pharmakol., 180, 140, 1956.
- . BEHMANN, F.W. : Die Regulierung der Wärmeproduktion bei Auskühlung homoiothermer Tiere - Arch. exp. Path.u. Pharmakol., 228, 126, 1956.
- . BEHMANN, F.W. : Wärmebilanzen bei künstlicher Hypothermie. Ein Beitrag zum Problem der Oekonomie des Kältezitterns - Pflügers Archiv, 263, 166-187, 1956.
- . BEHMANN, F.W. : Ein Modell zur Temperaturregulation des bekleideten und unbekleideten Menschen - 20 Internat. Physiol. Congr., Brussel, 77, 1956.
- . BEHMANN, F.W. and BONTKE, E.: Intravasale Kühlung. Eine Methode zur Erzielung steuerbarer Hypothermie - Pflügers Archiv, 263, 145-165, 1956.
- . BEHMANN, F.W. and THAUER, R.: Die Druck-Stromstärke-Beziehung des Gesamtkreislaufes bei künstlicher Hypothermie - Pflügers Archiv, 263, 145, 1956.
- . BEHMANN, F.W. : Die Bedeutung der kapillaren Gewebeeigenschaften für das Wärmeschutzvermögen einer Bekleidung - Melliand Textilberichte, 38, 12, 1957.
- . BEHMANN, F.W. : Bestimmung des Energieumsatzes am lebenden Organismus - ATM V 663-2, 1957.
- . BEHMANN, F.W., HEERD, E. and HEINRICHS, H.J.: Das thermische Schwitzen beim Menschen in Abhängigkeit von Haut- und Rectaltemperatur - Pflügers Archiv, 266, 1, 1957.
- . BEHMANN, F.W. : Einfluss der Hydrophobierung auf das bekleidungsphysiologische Verhalten von Textilien - Melliand Textilber., 39, 786, 1958.
- . BEHMANN, F.W. and BONTKE, E.: Die Regelung der Wärmebildung bei künstlicher Hypothermie I. Experimentelle Untersuchungen über den Einfluss der Narkosetiefe - Pflügers Archiv, 266, 408-421, 1958.
- . BEHMANN, F.W. : Die Regelung der Wärmebildung bei künstlicher Hypothermie. II. Theoretische Untersuchungen - Pflügers Archiv, 266, 422-446, 1958.
- . BERG, H. : Zur Bedeutung von Temperaturen und Feuchtigkeit für die Behaglichkeit - Wetter und Leben, 9, 3/5, 47-50, 1957.
- . BREDEMEYER, G. : Untersuchungen über den Einfluss künstlicher Strahler auf den Wärmehaushalt des Menschen bei Freiluftliegekuren in der kalten Jahreszeit - Diss. med. Kiel, 1953.
- . BUETTNER, K. : Physikalisches zum Wärmehaushalt des Menschen insbes. über die Wärmeabgabe durch Leitung - Klin. Wochenschr. 11, 1508-1509, 1932.
- . BUETTNER, K. : Kritisches über Abkühlungs- und Austrocknungsgrösse - Meteorol.Zeit. 4, 126-130, 1933.
- . BUETTNER, K. : Wärme- und Strahlungshaushalt des Menschen und des Frigorimeters - Strahlenther., 48, 386, 1933.
- . BUETTNER, K. and PFLEIDERER, H.: Die physiologischen und physikalischen Grundlagen der Hautthermometrie - J.A. Barth, Leipzig, 1935.
- . BUETTNER, K. : Bedeutung und Messung der Pberflächenfeuchte für die Transpirationsanalyse - Biolog. Zentralbl., 55, 356, 1935.
- . BUETTNER, K. : Ueber den Einfluss der Blutzirkulation auf die Wärmeverfrachtung in der Haut - Strahlentherapie, 55, 333, 1936.
- . BUETTNER, K. : Feuchte und Verdunstung der menschlichen Haut - Forschungen u.Fortschritte, 12, 151, 1936.
- . BUETTNER, K. and PFLEIDERER, H.: Methodik der thermoelektrischen Hauttemperaturmessung - Abderhaldens Handbuch der biologischen Arbeitsmethoden, 4, 13, 1937.
- . BUETTNER, K. : Ueber die Wärmestrahlung und die Reflexeigenschaften der menschlichen Haut - Strahlenther.. 58, 345, 1937.

24. BUETTNER, K. : Wärmehaushalt des Menschen und Wärmeschutz im Flugzeug - Lilienthalges. vom 20.6.1940. Jahrb.d. Luftfahrtforschung, 1941.
25. BUETTNER, K. : Wärmehaushalt des Menschen und Raumklima im Flugzeug - Luftwissen, 11, 1944.
26. BUETTNER, K. : Neue physikalische Ergebnisse zum Wärme- und Wasserhaushalt des Menschen - Wiener Med. Wschr., 94, 373-374, 1944.
27. BUETTNER, K., FRANK, A. : Das anfrieren der Haut an kalten Metallen - Der Militärarzt, 1, 9, 1944.
28. BUETTNER, K. : Heat and Cold in aircraft - German Aviation Medicine, Dept. of the Air Force, Washington, 8, 757-765, 1950.
29. BUETTNER, K. : Physical heat balance in man - German Aviation Medicine, Dept. of the Air Force, Washington, 8C, 766-791, 1950.
30. BUETTNER, K. : Protective clothing for heat and cold - German Aviation Medicine, Department of the Air Force, Washington, 8H, 876-886, 1950.
31. BUETTNER, K. : Conflagration Heat - German Aviation Medicine, Dept. of the Air Force Washington, 13A, 1167-1187, 1950.
32. BUETTNER, K. : Effects of extreme heat on man - J.Am.Med.Ass., 144, 732-738, 1950.
33. BUETTNER, K. : Bioclimatology of space flight - Space Medicine, Univ. of Illinois Publ., 1951.
34. BUETTNER, K. : Effects of extreme heat and cold on human skin. 1. Analysis of temperature changes caused by different kinds of heat application - J. Appl. Physiol., 3, 691, 1951.
35. BUETTNER, K. : Effects of extreme heat on human skin. II. Surface temperature pain and heat conductivity in experiments with radiant heat - Jnl. Appl. Physiol., 3, 703, 1951.
36. BUETTNER, K. : Effects of extreme heat and cold on human skin. III. Numerical Analysis and pilot experiments on penetrating flash radiation effects - J. Appl. Physiol., 5, 207-270, 1952.
37. BUETTNER, K. : Thermal aspects of travel in the aeropause-problems of thermal radiation - Physics and Medicine of the Upper Atmosphere, Un. of New Mexico Press, Albuquerque, New Mexico, 1952.
38. BUETTNER, K. : Die menschliche Haut unter der Einwirkung der Hitzestrahlung - Umschau, 23, 1952.
39. BUETTNER, K. : Diffusion von Wasser und Wasserdampf durch die menschliche Haut - Naturwissensch., 40, 392, 1953.
40. BUETTNER, K. : Diffusion of water and watervapour through human skin - Journ. Appl. Physiol., 6, 229-242, 1953.
41. BUETTNER, K. : Thermal stresses in modern aircraft - Arch.Met.Geophys. and Bioklim., 5 B, 377-387, 1954.
42. BUETTNER, K. : Thermal comfort as a climatological criterion - Met. Monographs, 1954.
43. BUETTNER, K. : Water gain and loss through human skin - The Am. J. of Physiology, 3, 1955.
44. BUETTNER, K. : Skin water intake in normal persons and during menstrual cycle, pregnancy and cardiac adema, and concentration of sweat - Fed. Proc., 15, 1, 1956.
45. BURCH, G.E. and WINSOR, T. : Rate of insensible perspiration (Diffusion of water) locally through living and through dead human skin - Archives of Internal Medicine, 74, 437-444, 1944.
46. BURCH, G.E. : Study of water and heat loss from the respiratory tract of man - Archives of Internal Medicine, 76, 308-314, 1945.
47. BURCH, G.E. : Rate of water and heat loss from the respiratory tract of normal subjects in a subtropical climate - Archives of Internal Medicine, 76, 315-327, 1945.
48. BURCH, G.E. : The rate of water loss from the respiratory tract of man living in a subtropical climate - Science, 102, 619-620, 1945.
49. BURCH, G.E. : The influence of environmental temperature and relative humidity on the rate of water loss through the subtropical climate - Am. Journal of Med. Science, 211, 181-188, 1946.
50. BURCH, G.E. : The rates of water and heat loss from the respiratory tract of patients with congestive heart failure who were from a subtropical climate and resting in a comfortable atmosphere - Am. Heart Journal, 32, 88-99, 1946.
51. BURCH, G.E. : Environmental conditions which initiate sweating in resting men - Proc. of the Soc. for exp. Biology and Medicine, 521-523, 1948.
52. BURCH, G.E. and BERENSON, G.S. : Studies of diffusion of water through dead human skin: The effect of different environmental states and of chemical alterations of the epidermis - The Am. Journal of Tropical Medicine, 31, 842-853, 1951.

53. BURCH, G.E. and BERENSON, G.S.: A study of the sodium, potassium and chloride contents of thermal sweat of man collected from small isolated areas of skin - The Journal of Laboratory and clin. Med., 42, 58-77, 1953.
54. CARLSON, L.D. and BURNS, H.L.: A method suitable for application of moist heat therapy and applicable for controlling moist chambers (abstract) - Fed.Proc. 6, 86, 1947.
55. CARLSON, L.D., BURNS, H.L., QUINTON, W.E. and BARK, R.S.: Report of studies on acclimatization winter 1948-1949 - USAF, Techn.Rep. No. 5835, 1949.
56. CARLSON, L.D. : Research in environmental physiology at Ladd Field, Alaska. Minutes Subcommittee on shelter and clothing, 89-91, Nat. Res. Coun., 21 February 1950.
57. CARLSON, L.D., YOUNG, A.C. and BURNS, H.L.: Regional heat loss by temperature gradient calorimetry (abstract) - Fed.Proc., 9, 140, 1950.
58. CARLSON, L.D. : Acclimatization to cold. Physiological mechanisms. Proceedings of the Alaskan Science Conference Nov. 1950 (abstract) - Bull.nat.Res.Coun. Washington, 122, 89, 1951.
59. CARLSON, L.D., YOUNG, A.C., BURNS, H.L. and QUINTON, W.F.: Acclimatization to cold environment. Physiologic mechanisms - USAF Tech. Rep. no. 6247, 1951.
60. CARLSON, L.D., YOUNG, A.C., BURNS, H.L. and QUINTON, W.F.: Temperature gradient calorimetry - USAF Tech.Rep. No. 6248, 1951.
61. CARLSON, L.D., BURNS, H.L., YOUNG, A.C. and HOLMES, T.H.: Adaptive mechanisms in cold environments (abstract) - Fed. Proc. 11, 22, 1952.
62. CARLSON, L.D. and YOUNG, A.C.: Thermal gradient calorimeter - Arctic Aeromed. Lab.Proj.Rep. 22, 1301-1302, 1952.
63. CARLSON, L.D., BURNS, H.L., HOLMES, T.H. and WEBB, P.P.: Adaptive changes during exposure to cold - Arctic Aeromed.Lab.Proj.Rep., 22, 1301-1302 and J.appl.Physiol., 5, 672-676, 1953.
64. CARLSON, L.D. and COTTLE, W.: Adaptation of caloric balance in rat exposed to cold (abstr.) - Fed. Proc., 12, 23, 1953.
65. CARLSON, L.D. and COTTLE, W.: Adaptive changes in rats exposed to cold. Caloric exchange - Am.J. Physiol., 178, 305-308, 1954.
66. CARLSON, L.D., PEARL, D. and SCHEYER, W.: Effects of temperature and work on metabolism and heat loss in man (abstract) - Am. J. Physiol., 179, 625, 1954.
67. CARLSON, L.D. : Adaptation to cold (abstract) - Bull. Meteorol., 35, 281, 1954.
68. CARLSON, L.D. : Man in cold environment. A study in physiology - Office of Technical Services, U.S. Department of Commerce, Washington, 13, 1954.
69. CARLSON, L.D., SHERWOOD, W. and ELSNER, R.: Adequate stimulus for shivering (abstract) - Fed. Proc., 13, 23, 1954.
70. CARLSON, L.D. : Changes in peripheral circulation with exposure to cold - Peripheral Circulation in Man (Ciba Foundation Symposium), 92-104, 1954.
71. CARLSON, L.D. : The adequate stimulus for shivering - Proc.Soc.exp.Biol., N.Y., 85, 303-305, 1954.
72. CARLSON, L.D. : Interrelationship of circulatory and metabolic factors - Cold Injur. Transactions of the Third Conf., Febr. 22, 23, 24 and 25, 1954. Josiag Macy Jf. Foundation, 13-51, 1955. M.I. Ferrer, Ed. New York.
73. CARLSON, L.D., YOUNG, A.C. and BURNS, H.L.: Regional heat loss by temperature gradient calorimetry - Arctic Aeromed. Lab. Proj.Rep., 13, 1955.
74. CARLSON, L.D. : Heat production and cold adaptation (abstract) - Fed.Proc., 14, 26, 1955.
75. CARLSON, L.D. and COTTLE, M.: Turnover of thyroid hormone in cold-exposed rats determined by radioactive iodine studies - Endocrinology, 59, 1 - 11, 1956.
76. CARLSON, L.D. and COTTLE, W.: Chemical regulation of heat production in cold adapted rats (abstract) - Fed. Proc., 15, 41-42, 1956.
77. CARLSON, L.D. and COTTLE, W.: Regulation of heat production in cold-adapted rats - Proc. Soc. exp. Biol., N.Y., 92, 845-849, 1956.
78. CARLSON, L.D. and PEARL Jr., D.C.: The effects of temperature and work on metabolism and heat loss in man - Proc.Soc. exp. Biol., N.Y., 91, 240-244, 1956.
79. CARLSON, L.D., PEARL, D.C. and SHERWOOD, W.W.: Mechanism of oxygen deficit - Proc.Soc.exp. Biol., N.Y., 92, 277-281, 1956.
80. CARLSON, L.D. and WOODS, R.: Thyroxine secretion in rats exposed to cold - Endocrinology, 59, 323-330, 1956.
81. CARLSON, L.D. and BUETTNER, K.: Thermal stress and physiological strain - Fed.Proc. 16, 2, 1957.
82. DEPOCAS, F. and HART, J.S.: Glucose metabolism of rats exposed to cold environments - Am. Phys. Soc. Proc.; Am. J. Physiol. 183, 609, 1955.
83. DEPOCAS, F. : Metabolic response of warm and cold acclimated rats to very cold environments - Fed.Proc., March 1956.

84. DESMARAI, A. : Effet de l'ablation partielle du foie sur la résistance au froid et à la chaleur - Revue Canadienne de Biologie, 2, 332, 1943.
85. DESMARAI, A. : Le froid et le travail musculair ont-ils les mêmes exigences vis-à-vis des surrénales ? - Revue Canad. de Biologie, 7, 662, 1948.
86. DESMARAI, A. : La circulation périphérique chez le rat blanc exposé au froid - Revue Canad. de Biologie, 9, 207, 1950.
87. DESMARAI, A. : Circulation périphérique et teneur des surrénales en adrénaline et en noradrénaline chez le rat blanc exposé au froid - Can.J.Med., 29, 90, 1951.
88. DESMARAI, A. : Influence de l'administration d'adrénaline et de noradrénaline sur l'hypertrophie de la surrénale au froid - Can.Jl.Med.Sc., 29, 104, 1951.
89. DESMARAI, A. : Acide ascorbique, consommation d'oxygène et teneur en glycogène du foie chez le rat exposé au froid - Proc.Can.Physiol. Soc., 21, 1952.
90. DESMARAI, A. : Circulation périphérique et adaptation au froid - Rev.Can. de Biol., 12, 96, 1953.
91. DESMARAI, A. : Effets de l'acide ascorbique sur la consommation d'Oxygène et la teneur en glycogène du foie chez le rat blanc exposé au froid ou maintenu à la température normale - Rev.Can. de Biologie, 12, 99, 1953.
92. DESMARAI, A. : Activité oxydative de différents tissus chez le rat blanc exposé au froid - Rev.Can. de Biol., 13, 115, 1954.
93. DESMARAI, A. : Influence of exposure to cold and ascorbic acid administration on the weight and the metabolic activity of the adrenal gland of hypophysectomized rats given ACTH - Can. J. Biochem. & Physiol., 33, 677, 1955.
94. DESMARAI, A. : Skin metabolism in rats exposed to cold - Can.J. Biochem. & Physiol. 33, 758, 1955.
95. DESMARAI, A. : Relations thyroïde-surrénales dans l'adaptation au froid - Can. J. Biochem. & Physiol., 33, 1018, 1955.
96. DESMARAI, A. : Effet de l'hypophysectomie, de l'administration d'ACTH et de l'ACTH endogène sur le poids et l'activité métabolique de la surrénale du rat blanc - Rev. Can. de Biolog., 14, 108, 1955.
97. DESMARAI, A. : Relations between the thyroid, the adrenals and ascorbic acid administrations in animals exposed to cold - Areal report: Symposium Int. Soc. Bioclim. and Biomet., Paris, 29-31 August 1956.
98. DESMARAI, A. : Conditionnement par l'exposition au froid de la réponse thyroïdienne à la thyrotrophine chez le rat hypophysectomisé - Compt.rend. Soc. biol. Paris, 149, 1714, 1955.
99. DESMARAI, A. : Effet de l'hormone somatotrope (STH) chez le rat surrénalectomisé exposé au froid - Compt. rend. Soc. biol. Paris, 149, 1722, 1955.
100. DESMARAI, A. : Influence de l'exposition au froid sur l'activité thyroïdienne et sur le poids et la teneur en acide ascorbique de surrénales chez le cobaye carencé ou non en acide ascorbique - Compt. rend. Soc. biol., Paris, 149, 1726, 1955.
101. DRIEDGER, G. : Untersuchungen über den respiratorischen Gaswechsel unter dem Einfluss von Abhärtung und Verweichlichung - Diss.med.Kiel, 1948.
102. DRISCHER, H. : Zur physiologischen Problematik vegetativer Regulationsprüfungen - Z. ges.inn. Med., 7, 1060-1065, 1952.
103. DUGAL, L.P., DESMARAI, A. and LEBLOND, C.P.: Effet de l'ablation partielle du foie sur la résistance au froid d'animaux adaptés - Revue Can. de Biol., 2, 4, 453-454, 1943.
104. DUGAL, L.P. and LEBLOND, C.P.: Manifestations pathologiques produites par le froid au niveau des reins et des extrémités - Revue Can. de Biol., 2, 5, 542-545, 1943.
105. DUGAL, L.P., LEBLOND, C.P. and THERIEN, M.: Les aliments choisis par le rat blanc au froid et à la chaleur - Rev. Can. de Biol., 3, 1, 127-129, 1944.
106. DUGAL, L.F. and THERIEN, M.: The increase of ascorbic acid contents of the tissues of rats exposed to cold temperature - Proc. of the Can. Physiol. Soc. Can. Med. Jl., 54, 69-72, 1945.
107. DUGAL, L.P., LEBLOND, C.P. and THERIEN, M.: Resistance to extreme of temperatures in connection with different diets - Can.Jl. of Research, E 23, 244 - 258, 1946.
108. DUGAL, L.P. and THERIEN, M.: Ascorbic acid and acclimatisation to cold environment - Can. Jl. of Research, E.25, 111-136, 1947.
109. DUGAL, L.P. and THERIEN, M.: La nécessité de l'acide ascorbique pour l'acclimatation au froid du cobaye - Rev. Can. de Biol., 6, 1, 548-551, 1947.
110. DUGAL, L.P. and THERIEN, M.: Relation entre la teneur en acide ascorbique des organes et l'acclimatation au froid chez le cobaye - Rev.Can. de Biol., 6, 3, 552-553, 1947.

11. DUGAL, L.P. and DESMARAIS, A. : Exigences différentes du froid et du travail musculaire vis-à-vis des surrénales - Rev.Can. de Biol., 7, 5, 662-665, 1948.
12. DUGAL, L.P. and THERIEN, M.: The influence of ascorbic acid on the adrenal weight during exposure to cold - Endocrinology, 44, 420-427, 1949.
13. DUGAL, L.P. and THERIEN, M.: Excrétion urinaire de l'acide ascorbique chez le rat et le cobaye exposés au froid - Rev. Can. de Biol., 8, 3, 248-257, 1949.
14. DUGAL, L.P. and THERIEN, M.: Effet de l'acide ascorbique sur l'hypertrophie de la surrénale pendant l'exposition au froid: répercussions sur le thymus et la thyroïde - Rev. Can. de Biol., 8, 3, 309, 1949.
15. DUGAL, L.P. and HEROUX, O.: L'acide ascorbique est-il un régulateur de la pression artérielle dans un climat froid ? - Rev.Can. de Biol., 8, 3, 310, 1949.
16. DUGAL, L.P. and THERIEN, M.: Influence de l'acide ascorbique sur l'activité de la surrénale au froid - Rev. Can. de Biol., 8, 3, 315-316, 1949.
17. DUGAL, L.P. and LEBLANC, J.: Influence de l'acide ascorbique sur les variations du cholestérol surrénalien d'animaux exposés au froid - Rev. Can. de Biol., 8, 4, 440-443, 1949.
18. DUGAL, L.P. and THERIEN, M.: Teneur des tissus en acide ascorbique chez le rat partiellement exposé au froid intense - Rev.Can. de Biol., 8, 4, 1949.
19. DUGAL, L.P., THERIEN, M., LEBLANC, J. and HEROUX, O.: Effet de l'acide ascorbique sur plusieurs variables biologiques normalement affectées par le froid - Can. J. of Research, E. 27, 6, 1949.
20. DUGAL, L.P. and DESMARAIS, A.: La circulation périphérique chez le rat blanc exposé au froid - Rev. Can. de Biol., 9, 3, 206, 1950.
21. DUGAL, L.P. : Effects of cold, aged and ascorbic acid on formaldehyde-induced arthritis - Can. J. Med. Sciences, 29, 35, 1951.
22. DUGAL, L.P. and DESMARAIS, A.: Variations de la circulation périphérique chez le rat blanc adapté au non au froid - Can. J. Med. Sc., 29, 3, 1951.
23. DUGAL, L.P. and DESMARAIS, A.: Variations de la teneur en adrénaline et artérénol chez le rat blanc exposé au froid - Can. J. Med. Sc., 29, 3, 90, 1951.
24. DUGAL, L.P. and DESMARAIS, A.: Circulation périphérique et teneur des surrénales en adrénaline et en artérénol (noradrénaline) chez le rat blanc exposé au froid - Can. J. Med. Sc., 29, 90-99, 1951.
25. DUGAL, L.P. and DESMARAIS, A.: Influence de l'administration d'adrénaline et d'artérénol sur l'hypertrophie de la surrénale au froid - Can.J. Med. Sc., 29, 104-107, 1951.
26. DUGAL, L.P. and FORTIER, G.: Ascorbic acid and acclimatization to cold in monkeys - Proc. Can. Physiol. Soc. Rev. Can. Biol., 11, 59, 1952.
27. DUGAL, L.P. and FORTIER, G.: Adrénaline et résistance au froid - Rev.Can.de Biol., 11, 61, 1952.
28. DUGAL, L.P. and THERIEN, M.: Effets de l'acide glutanique et d'extraits de foie sur la résistance du rat blanc à un froid intense - Rev.Can. de Biol., 11, 180, 1952.
29. DUGAL, L.P. and FORTIER, G.: Ascorbic acid and acclimatization to cold in monkeys - Journ. of Applied Sc., 5, 3, 1952.
30. DUGAL, L.P. and FORTIER, G.: Nécessité de l'acide ascorbique pour l'acclimatation du singe (Rhésus) au froid - Comptes Rendus Acad.Sc., Paris, 234,138, 1952.
31. DUGAL, L.P. and THERIEN, M.: L'acide ascorbique et l'activité de la surrénale chez le rat hypophysectomisé - Comptes Rendus Acad. Sc. Paris, 234, 245, 1952.
32. DUGAL, L.P. and THERIEN, M.: L'acide ascorbique et l'activité de la surrénale chez le rat normal et chez le rat hypophysectomisé - Annales d'Endocrinologie, 13, 4, 578-588, 1952.
33. DUGAL, L.P. and DESMARAIS, A.: Circulation périphérique et adaptation au froid - Revue Can. de Biol., 12, 1, 96-99, 1953.
34. DUGAL, L.P. and FORTIER, G.: Prévention des gelures par l'acide ascorbique chez le singe - Rev. Can. de Biol., 12, 1, 97, 1953.
35. DUGAL, L.P. and FORTIER, G.: Résistance et acclimatation au froid chez le singe - Gazette Médicale de France, 60, 1, 1953.
36. DUGAL, L.P. : Ascorbic acid and resistance to cold - Cold Injury. Second Conference Josiah Macy Foundation, New York, 85, 134, 1954.
37. DUGAL, L.P. : Ascorbic acid and acclimatization to cold - Symposium on Nutrition and climatic Stress, Washington, Nat Academy of Science, National Research Council, 70-81, 1954.
38. DUGAL, L.P. and DUFOUR, D.: Maintien, par l'hormone somatotrope de la croissance normale du rat exposé au froid - C.R. Soc. Biol., 148, 1521, 1954.
39. DUGAL, L.P., DESMARAIS, A. and CAGNON, P.M.: Influence of exposure to cold and ascorbic acid administration on the weight and the metabolic activity of the adrenal gland of hypophysectomized rats given ACTH - C.J.Bioch. Physiol., 33, 677, 1955.

140. DUGAL, L.P., DESMARAIS, A. and CAGNON, P.M.: Effets de l'hypophysectomie, de l'administration d'ACTH et de l'ACTH endogène sur le poids et l'activité métabolique de la surrénale du rat blanc - *Rev.Can.Biol.*, 14, 108, 1955.
141. ELSNER, R.W. : Some acclimatizing responses of man to prolonged cold exposure - *Arctic Aeromed. Lab.*, Ladd AFB, 1, 1955.
142. FAHNESTOCK, M.K. : Thermocouples and the measurement of body temperatures in the laboratory and in flight - *The Jl. of Aviation Med.*, 17, 2, 1946.
143. FAHNESTOCK, M.K., MITCHELL, H.H., GLICKMAN, N., LAMBERT, E.H. and KEETON, R.W.: The tolerance of man to cold as affected by dietary modification: Carbohydrate versus fat and the effect of the frequency of meals - *The Am. Jl. of Physiol.*, 146, 1, 1946.
144. FAHNESTOCK, M.K., KEETON, R.W., LAMBERT, E.H., GLICKMAN, N., MITCHELL, H.H. and LAST, J.H.: The tolerance of man to cold as affected by dietary modifications. Proteins Versus carbohydrates, and the effect of variable protective clothing - *The Am.Jl. of Physiol.*, 146, 1, 1946.
145. FAHNESTOCK, M.K., GLICKMAN, N., KEETON, R.W. and MITCHELL, H.H.: The tolerance of man to cold as affected by dietary modifications: High versus low intake of certain water soluble vitamins - *The Am.Jl. of Physiol.*, 146, 4, 1946.
146. FAHNESTOCK, M.K. a.o.: Physiological adjustments of human beings to sudden change in Environment - *ASHVE Journal Section of Heating, Piping and Air Conditioning*, 19, 7, 101-115, 1947.
147. FAHNESTOCK, M.K. a.o.: Comparison of physiological adjustments of human beings during summer and winter - *ASHVE Journal Section of Heating, Piping and Air Conditioning*, Aug. 1948.
148. FAHNESTOCK, M.K. a.o.: Physiologic adjustments of normal subjects and cardiac patients to sudden change in environment - *ASHVE Jl. Section of Heating, Piping, and Air Conditioning*, Febr. 1949.
149. FAHNESTOCK, M.K. a.o.: Physiologic examination of the effective temperature Index - *ASHVE Jl. Section of Heating, Piping, and Air Conditioning*, 22, 1, 157, 1950.
150. FAHNESTOCK, M.K. a.o.: Physiological adjustment of clothed human being to sudden change in environment - *ASHVE Jl. Section Heating, Piping, and Air Conditioning*, 2, 107, 1952.
151. FAHNESTOCK, M.K. a.o.: Controlling the environment for clinical research - *ASHVE Jl. Section of Heating, Piping and Air Cond.*, 1, 123, 1953.
152. FREGLY, M.J. : Minimal exposures needed to acclimatize rats to cold - *Am.J.Physiol.*, 173, 393-402, 1953.
153. FREGLY, M.J. : Minimal exposures needed to acclimatize rats to cold atmospheres - *J. Nat. Cancer Inst.*, 13, 1390, 1953.
154. FREGLY, M.J. : Cross-acclimatization between cold and altitude in rats - *Am. Jl. Physiol.*, 176, 267, 1954.
155. FREGLY, M.J. : The effects of extremes of temperature on hypertensive rats - *Am. Jl. Physiol.*, 176, 275, 1954.
156. FREGLY, M.J. and KNOBIL, E.: Effect of ascorbic acid on the adrenal gland after hypophysectomy and after exposure to cold - *Endocrin.*, 56, 614, 1955.
157. FREGLY, M.J., IAMPIETRO, P.F. and BUSKIRK, E.R.: Effect of administration of adrenal cortical hormones on cooling rate of rats - *Am.J. Physiol.*, 183, 629, 1955.
158. FUKUDA, M. : Studies on seasonal variation of thermoregulatory function of human body, Part.I. - *J. Physiol. Soc. of Japan*, 15, 68-80, 1953.
159. FUKUDA, M. : Studies on seasonal variation of thermoregulatory function of human body. Part.II. On sexual differences of thermoregulatory functions - *J. Physiol. Soc. of Japan*, 15, 150-157, 1953.
160. FUKUDA, M. : Studies on seasonal variation of thermoregulatory function of human body. Part. III. Changes of thermoregulatory function with age - *J. Physiol. Soc. of Japan*, 15, 417-427, 1953.
161. GELINEO, S. and GIAJA, I.: L'Hypothermie et la thermogénèse - *Arch. intern. Physiol.*, 32, 237-250, 1930.
162. GELINEO, S. and GIAJA, I.: Influence d'un séjour prolongé à une température élevée sur la puissance thermogénétique - le métabolisme du sommet - de l'homeotherme - *Annales de Physiologie*, 7, 163-168, 1931.
163. GELINEO, S. and GIAJA, I.: La thermogénèse aux températures inférieures à celle de la résistance homéotherme - *Annales de Physiologie*, 7, 172-182, 1931.
164. GELINEO, S. : Facteur thermique de la croissance des homéothermes - *Bull. Acad. r. serbe*, B. 1, 103-110, 1933.
165. GELINEO, S. and GIAJA, I.: Sur la résistance au froid de quelques homéothermes - *Arch. intern. de Physio.*, 37, 1-15, 1933.
166. GELINEO, S. : Influence du milieu thermique d'adaptation sur la thermogénèse des homéothermes - *Annales de Physiologie*, 10, 1083-1115, 1934.

67. GELINEO, S. : La thermogénèse dans l'hyperthermie exogène - C.r. Soc.Biol., 115, 865, 1934.
68. GELINEO, S. : Influence du milieu thermique d'adaptation sur la courbe de la thermorégulation - C.r. Soc. Biol., 117, 40, 1934.
69. GELINEO, S. and GIAJA, I.: Alimentation et résistance au froid - C.r. des séances de l'Académie des Sciences Paris, 198, 2277, 1934.
70. GELINEO, S. : Der Sauerstoffverbrauch bei Behaglichkeitstemperatur nach exogener Hyperthermie - Biologia Generalis, 12, 243-244, 1936.
71. GELINEO, S. and GIAJA, I.: Sur la résistance au froid de quelques homéothermes - Bull. de l'Acad.r. serbe. B.2, 53-70, 1935.
72. GELINEO, S. and GIAJA, I.: Sur la résistance de quelques homeothermes aux basses températures - C.r. des séances de l'Acad. des Sciences, Paris, 200, 2115, 1935.
73. GELINEO, S. and GIAJA, I.: Sur les causes de la mort par inanition - C.r. Soc.Biol., 119, 641, 1935.
74. GELINEO, S. : Sur l'adaptation de la thermogénèse aux températures provoquant l'hyperthermie - C.r. Soc. Biol., 119, 643, 1935.
75. GELINEO, S. : La métabolisme de base et l'hyperthermie exogene de longue durée - C.r. Soc. Biol., 119, 645, 1935.
76. GELINEO, S. : Tension d'oxygène et resistance au froid - Bull. de l'acad.r.serbe , B. 3, 189-192, 1936.
77. GELINEO, S. : Sur la "seconde thermorégulation chimique" - C.r. Soc. Biol., 122, 337, 1936.
78. GELINEO, S. and GIAJA, I.: Sur la courbe réèle de la thermogénèse des homéothermes - C. r. Soc. Biol., 122, 343, 1936.
79. GELINEO, S. and GIAJA, I.: Pression barométrique et résistance au froid - C.r. Acad. des Sciences, Paris, 202, 1613, 1936.
80. GELINEO, S. : Pression barométrique et thermogénèse - Bull. Acad.r. serbe,B3, 193-199, 1937.
81. GELINEO, S. : Sur la thermorégulation chimique aux températures provoquant l'hyperthermie - Bull. Acad. r. serbe, B. 3, 209-234, 1937.
82. GELINEO, S. : Anpassungstemperatur und Grundumsatz - XVe Congrès internat. d'Hydrologie, de Climatologie et de Geologie-médicales, Beograd,341-349, 1939.
83. GELINEO, S. and RAJEVSKAJA, T.:Influence d'un régime alimentaire de maïs sur la concentration de l'hémoglobine à des températures différentes - C.r.Soc.Biol., 147, 136, 1953.
84. GELINEO, S. : Influence du climat de la cote orientale d'Adriatique sur les échanges gazeux de l'homme - Congrès internat. d'Hydro-climatisme et de thalasso-thérapie, Resumés, Opatija, 8-13 mai 1954, 5-6.
85. GELINEO, S. : Homéothermie et métabolisme de base dans l'ontogénie - Archives des Sciences biologiques Beograd, 6, 227-230, 1954.
86. GELINEO, S. : Température d'adaptation, concentration d'hémoglobine et production de chaleur chez le chien - Arch. des Sci. Biol., Beograd,6, 235-243, 1954.
87. GELINEO, S. : Contribution to the study of the effect of climate on basal metabolism - Bull. de l'Acad. serbe des Sciences, 11, 56-70, 1954.
88. GELINEO, S. : The effect of the climate of the Croatian Littoral on the respiratory quotient in man - Bull. de l'Acad. serbe des Sciences, 11,71-73,1954.
89. GELINEO, S. : La concentration d'hémoglobine chez les travailleurs yougoslaves en 1950 - Bull. de l'Acad. serbe des Sciences, 11, 109-119, 1954.
90. GELINEO, S. : Le développement ontogénétique de la thermorégulation chez le chien - C.r. Soc. Biol., 148, 1483-1485, 1954.
91. GELINEO, S. : La température d'adaptation, la concentration de l'hémoglobine et la production de chaleur chez le chien - C.r. Soc.Biol. 148, 1954-1956, 1954.
92. GELINEO, S. and KOCAREV, R.: Influence de le température d'adaptation sur l'intensité des échanges respiratoires chez le rat au cours des différentes saisons - C.r. Soc. Biol., 149, 1652-1654, 1955.
93. GLASER, E.M. : The effects of cooling and of various means of warming on the skin and body temperature of men - J. Physiol., 109, 366, 1949.
94. GLASER, E.M. : The effects of cooling and warming on the vital capacity, forearm and hand volume, and skin temperature of man - J. Physiol., 109,421,1949.
95. GLASER, E.M. : Immersion and survival in cold water - Nature, 166, 1068, 1950.
96. GLASER, E.M., BERRIDGE, F.R. and PRIOR, K.M.: Effects of heat and cold on the distribution of blood within the human body - Clin.Sci., 9, 2, 1950.
97. GLASER, E.M. and HOLMES JONES, R.V.: Initiation of shivering by cooled blood returning from the lower limbs - J. Physiol., 114, 277, 1951.

198. GLASER, E.M. : Principles underlying the regulation of animal heat - XIX Internat. Physiol. Congr. 1953, Abstract of communications, 395.
199. GLASER, E.M. and NEWLING, P.S.B.: Thermal balance in man - J. Physiol. Proc., 1955.
200. GOETZ, A. : Reaction: Live matter to low temperatures - Refrigerating Engineering Nov., 1955.
201. GROBER, J. : Die Akklimatisation - G. Fischer, Jena 1934.
202. GROBER, J. : Der weisse Mensch in Afrika und Südamerika - G. Fischer, Jena 1938.
203. GROBER, J. : Die Akklimatisation der Europäer in Afrika - Handbuch: Afrika, herausgeg. von E. Obst, Verlag de Gruyter, Berlin, 1940.
204. GROBER, J. : Grundfragen der tropischen Bioklimatik - Ber.d. dtsh. Wetterdienst i.d. US-Zone, 42, 1952.
205. GROBER, J. : Bioklimatische Entsteppung - Wiss. Zeitschr. d. Univ. Jena, 3/4, 1953/1954.
206. GROBER, J. : Schleimhautleistung und Akklimatisation - Zeitschr. f. Trop.-Med. u. Parasitolog., 1, 1954.
207. GROBER, J. : Triennium Bioclimaticum Tropicale - Forschungen und Fortschritte, 6, 1955 und Forschungs- und Sitzungsberichte der Akademie für Raumforschung und Landesplanung, 5, 2, 1956.
208. GUTTMAN et al. : Thermoregulation in spinal man - Journal of Physiology, 142, 3, 406-420, 1958.
209. HALE, H.B. and MEFFERD Jr., R.B.: Factorial study of environmentally-induced metabolic changes in rats - Am.J. Physiol., 194, 469-475, 1958.
210. HANDLINGER, K. : Der Einfluss der Umwelttemperatur auf die Höhe der ED 50 (Stilböstrol) im Allen-Doisy-Test der Maus - Med.Vet.Diss. Wien, 1955.
211. HANNON, J.P., LARSON, A.F., DRURY, H.F., VAUGHAN, D.A. and VAUGHAN, L.N.: Arctic survival rations. VII. Alterations in serum electrolytes during winter field tests of survival rations - Alaskan Air Command, Arctic Aeromed. Lab., Ladd Air Force Base, Techn. Report, 58, 25, Sept. 1959.
212. HANNON, J.P. and VAUGHAN, L.N.: Gross liver composition of the rat following one-month cold exposure - Alaskan Air Command, Arctic Aeromed. Lab., Ladd Air Force Base, Techn. Report 58, 26, Sept. 1959.
213. HANNON, J.P. : The respiration of rat liver homogenates following prolonged cold exposure - Alaskan Air Command, Arctic Aeromed. Lab., Ladd Air Force Base, Techn. report 57, 59, Sept. 1959.
214. HANNON, J.P., LARSON, A.M. and YOUNG, D.W.: Blood chemistry of the cold-acclimatized rat. I. Plasma electrolytes - Alaskan Air Command, Arctic Aeromed. Lab., Ladd Air Force Base, Techn. Report 57, 60, Nov. 1959.
215. HENSEL, H. : An apparatus for the continuous recording of the integral skin temperature and the skin temperatures of various places of the body - Submarine Med. Mon. Annex II, S.I. 1948.
216. HENSEL, H. and SCHAEFFER, K.E.: The heat regulation aboard submarines - Submarine Med.Mon., C 8, 27, 1948.
217. HENSEL, H. : Ein vollautomatischer Thermointegralschreiber - Pflügers Arch., 252, 103-106, 1949.
218. HENSEL, H. : Auslösung von Kältezittern durch Kohlensäureatmung - Pflügers Arch., 252, 107-110, 1949.
219. HENSEL, H. : Ein Gerät zur fortlaufenden Registrierung der integralen Hauttemperatur und der Hauttemperaturen einzelner Körperstellen - Pflügers Arch., 251, 388-397, 1949.
220. HENSEL, H. : Die intracutane Temperaturbewegung bei Einwirkung äusserer Temperaturereize - Pflügers Arch., 252, 146-164, 1950.
221. HENSEL, H. : Temperaturempfindung und intracutane Wärmebewegung - Pflügers Arch., 252, 165-215, 1950.
222. HENSEL, H. and ZOTTERMAN, Y.: The cold receptor discharge and the intracutaneous temperature gradient - Abstr. XVIII, Internat. Physiol. Congr. Copenhagen, 249, 1950.
223. HENSEL, H. and ZOTTERMAN, Y.: The persisting cold sensation - Abstr. XVIII. Internat. Physiol. Congr. Copenhagen, 250, 1950.
224. HENSEL, H. : Intracutane Wärmeleitfähigkeits- und Temperaturmessungen - Abstr. Intern. Physiol. Congr. Copenhagen, 541, 1950.
225. HENSEL, H. : Ein Strömungscalorimeter für beliebige Körperstellen - Z.Exper. Med. 117, 587-597, 1951.
226. HENSEL, H. and ZOTTERMAN, Y.: The effect of menthol on the thermoreceptors - Acta Physiol. Scand., 24, 27-31, 1951.
227. HENSEL, H. and ZOTTERMAN, Y.: The response of the cold receptors to constant cooling - Acta Physiol. Scand., 22, 96-105, 1951.
228. HENSEL, H. and ZOTTERMAN, Y.: The persisting cold sensation - Acta Physiol.Scand., 22, 106-113, 1951.

29. HENSEL, H. and ZOTTERMAN, Y. : Quantitative Beziehungen zwischen der Entladung einzelner Kältefasern und der Temperatur - *Acta Physiol. Scand.* 23, 291 - 319, 1951.
30. HENSEL, H. and ZOTTERMAN, Y.: Action potentials of cold fibres and intracutaneous temperature gradient - *J. of Neurophysiol.*, 14, 377-385, 1951.
31. HENSEL, H. a.o. : Electro-physiological measurements of depth of thermoreceptors - *J. of Neurophysiol.*, 14, 423-429, 1951.
32. HENSEL, H. and ZOTTERMAN, Y. : The response of mechanoreceptors to thermal stimulation - *J. of Physiol.*, 115, 16-24, 1951.
33. HENSEL, H. : Physiologie der Thermoreception - *Erg. Physiol.*, 47, 166-368, 1952.
34. HENSEL, H. : Wärme- und Temperaturleitung in der lebenden Haut - *Experientia*, 8, 477-481, 1952.
35. HENSEL, H. : Afferente Impulse aus den Kälterezeptoren der äusseren Haut - *Pflügers Arch.*, 256, 195-211, 1952.
36. HENSEL, H. : Ein neues Verfahren zur peripheren Durchblutungsregistrierung an beliebigen Körperstellen - *Z. Kreislaufforschg.*, 41, 251-261, 1952.
37. HENSEL, H. : The time factor in thermoreceptor excitation - *Acta Physiol. Scand.* 29, 109-116, 1953.
38. HENSEL, H. and BRUECK, K.: Wärmedurchgang und Innentemperatur der menschlichen Extremitäten - *Pflügers Arch.* 257, 70-86, 1953.
39. HENSEL, H. : Das Verhalten der Thermorezeptoren bei Ischämie - *Pflügers Arch.* 257, 371-383, 1953.
40. HENSEL, H. : Das Verhalten der Thermorezeptoren bei Temperatursprüngen - *Pflügers Arch.*, 256, 470-487, 1953.
41. HENSEL, H. : Der Wärmehaushalt des Menschen und seine Unterstützung durch Wohnung und Kleidung - Mensch und Wärme, Bericht über eine Vortragsstagung des Warmwirtschaftsverbandes Baden-Württemberg, Stuttgart, 11-25, 1954.
42. HENSEL, H. : Biologische Regelungsvorgänge - *Umschau*, 54, 289-292, 1954.
43. HENSEL, H. : Mensch und Warmblütige Tiere - in: H. Precht, J. Christophersen und H. Hensel: Temperatur und Leben, 329-466. Springer Verlag Berlin-Göttingen-Heidelberg, 1955.
44. HENSEL, H. : Temperatur und Leben. Teil III. - Berlin-Göttingen, Springer Verlag, 1955.
45. HENSEL, H. : Ueber die Steuerung der peripheren Durchblutung - *Arch. Physikal. Ther.* 7, 60-74, 1955.
46. HENSEL, H. a.o. : Die Muskeldurchblutung des Menschen bei indirekter Erwärmung und Abkühlung - *Pflügers Arch.*, 261, 199-200, 1955.
47. HENSEL, H. and BOCK, K.D.: Durchblutung und Wärmeleitfähigkeit des menschlichen Muskels - *Pflügers Arch.*, 260, 361-367, 1955.
48. HENSEL, H. : Fortlaufende Aufzeichnung der Muskeldurchblutung beim Menschen - *Umschau*, 55, 132-133, 1955.
49. HERMESMEIER, W. : Ueber normale Temperaturdifferenzen an symmetrischen Hautstellen - *Diss. med. Kiel*, 1937.
50. HEROUX, O. and HART, J.S.: A comparison of some seasonal and temperature-induced changes in *Peromyscus*: cold resistance, metabolism and pelage insulation - *Can. Jr. of Zool.*, 31, 528-534, 1953.
51. HEROUX, O. and HART, J.S.: Effect of low temperature and work on blood lactic acid in deer mice - *Am. Jl. Physiology*, 176, 452-454, 1954.
52. HEROUX, O. and HART, J.S.: Restraint hypothermia and its inhibition by cold acclimation - *Am. Jl. Phys.*, 177, 219-221, 1954.
53. HEROUX, O. and HART, J.S.: Comparison of four indices of adrenal activity in rats acclimated to 30°, 15°, 1° C. - *Am. Jl. Phys.*, 178, 445-448, 1954.
54. HEROUX, O. and HART, J.S.: Adrenal cortical hormone requirement of warm and cold acclimated rats after adrenalectomy - *Am. J. Phys.*, 178, 449-452, 1954.
55. HEROUX, O. and HART, J.S.: Cold acclimation and adrenal cortical activity as measured by eosinophil levels - *Am. J. Phys.*, 178, 453-456, 1954.
56. HEROUX, O. : Acclimation of adrenalectomized rats to low environmental temperature - *Am. J. Phys.*, 181, 75-78, 1955.
57. HEROUX, O. and HART, J.S.: Exercise and temperature regulation in lemmings and rabbits - *Can. J. of Bioch. and Physiol.*, 33, 428-435, 1955.
58. HEROUX, O. : Physiology of cold Acclimation - Areal report: Symposium Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956.
59. HERRINGTON, L.P. and HARDY, J.D.: Temperature and humidity in relation to the thermal interchange between the human body and the environment - *Human Factors in Undersea Warfare*. National Research Council, 1949.
60. HERRINGTON, L.P. : Biophysical adaptations of man under climatic stress - *Meteorological Monographs*, 2, 8, 30-42, 1954.

261. HICK, F.K. : Partial pressure of oxygen in arterial blood of patients. Description of an Aerotonometer Method - Proc. Soc. of Exp. Biol. & Med., 33, 582-587, 1936.
262. HICK, F.K. a.o. : Physiologic response of man to environmental temperature - Heating, Piping and Vent., 145, 1938.
263. HICK, F.K. a.o. : Circulation through the periphery to facilitate heat losses - Proc. Cent. Soc. for Clin. Res., 11, 1938.
264. HICK, F.K. a.o. : Cardiac output, peripheral blood flow and blood volume changes in normal individuals subjected to varying environmental temperatures - Heating, Piping and Air Conditioning, 123, 1939.
265. HICK, F.K. a.o. : Mechanism of circulatory collapse in heat exhaustion - Proc. Cent. Soc. f. Clin. Res., 12, 1939.
266. HICK, F.K. a.o. : The peripheral type of circulatory failure in experimental heat exhaustion - Heating, Piping and Air Conditioning, 157, 1940.
267. HICK, F.K. a.o. : Adjustments in circulating blood volume in human subjects exposed to hot environments - Proc. Cent. Soc. f. Clin. Res., 13, 1940.
268. HICK, F.K. a.o. : Blood volume changes in man exposed to hot environmental conditions for a few hours - Am. Jl. Physiol., 134, 165, 1941.
269. HICK, F.K. a.o. : Physiological adjustments of human beings to sudden change in environment - ASHVE Journal Section of Heating, Piping and Air Conditioning, 19, 7, 101-115, 1947.
270. HICK, F.K. a.o. : Physiologic adjustments of human being to sudden changes in environment - Heating, Piping and Air Cond., 1948.
271. HICK, F.K. a.o. : Physiological responses to sudden changes in atmospheric environment - Am. Soc. for Heat & Vent. Engineers Journal, Section Heating, Piping and Air Cond., 1954.
272. HILDES, J.A. : Cold acclimatization in humans - Areal report: Symposium Int. Soc. Bioclim. and Biomet., Paris 29-31 August 1956.
273. HILDES, J.A. : Some physiological aspects of arctic warfare - Can. Services Med. Jl. 12, 776-786, 1956.
274. HUNTER, J., CLARK, W.D. and WHILLANS, M.G. (introduced by JOHNSON, W.H.): A study of the effect of Trafuril on fingers exposed to cold - Proc. Can. Physiol. Soc., 14th Ann. Meeting, Univ. of Ottawa, Ottawa, Ont., 16 and 17 Oct. 1950, 23.
275. HUNTER, J. and WHILLANS, M.G. (Introduced by JOHNSON, W.H.): A study of the effect of cold on joints - Proc. Can. Physiol. Soc., 14th Ann. Meeting, Univ. of Ottawa, Ottawa, Ont., 16 and 17 October 1950.
276. HUNTER, J. and WHILLANS, M.G.: A study of the effect of cold on joint temperature and mobility - Can. Jnl. of Med. Sciences, 29, 255-262, 1951.
277. HUNTER, J. and WHILLANS, M.G.: Effect of cold on joints - Fed. Proc., 10, 68, 1951.
278. HUNTER, J., KERR, E.H. and WHILLANS, M.G.: The relation between joint stiffness upon exposure to cold and the characteristics of synovial fluid - Can. Jnl. of Med. Sciences, 30, 367-377, 1952 ; DRB 32rd Symposium, 243, 1951.
279. HUNTER, J. : Effects of cold on hand activities with special reference to joints and fluid viscosities - Paper presented at Symposium on Protection and Functioning of the Hands in Cold Climates, Natick, Mass., April 1956.
280. HUNTER, J. : Reanimation of rats after cardiac and respiratory arrest due to low body temperature - Can. Jnl. of Biochem. and Physiol., 35, 605 - 613, 1957.
281. HUNTER, J. : Some studies in Hypothermia - A paper presented at the Conf. on Environmental Physiology. U.S. Army Chem. Warfare Lab., Army Chem. Center, Edgewood, Md., 10 and 11 April 1958.
282. IAMPIETRO, P.F., FREGLY, M.J. and BUSKIRK, E.R.: Maintenance of body temperature of res-trained adrenalectomized rats exposed to cold: effect of adrenal cortical hormones - Can. J. Biochem. & Physiol., 34, 721-729, 1956.
283. IIDA, T. : Studies on the temperature reaction of human skin to cold. I. Report. On physiological rôles of the vascular reaction to cold - Journ. Physiol. Soc. of Japan, 11, 73-78, 1949.
284. IIDA, T. : Studies on the temperature reaction of human skin to cold. II. Report. Blood vessels which contribute to the temperature reaction to cold - J. Physiol. Soc. of Japan, 11, 78-85, 1949.
285. IIDA, T. : Studies on the temperature reaction of human skin to cold. III. Report. On reflex nerves of the temperature reaction to cold and mechanism of reaction - Journ. Physiol. Soc. of Japan, 13, 513-517, 1951.
286. IRVING, L. : Temperature of skin in the Arctic as a regulator of heat - Jnl. of Applied Physiology, 7, 4, 1955.

87. ITOH, S. : Examination on the method to determine local sweating (Japanese) - J. Physiol. Soc. Japan, 7, 108, 1942.
88. ITOH, S. : Is sweat inhibition due to cold stimulus observed in pilocarpine sweating ? (Japanese) - J. Physiol. Soc. Jap., 7, 114, 1942.
89. ITOH, S. : Studies on the acclimatization of the sweat glands to tropical climate - J. Physiol. Soc. Jap., 8, 648, 1943.
90. ITOH, S. : Regional variation of sweat volume in tropical inhabitants (Japanese) - Tropical Med., 1, 499, 1943.
91. ITOH, S. : Observation on a patient with thermal anhidrosis (Japanese) - Jnl. Physiol. Soc. Japan., 9, 493, 1944.
92. ITOH, S. : Some observations on the shape of sweat gland in human finger - Nippon Seiri, Z. 10, 123, 1947.
93. ITOH, S. : Periodicity of the elimination of sweat and the rate of its flow - Nippon Seiri, Z. 10, 126, 1947.
94. ITOH, S. : Iontophoresis of several dyestuffs into sweat glands - Nippon Seiri, Z. 10, 128, 1947.
95. ITOH, S. : Penetration of safranin into skin tissues by iontophoresis. Consideration on the function of sweat duct - Nippon Seiri, Z. 10, 130, 1947.
96. ITOH, S. : Histamine in human sweat - Nippon Seiri, Z. 10, 187, 1948.
97. ITOH, S. : Sugar and pyruvic acid in human sweat - Nippon Seiri, Z. 11, 12, 1948.
98. ITOH, S. : Changes of pyruvic acid content in sweat after administration of thiamine and riboflavin - Nippon Seiri, Z. 11, 16, 1948.
99. ITOH, S. : The secretion of sugar and its intermediate substances in sweat and saliva - Jap. J. Physiol., 3, 10, 1952.
100. ITOH, S. : Amino acids in human sweat - Jap. J. Physiol., 2, 3, 248, 1952.
101. ITOH, S. : Chemistry of human sweat - Igaku Syoin, Tokyo, 1953.
102. ITOH, S. : Ammonia in human sweat and its origin - Jap. J. Physiol., 3, 2, 133, 1953.
103. ITOH, S. : The water loss and blood changes by prolonged sweating without intake of food and drink - Jap. J. Physiol., 3, 2, 148, 1953.
104. ITOH, S. : Occurrence of antidiuretic substance in human urine after heavy sweating - Jap. J. Physiol., 3, 4, 277, 1953.
105. ITOH, S. : The release of antidiuretic hormone from the posterior pituitary body on exposure to heat - Jap. J. Physiol., 4, 185, 1954.
106. ITOH, S. : ADH discharge on exposure to heat (Japanese) - Modern Med. 9, 985, 1954.
107. ITOH, S. : Water excretion of diabetes insipidus rats in hot environment - The Nagoya Journal of Med. Science, 19, 4, 1957.
108. JANTZEN, G. : Ueber die Beziehungen zwischen Perspiration und Frigorigraphen bzw. Katathermometerwerten bei verschiedenartiger Exposition des menschlichen Körpers bezüglich Wind- und Sonnenstrahleneinfall - Diss. med. Kiel, 1948.
109. JENSEN, M. : Shelter effect - Investigations into the Aerodynamics of shelter and its effects on climate and crops - Copenhagen, 1954.
110. JOHNSON, B.C., HAMILTON, T.S. and MITCHELL, H.H.: The excretion of pyridoxine, pseudo-pyridoxine and 4-pyridoxic acid in the urine and sweat of normal individuals - J. Biol. Chem., 158, 619, 1945.
111. JOHNSON, B.C., HAMILTON, T.S. and MITCHELL, H.H.: The effect of choline intake and environmental temperature on the excretion of choline from the human body - J. Biol. Chem., 159, 5, 1945.
112. JOHNSON, B.C., SHIELDS, J.B., HAMILTON, T.S. and MITCHELL, H.H.: The excretion of ascorbic acid and dehydroascorbic acid in sweat and urine under different environmental conditions - J. Biol. Chem., 161, 351, 1945.
113. JOHNSON, B.C., MITCHELL, J.B. and HAMILTON, T.S.: The occurrence of inositol and of p-aminobenzoic acid in sweat - J. Biol. Chem., 161, 357, 1945.
114. JUSTUS, M. : Hautfeuchte-Untersuchungen unter kühlen Bedingungen - Diss. med. Kiel, 1952.
115. KANTER, G.S. : Thirst and excretion after salt administration - Fed. Proc. 11, 80, 1952.
116. KANTER, G.S. : Excretion and drinking after salt loading in dogs - Am. J. Physiol. 174, 87, 1953.
117. KANTER, G.S. : Heat and hydropenia; their effects on thirst and chloride regulation in dogs - Am. J. Physiol., 174, 95, 1953.
118. KANTER, G.S. : Volume of distribution of added chloride - J. Applied Physiol., 6, 124, 1953.
119. KANTER, G.S. : The effect of heat on the regulation of body fluids and electrolytes in dogs - Am. J. Physiol., 178, 259, 1954.

320. KANTER, G.S. and LUBINSKI, R.H.: Effect of heat on inorganic phosphate regulation in dogs - *Am.J. Physiol.*, 180, 559, 1955.
321. KANTER, G.S. : Heat and excretion in man - *J. Applied Physiol.*, 7, 533, 1955.
322. KAWAHATA, A. : Observations on the adaptability of human being to the winter climate in Manchuria - *J. Physiol. Soc. Japan*, 1939.
323. KAWAHATA, A. : Amount of chlorides contained in Micronesian sweat - *J. Physiol. Soc. Japan* 1942.
324. KAWAHATA, A. : Further observations on the acclimatization of the human sweat organs to tropical climate - *J. Physiol. Soc. Japan*, 1943.
325. KAWAHATA, A. : Comparative studies of the constitution of natives, native born Japanese children in tropics and home born children - *Jap.Sc.Hyg.*, 1948.
326. KAWAHATA, A. : Studies on the function of human sweat organs. I. Statistical research of the active sweat glands of the human body - *J.Mie.Med.Coll.* 1950.
327. KAWAHATA, A. : Studies on the function of human sweat organs. II. The relation between the amount of chlorides contained in sweat and the function of the sweat glands - *J. Mie.Med. Coll.*, 1950.
328. KAWAHATA, A. : Studies on the function of human sweat organs. III. Excitabilities of the autonomic nervous system as an indicator of the ability to perspire - *Jap. Soc. Hyg.*, 1950.
329. KAWAHATA, A. : Studies on the function of human sweat organs. IV. Normal concentration of the dissolved constituents in human sweat - *Jap.Soc. Hyg.*, 1950.
330. KAWAHATA, A. : Studies on the function of human sweat organs. V. Sweat glands as an excrete organ of lactic and pyruvic acid - *J.Mie.Med.Coll.*, 1951.
331. KAWAHATA, A. : Some observations on sweating of the Aino - *Jap.J.Physiol.*, 1951.
332. KAWAHATA, A. : Concentration of the dissolved constituents in human sweat related to the rate of sweating - *Med. & Biol.*, 1951.
333. KAWAHATA, A. : Concentration of the dissolved constituents in human sweat related to seasons of the year - *Med. & Biol.*, 1951.
334. KAWAHATA, A. : Studies on the function of human sweat organs. VI. Some factors influencing the concentration of the dissolved constituents in human sweat - *Mie. Med. Journal*, 1953.
335. KENNEY, R.A. : The effect of hot, humid environments on the renal function of West Africans - *J. Physiol.*, 118, 25-26, 1952.
336. KUNO, Y. : The physiology of human perspiration - *J. & A. Churchill Ltd.*, London, 1934.
337. KUNO, Y. : Human perspiration - Charles C. Thomas, Springfield, Illinois, 1956.
338. LADELL, W.S.S. : Thermal Comfort - Practitioner, 1950.
339. LADELL, W.S.S. : Physiology of Life and work in high ambient temperature - Symposium, Desert Research, Jerusalem, 1953.
340. LADELL, W.S.S. : Biology of man in arid zones - UNESCO and Zone Committee, 1955.
341. LADELL, W.S.S. : Myth about the tropics - UNESCO Courier, 1955.
342. LAKES, J.V. : The nocturnal heat balance - *Nature*, 176, 32, 1955.
343. LANDSBERG, H.E. : Eine Bett-Temperatur Studie - *Bioklim. Beibl.*, 5, 2, 66-68, 1938.
344. LEBLANC, J.A. : Effet de l'acide ascorbique sur plusieurs variables biologiques normalement affectées par le froid - *Can.J. Med. Sci.*, 27, 349, 1949.
345. LEBLANC, J.A. : Physiological response of men to cold - DRB Symposium, 1951.
346. LEBLANC, J.A. : Subcutaneous fat and skin temperature - *Can.J. Biochem. and Physiol.* 32, 354, 1954.
347. LEBLANC, J.A. : Studies on acclimatization and on the effect of ascorbic acid in men exposed to cold - *Can.J. Biochem. and Physiol.*, 32, 407, 1954.
348. LEBLANC, J.A. : Physiological evaluation of clothing and equipment at Fort Churchill in Winter 1954-1955. - DRB Symposium, 1955.
349. LEBLANC, J.A. : Blood ascorbic level of a northern population - DRNL report, 3/55, 1955.
350. LEBLANC, J.A. : Insulation of sleeping bags and pads - DRNL Technical Memorandum 3/55, 1955.
351. LEBLANC, J.A. : Efficiency of arctic clothing in providing thermal comfort - DRNL Technical Memo 8/55, 1955.
352. LEBLOND, C.P. and GROSS, J.: Effect of thyroidectomy on resistance to low environmental temperature - *Endocrinology*, 33, 155-160, 1943.
353. LEBLOND, C.P., DESMARAIS, A. and DUGAL, C.P.: Effet de l'ablation partielle du foie sur la résistance au froid et à la chaleur - *Rev.Can. de Biol.*, 2, 332, 1943.
354. LEBLOND, C.P., DESMARAIS, A. and DUGAL, L.P.: Effet de l'ablation du foie sur la résistance au froid d'animaux adaptés - *Rev. Can. de Biol.*, 2, 1943.

55. LEBLOND, C.P., DUGAL, L.P. and THERIEN, J.: Les aliments choisis par le rat blanc au froid et à la chaleur - Rev.Can. de Biol., 3, 127, 1944.
56. LEBLOND, C.P., GREGOIRE, F. and ROBILLARD, E.: Effect of partial hepatectomy on general resistance and blood sugar level in rats exposed to anoxia - J. of Av. Med., 15, 158, 1944.
57. LEBLOND, C.P., GROSS, J., PEACOCK, W. and EVANS, R.D.: Metabolism of radi-iodine in the thyroids of rats exposed to high or low temperatures - Am.J.Physiol., 140, 671, 1944.
58. LEBLOND, C.P., DOUGHERTY, J. and GROSS, J.: Steady state of the thyroidal iodine - Endoc., 48, 700-713, 1951.
59. LEE, D.H.K. : The human organism and hot environments - Trans. Roy. Soc. Trop.Med. & Hyg., 29, 7, 1935.
60. LEE, D.H.K. : The regulation of cardiac rate during exposure to heat - Jl.of Physiol., 85, 39, 1935.
61. LEE, D.H.K. : Some immediate effects of reduced cooling powers on human subjects - J. Physiol., 279, 1935.
62. LEE, D.H.K. : Some immediate effects of reduced cooling powers upon the water balance and related effects in the human subject - J. Physiol., 410, 1935.
63. LEE, D.H.K. : The effect of water intake upon human reactions to reduced cooling powers - Jl. Physiol., 86, 204, 1936.
64. LEE, D.H.K. : Physiology and the tropics.- Malayan Med. J., 11, 105, 1936.
65. LEE, D.H.K. : The settlement of tropical Australia - Med.J. Aust., 11, 707, 1936.
66. LEE, D.H.K. : Biochemical factors in adaptation - Aust.Chem.Inst.J.Proc., 4, 426, 1937.
67. LEE, D.H.K. : The effect of exercise in hot atmospheres upon the pulse rate - Med. J. Aust., 11, 664, 1940.
68. LEE, D.H.K. : Basis for the study of Man's reaction to tropical climates - Univ. Qld. Papers Dept. Physiol., 1, 5, 1940.
69. LEE, D.H.K. : The effect of exercise in hot atmospheres upon the salt water balance of human subjects - Med.J. Aust., 11, 249, 1941.
70. LEE, D.H.K. : Field trials of methods of tank personnel cooling carried out in New Guinea - AFV 48, Nov. 1943.
71. LEE, D.H.K. : Tropical fatigue and warfare - For Meeting Aust. N.Z. Assoc.Adv.Sci., 1946.
72. LEE, D.H.K. : Human climatology and tropical settlement - John Thompson Lecture, Univ. Qld. Pub., 1946.
73. LEE, D.H.K. : Heat and cold - Ann. Revs. Physiol., 10, 365, 1948.
74. LEE, D.H.K. : Thermal assessment of the environment. A review of current and proposed methods - Heating, Piping and Air Cond., 1948.
75. LEE, D.H.K. : Interactions of tropical environments and human performance - Environmental Protection Section OQMG Rpt., 1949.
76. LEE, D.H.K. and LEMONS, H.: Clothing for global man - Geog. Rev., 39, 181, 1949.
77. LEE, D.H.K. : Physiology as a guide to combating tropical stress - Med.Eng.Jl. of Med., 243, 723-730, 1950.
78. LEE, D.H.K. : Studies on clothing for hot environments: Death Valley 1950; Part I. Experiments and results - Environmental Protection Section rep. 178, OQMG, 1951.
79. LEE, D.H.K. : Repeatability of experimental heat tolerance test and the influence of season - J. Anim. Sci., 12, 757, 1953.
80. LEE, D.H.K., MCDOWEL, R.E. and FOHRMAN, M.H.: The measurement of water evaporation from limited areas of a normal body surface - J. Anim.Sci., 13, 2, 405-416, 1954.
81. LEE, D.H.K., MCDOWEL, R.E., SCHEIN, M.W. and FOHRMAN, M.H.: A method for the analysis and comparative presentation of temperature and humidity regimes - J. Anim. Sc., 14, 1, 1955.
82. LEE, D.H.K. : Nomographic Assessment of Thermal Strain in man - Lecture, 31 Aug. 1956 at the Symposium on Bioclim. at UNESCO H.Q., Paris.
83. LEE, D.H.K. : Climate and economic development in the Tropics - Council on Foreign relation, Inc. New York 21, N.Y., Febr. 1957.
84. LEE, D.H.K. a.o. : Atmospheric conditions encountered during a survey of RAAF ground crew on tropical service and their relationship to both sweat loss and the results obtained with the Harvard Pack test - Army report : FLOR 9.
85. LEE, D.H.K. : Clothing tests - Interim Report AFV 8.
86. LEE, D.H.K. : Fatigue in jungle operations - Army report: FRC 2.
87. LEE, D.H.K. : The principles of clothing for hot environments - Army report: FLOM 2.

388. LEE, D.H.K. : Tropical climates and the soldier - Army Report: PRC 1.
389. LEE, D.H.K. : Studies of evaporation from an artificial "skin" underneath surface cloths - Army Report: FLOR 2.
390. LEE, D.H.K. : Psycho-motor efficiency under hot conditions - Army Report: FLSR 3.
391. LEE, D.H.K. : The effect upon AFV Personnel of operations in the humid tropics with especial reference to the M3 medium - Army Report: AFV 24a.
392. LEE, D.H.K. a.o. : Effect of tropical service upon some physical characteristics of RNZAF ground-crew - Army Report: FLSR 5.
393. LEE, D.H.K. : Field investigation into the incidence of tropical fatigue - Army Report: FLSR 1.
394. LEE, D.H.K. a.o. : Variations in human reactions to hot atmospheres as functions of temperature and humidity - Army report: FLOR 5.
395. LEE, D.H.K. : Atmospheric conditions in troop decks of H.M.A.T. "KATOOMBA" - Army Report: AFV 40.
396. LEE, D.H.K. a.o. : Effect of tropical service upon some physical characteristics of RAAF ground-crew - Army Report: FLSR 4.
397. LEE, T.S. : Observation on man exposed to excessive humid heat (Eng.)- Jap. J. Physiol., 2, 103, 1951.
398. LEE, T.S. : The standardisation of conditions in the comparative study of sweating under heat stress - Med. J. Malaya, 7, 90-99, 1952.
399. LEE, T.S. : Daily variations in physiological values for european and asian men working at high temperatures in the tropics - Rep. R.N.P. 53/749 to R.N. Personnel Research Committee, med. Res. Coun. London, 1953.
400. LEE, T.S. : Periodic and diurnal variations in physiological values for acclimatised naval ratings working at high temperatures for 5 months in the tropics - Rep. R.N.P. 53/750, R.N. Personnel Research Committee, Med. Res. Council, London, 1953.
401. LEE, T.S. : Activity of human sweat glands during exposure to cold - J. Physiol., 122, 59-65, 1953.
402. LEE, T.S. : Physiological gustatory sweating in a warm climate - J. Physiol., 124, 528-542, 1954.
403. LEHMANN, G. and SZAKALL, A.: Anpassungserscheinungen an Hitzearbeit - Arbeitsphysiologie 9, 678-705, 1937.
404. LEHMANN, G. : Hitzearbeit und Durst - Der Ruhrarbeiter, August 1939.
405. LEHMANN, G. and SZAKALL, A.: Schweißverlust und Getränkeaufnahme bei Bergleuten und Hitzearbeitern - Arbeitsphysiologie, 11, 73-100, 1940.
406. LEHMANN, G. : Schwitzen und Trinken bei Hitzearbeit - Zentralblatt für Arbeitswissenschaft und soziale Betriebspraxis, 4, 129-135 u. 149-153, 1950.
407. LEHMANN, G. : Arbeit bei hohen Temperaturen - Schriftenreihe der Arbeitsgemeinschaft für Forschung des Landes Nordrhein-Westfalen, Düsseldorf - Heft Westdeutscher Verlag Köln und Opladen, Opladen, 1956.
408. LEMAIRE : Les réactions thermo-régulatrices en milieu tropical - Gaz. Méd. de France, 58, 15, 933, 1951.
409. LEMAIRE : L'adaptation au climat africain - Ann. de l'Institut d'Hydrologie et Climatologie, 23, 72, 65, 1952.
410. LEMAIRE : Les troubles psychiques imputables à la déshydratation - Bull. Méd. d'A.O.F., 9, 1, 113, 1952; Le Méd. Aéronautique, 7, 3, 299, 1952.
411. LEMAIRE : Déshydratation et Albuminurie - Bull. Méd. d'A.O.F., 9, 1, 115, 1952 ; La Méd. Aéronautique, 7, 3, 299, 1952.
412. LEMAIRE : Les variations des cations alcalins du sang au cours des états de déshydratation mixte - Bull. de la Soc. de Chimie Biol., 35, 9, 1027, 1953.
413. LEMAIRE : Les états de déshydratation mixte - Bull. et Mém. de l'Ecole de Méd. Dakar, 1, 150, 1953.
414. LEMAIRE : Considérations sur l'étude physiologique des climats - Le Maroc Médical, 342, 1076, 1953.
415. LEMAIRE : Les réactions circulatoires aux climats tropicaux - La Médecine Aéronautique, 8, 3, 256, 1953.
416. LEMAIRE : Les conditions de l'adaptation aux pays tropicaux - La semaine des Hôpitaux, 64, 29, 3270, 1953.
417. LEMAIRE : Les variations des électrolytes sanguins au cours des états de déshydratation mixte - J. de Physiol., 46, 1, 418, 1954. 22-ème Congrès de l'association des Physiol. de Langue Française, Lille, 1954.
418. LEMAIRE : Les effets du climat tropical sur certaines réactions circulatoires - Bull. Médical de l'A.O.F., 2, 57, 1954.
419. LEMAIRE : La sudation de réhydratation - Bull Méd. de l'A.O.F., 2, 1, 135, 1954.
420. LEMAIRE : L'adaptation aux climats tropicaux du sujet transplanté - Bull. et Mémoire Ecole de Médecine de Dakar, 2, 180, 1954.

21. LEMAIRE : Le débit sudoral du sujet adapté au climat désertique - Bull. et Mémoire de l'Ecole de Médecine de Dakar, 2, 198, 1954.
22. LEMAIRE : La pratique des sports dans les climats tropicaux - Médecine Ed. Physique et Sports, 1, 3, 1954.
23. LEMAIRE : Les variations de la concentration des cations magnésium du sang au cours des états de déshydratation mixte - C.R. Soc. Biol., Nancy, 148, 1-2, 144, 1954.
24. LEMAIRE : La part des réactions neuro-hormonales dans l'acclimatement aux pays tropicaux - La Presse Thermale et Climatologique, 76, 3-4, 1954.
25. LEMAIRE : Influence du facteur racial sur l'adaptation au climat saharien - Bull. de l'Institut français d'Afrique Noire, 17, 4, 1221, 1955.
26. LEMAIRE : Les syndrômes d'inadaptation primitive aux climats tropicaux - Comm. Congrès Aéro Médical Assoc. Paris 1955. Med. Aéronautique, 10, 4, 465 et 467, 1955.
27. LEMAIRE : Les échanges hydriques au cours des états de déshydratation mixte - 23ème Congrès de l'association des Physiologistes de Langue Française Gand, Avril 1955 and: J. Physiol., 47, 218-220, 1955.
28. LEMAIRE : Conduite à tenir dans les états de déshydratation mixte - Méd. Aéronautique, 10, 4, 469-473, 1955.
29. LEMAIRE : Influence du facteur ethnique sur la sécrétion sudorale en climat tropical - La Presse Thermale et Climatologique, 152, 5-6, 1955.
30. MCFARLANE, W.V. : Effects of temperature and surgery upon the elimination of radioiodine in the rat - J. Clin. Endo. and Met., 1952.
31. MCFARLANE, W.V. and ROBINSON, K.W.: The influence of environmental temperature on the level of plasma antidiuretic substances in the rat - Aust.J. Biol.Sci., 9, 130, 1955.
32. MCFARLANE, W.V., ROBINSON, K. and HOWARD, B.: Observations on seasonal fluctuations of 17 ketosteroid output in the subtropics and tropics - Med.J. Aust., 1, 756, 1955.
33. MCFARLANE, W.V. and MEARES, J.: Modification of the after potential of single muscle fibres by 2:4 DNP - Nature, 176, 403-404, 1955.
34. MCFARLANE, W.V. : (a) Human functions in the tropics; (b) The water economy and heat tolerance of tropical Merinos - Symposium Man and Animals in the tropics. Aust. Acad. Sci., 1956.
35. MCFARLANE, W.V. : Urinary sodium-potassium ration in hot environments - Med.J. Aust., 2, 130, 1956.
36. MCFARLANE, W.V. and ROBINSON, K.: Seasonal changes in amount of plasma antidiuretic activity released by a standard heat stimulus - J. Physiol., 135, 1, 1957.
37. MCFARLANE, W.V., PENNY CUIK, P.R. and THRIFT, E.: Resorption and loss of foetuses in rats living at 35°C. - J. Physiol., 135, 1957.
38. MEFFERD Jr., R.B., HALE, H.B. and MARTENS, H.H.: Nitrogen and electrolyte excretion of rats chronically exposed to adverse environments - Am.J. Physiol., 192, 209-213, 1958.
39. MEFFERD Jr., R.B., and HALE, H.B.: Effects of abrupt temperature changes on excretion characteristics of rats acclimated to cold, neutral, or hot environments - Am.J. Physiol., 195, 726-734, 1958.
40. MOLFINO, F. : Alterazioni aortiche e lavoro in ambienti a temperatura elevata - Minerva Medica, 24, 1934.
41. MONGE, C. : Physiological adaptations of dwellers in the tropics - Proc. of the IV Intern. Congr. on Tropical Med. and Malariae, Washington U.S. Government. Print. Off., 1, 136-147, 1948.
42. MONGE, C. : Acclimatization in the Andes. Historical confirmations of climatic aggression in the development of Andean Man - Baltimore, Johns Hopkins Press, 1948.
43. MONGE, C. : Physiological anthropology of the Dwellers in America's High Plateaus - Proc. of the 29th Intern. Congr. of Americanists, Chicago, The Univ. Chicago Press., 3, 361-373, 1952.
44. MORIKOFER, W. : Einiges über die Abkühlungsgrösse. Messung der Abkühlungsgrösse - Aus dem Sammelwerke Prof. Mladejovsky, 15, 334-349, 1935.
45. MUELLER, T. : Klimakammeruntersuchung zur Höhenfliegertauglichkeitsbestimmung - Wiener.Med. Wschr., 50, 1033, 1955.
46. OGATA, K. : The effect of the application of cold on the metabolism at high surrounding temperatures - J. Orient Med., 22, 6, 1173, 1935.
47. OGATA, K. : Functional variations in human sweat glands, with remarks upon the regional difference of the amount of sweat - J. Orient Med. 23, 6, 1155, 1935.

448. OGATA, K. et al. : Observations on the daily amount of human perspiration in the winter and summer, and its relationship with the intake of food and drink and with the amount of urine - J. Orient. Med., 23, 1, 13, 1935.
449. OGATA, M. et al. : Racial variations in the sweat secretion on the axilla - J. Orient. Med., 25, 6, 1613, 1936.
450. OGATA, K. et al. : Influence of the exposure to cold on the regulation of human temperature - J. Orient. Med. 36, 2, 369, 1942.
451. OGATA, M. : Body temperature regulation in human under cold environment - J. Orient. Med., 37, 3, 647, 1942.
452. OGATA, K. : Climate and body temperature regulation - Kagaku Chishiki, 2818, 12, 1948.
453. OGATA, K. : Acclimatization of regulatory function of body temperature regulation - Rinsho to Kenkyu, 26, 1, 1949 (Jap.).
454. OGATA, K. : Acclimatization of the body temperature regulation - Rinsho to Kenkyu, 26, 2, 57, 1949.
455. OGATA, K. : Regulation of body temperature under cold environment - Nanjo Publ. Co., Tokyo, 1949.
456. OGATA, K. et al. : Topographic determination of the human surface temperatures in every season - Bull. Res. Inst. Diath. Med., 2, 1, 57, 1951.
457. OGATA, K. : Significance of thermal sensation and thermal gradients in relation to regulation of body temperature - Seitai no Kagaku, 3, 3, 128, 1951.
458. OGATA, K. et al. : Influence of a large amount of sodium chloride ingestion on the basal metabolism and on resistance to cold and frost-bite - Jap. J. Physiol., 2, 4, 303, 1952.
459. OGATA, K. et al. : Responses to cold observed in regard to physical build and native place - J. Orient. Med. 39, 6, 1021, 1953.
460. OGATA, K. : Regulation of the body temperature in childhood - J. Physiol. Soc. Jap. 15, 175, 1953.
461. PEEL : Thermal conditions in traditional mud dwellings in Northern Nigeria - Journal of Tropical Medicine & Hygiene, 61, 8, 189-204, 1958.
462. PETR, B. : Attempts to determine the comfort zone of the human organism in relation to meteorological factors - Cs. HEMI, IV/2, 1955.
463. PETR, B. : Biometeorology in Hygiene - Cs. HEMI IV/9, 1955.
464. PETR, B. : The influence of refrigeration upon skin temperature and total volume of expired air on human organism - Ceskoslovenska hygiena, 1/6, 1956.
465. PETR, B. : Communal Hygiene - Praha, 1956.
466. PFLEIDERER, H. : Wärme- und Wasserverlust des Gesunden im Zimmer und im Freien - Verh. d. Ges. inn. Med., Wiesbaden, 372, 1933.
467. PFLEIDERER, H. : Studien über den Wärmehaushalt des Menschen - Z. exp. Med. 90, 245, 1933.
468. PFLEIDERER, H. : Meteorische Einflüsse auf die Temperatur der Expirationsluft - Tagungsbericht. Ged. D. Naturforscher u. Aerzte, Hannover, 158, 1934.
469. PFLEIDERER, H. and BUETTNER, K.: Die physiologischen und physikalischen Grundlagen der Hautthermometrie - J.A. Barth, Leipzig, 1935.
470. PFLEIDERER, H. : Meteorophysiologie des Wärmehaushaltes - Verh. D. Ges. Inn. Med., 47. Kongress Wiesbaden, 492. 1935.
471. PFLEIDERER, H. : Haut und Klima - D. Med. Wschr., 635, 1937.
472. PFLEIDERER, H. and BUETTNER, K.: Methodik der thermoelektrischen Hauttemperaturmessung - Abderhaldens Handbuch der biologischen Arbeitsmethoden, Abtlg. IV, 16, 1937.
473. PFLEIDERER, H. : Bioklimatologie unter besonderer Berücksichtigung der Haut - Strahlentherapie 66, 627, 1939 and Arch. f. Dermatol. u. Syphilis, 180, 251, 1940.
474. PFLEIDERER, H. : Die Bedeutung der Trifrigorigraphie für die bioklimatische Forschung - Bioklim. Beibl., 121, 1943.
475. PFLEIDERER, H. : Zur Bioklimatik des menschlichen Wärmehaushaltes - Ber. Dtsch. Wetterdienstes i.d. US-Zone, 42, 338-341, 1952. (Knoch-Festschrift).
476. PFLEIDERER, H. : Kritische Betrachtungen über die Abkühlungsgrösse - Ber. Dtsch. Wetterdienstes i.d. US-Zone, 38, 267-270, 1953 (Weickmann-Festschrift).
477. PFLEIDERER, H. : Biologische und therapeutische Infrarotprobleme - Vortrag 62. Kongr. der Dtsch. Ges. für Balneologie, Bioklimatologie und Physikalische Therapie, Bad Nauheim, 1954. Arch. Physik. Therapie, 7, 201-205, 1955.
478. PFLEIDERER, H. : Extrarenaler Wasserhaushalt und dessen Abhängigkeit von klimatischen Bedingungen - Vortrag 63. Kongress der Dtsch. Ges. f. Balneologie, Bioklim. & Phys. Ther., Lugano, 1955; Z. angew. Bäder- u. Klimaheilk., 1, 71-86, 1956.
479. PFLEIDERER, H. : Zum Begriff der Abhärtung - Med. Klin. 51, 1574, 1956.

30. PICHOTKA, J. : Ueber die histologischen Veränderungen der Lunge nach Atmung von hochkonzentriertem Sauerstoff im Experiment - Diss.Freiburg, 1941 ; Beitr. path. Anat., 105, 381, 1944.
31. PICHOTKA, J. and BECKER, FREYSENG, H.: Der Kohlensäurehaushalt bei Sauerstoffüberdruck - Klin. Wschr., 1, 339, 1944.
32. PICHOTKA, J. and KUEHN, H.A.: Untersuchungen über die langfristige Einwirkung einer hochprozentigen Sauerstoffatmosphäre bei vermindertem Gesamtdruck - Arch. exper. Path.u. Pharmakol., 205, 659, 1948.
33. PICHOTKA, J. and LEWIS, R.B.: Use of Heparin in Treatment of Experimental Frostbite - Proc. of the Soc. for Experimental Biol. and Med., 72, 130-136, 1949.
34. PICHOTKA, J. and REICHEL, H.: Determination of the normal clotting time and its variations determined with a modified capillary method - Americ.Jul. Physiol., 162, 632, 1950.
35. PICHOTKA, J., LEWIS, R.B. and FREYTAG, E.: Sequence of increasing cold injury - Texas Reports on Biol. and Med., 9, 613, 1951.
36. PICHOTKA, J. and LEWIS, R.B.: Effect of rapid and prolonged rewarming on local cold injury - U.S. Armed Forces M.J., 2, 1293, 1951.
37. PICHOTKA, J. : Ueber den Wirkungsmechanismus der Antikoagulantien - Ver.Dtsch. Ges. Inn. Med., 58. Kongress 1952.
38. PICHOTKA, J. : Das Verhalten der Schilddrüse und der Körpertemperatur bei der Adaptation an niedere Umgebungstemperaturen - Arch.exper.Path.u.Pharmakol., 215, 299, 1952.
39. PICHOTKA, J. : Das Verhalten der Schilddrüse bei Versagen der Wärmeregulation nach protahierter Belastung - Arch.exper. Path. u, Pharmakol., 215, 317, 1952.
40. PICHOTKA, J. : Das Verhalten der Schilddrüse bei akuter Insuffizienz der Wärmeregulation - Arch. exper. Path.u. Pharmakol., 216, 268, 1952.
41. PICHOTKA, J. : Ueber den Durchtritt von Gallensubstanzen aus dem Gallengangssystem ins Leberparenchym bei Sauerstoffmangel - Klin.Wschr., 143, 1952.
42. PICHOTKA, J. : Experimentelle Untersuchungen über die Heparinwirkung - Klin. Wschr., 189, 1952.
43. PICHOTKA, J. and CAROLI, G.: Beeinflussung der normalen Blutgerinnungszeit durch das Wetter - Naturwissenschaften, 39, 193, 1952.
44. PICHOTKA, J. : Untersuchungen über Gefrierpunkte des lebenden Gewebes - Z. Biol., 105, 181, 1952.
45. PICHOTKA, J. : Die Morphogenese der Schilddrüsenveränderungen während der Anpassung an niedere Umgebungstemperaturen - Beitr. path. Anat., 113, 169, 1953.
46. PICHOTKA, J., KUGELGEN, B.von and DAMMANN, R.: Die Bedeutung der Schilddrüse für die Temperaturregulation - Arch.exper.Path.u. Pharmakol., 220, 398, 1953.
47. PICHOTKA, J. : Tierexperimentelle Untersuchungen zur Auswertung und Anwendung von Antikoagulantien - Arch.exper.Path.u. Pharmakol., 222, 159, 1954.
48. PICHOTKA, J. and CAROLI, G.: Weitere Untersuchungen zur Beziehung zwischen Blutgerinnung und Wetter - Arch. Meteorol., Geophys. u. Bioklimat., 5, 403, 1954.
49. PICHOTKA, J. and MAYER, K.: Experimentelle Untersuchungen über die perkutane Wirksamkeit gerinnungshemmender Substanzen - Arzneimittelforschung, 4, 277, 1954.
50. PICHOTKA, J., CREUTZFELDT, O. and HOEFLE, W.: Das Verhalten der Körpertemperatur im Sauerstoffmangel - Arch. exper. Path. u. Pharmakol., 225, 314-322, 1955.
51. PICHOTKA, J., CREUTZFELDT, O. and HOEFLE, W.: Die Bewegungen von Körpertemperatur und Sauerstoffaufnahme beim Wechsel zwischen normalen und erniedrigten Sauerstoffspannungen - Arch. exper. Path.u. Pharmakol., 225, 323-334, 1955.
52. PICHOTKA, J., CREUTZFELDT, O. and HOEFLE, W.: Die Bewegungen der Körpertemperatur und der Sauerstoffaufnahme bei veränderlicher Wärmeabgabe im Sauerstoffmangel - Arch. exper. Path. u. Pharmakol., 225, 335-341, 1955.
53. PICHOTKA, J. : Das physiologische Experiment in der Medizin-Meteorologischen Forschung - Hauptvortrag auf der Medizin-Met. Tagung in Hamburg 1955 ; Med.-Met. Hefte, 11, 54, 1956.
54. PICHOTKA, J. : Ueber die Beziehung von Körpertemperatur und Sauerstoffaufnahme im Sauerstoffmangel - Internat. Physiologischer Kongress, Brüssel, 1956.
55. PICHOTKA, J. : Der Gesamtorganismus im Sauerstoffmangel - Handbuch d. allgem.Pathologie, 4, 2, 497-568, 1957.
56. PICHOTKA, J. : Ist die O_2 -Aufnahme spannungsabhängig ? - Klin.Wschr., 649, 1957.
57. PIRLET, K. : Individualpsychologische Studien des Wärmehaushaltes - Vortrag 63. Kongr. der Dtsch. Ges. für Balneologie, Bioklim. & Phys. Ther., Freudenstadt 1955 - Arch. Physik. Therapie, 8, 162-169, 1956.
58. PITESKY, I. and LAST, J.H.: Effects of seasonal heat stress on glomerular and tubular functions in the dog - Am.J. Physiol., 164, 497-501, 1951.

509. RADIGAN, L.R. and ROBINSON, S.: Effects of environmental heat stress and exercise on renal blood flow and filtration rate - J. Appl. Physiol., 2, 185-191, 1949.
510. RALFS, R. : Untersuchungen über die Perspirationsgrösse von Abgehärteten und Verweichtlichten - Diss.med. Kiel, 1948.
511. RALLI, E.P., KUHL Jr, W.J., GERSHBERG, H., BECK, E.M., STREET, E.R. and LAKEN, B.: Effects of vitamin supplementation of the diet on reaction to short - term cold stress in normal young male adults - Metabolism, 5, 170-196, 1956.
512. REIFFERSCHIED, H. : Prüfung der Wärmehaltfähigkeit einiger natürlicher und künstlicher Faserstoffe - Vortrag Int. Kongress Angew. Met., Obertraun, 1955.
513. REIFFERSCHIED, H. : Untersuchung der Wärmehaltfähigkeit einiger natürlicher und künstlicher Faserstoffe mit Hilfe des DAVOSer Frigorimeters - Vortrag Bädertag Freudenstadt-Lugano, 1955.
514. REIFFERSCHIED, H. : Untersuchungen der Wärmehaltfähigkeit einiger natürlicher und künstlicher Faserstoffe mit Hilfe des Davoser Frigorimeters - Archiv. f. Physikalische Therapie, 8, 3, 1956.
515. REIFFERSCHIED, H. : Messwerte und theoretische Überlegungen zum Wärmedurchgang durch Gewebe aus natürlichen und synthetischen Fasern - Melland Textilberichte, Heidelberg, 38, 2, 189-191, 1957 and: Ber. des Met. Geophys. Inst. zu Frankfurt a.M., 6, 1956.
516. RENBOURN, E.T. : Observations on normal body temperature in North India - Brit. Med. J., 2, 99, 1946.
517. RENBOURN, E.T. : Variations diurnal and over longer periods of time in blood etc. - J. Hyg., 45, 455, 1947.
518. RIEDEL, G. and MEITNER, H.J.: Veränderung des Stoffwechsels bei Klimabelastung - Klin. Wschr., 44/45, 677, 1943.
519. RIEDEL, G. : Klimatologisch-physiologische Studie über die Grenzen der Hitzeerträglichkeit - Berichte des Deutschen Wetterdienstes in der US-Zone, 38, 273-277, 1952.
520. RIEDEL, G. : Wärmeabgabesonde (Diskussionsbeitrag)- VDI-Berichte, 21, 27-28, 1957.
521. ROBINSON, K.W., HOWARD, B. and MACFARLANE, W.V.: Observations on seasonal fluctuations in 17-ketosteroid output in the subtropics and tropics - Med. J. Aust., 2, 756-759, 1955.
522. SANCETTA et al : Effects of "dry" heat on the circulation of man. General hemodynamics - Am. Heart Journal, 56, 2, 212-222, 1958.
523. SAPIN-JALOUSTRE, J. : L'établissement de la thermo-régulation chez le Manchot Adélie - C.R. Acad. Sci, 1443-1445, 1953.
524. SAPIN-JALOUSTRE, J. : Contribution à l'étude de l'acclimatation de l'homme et des vertèbres supérieurs dans l'Antarctique et plus spécialement en Terre Adélie - La Biologie Médicale, 44, 1, 15-72 and 2, 133-208, 1955.
525. SAPIN-JALOUSTRE, J. : Enquête sur les Gelures - Hermann, Paris, 1, 1956.
526. SARGENT, F., Robinson, P. and JOHNSON, R.E.: Water soluble vitamins in sweat - J.Biol.Chem., 153, 285, 1944.
527. SARGENT, F. : Acclimatization in the Andes. Historical confirmation of "Climatic Aggression" in the development of Andean man - Bull. Am.Meteor. Soc., 30, 298, 1949. Book review.
528. SARGENT, F. : Role of the field test in nutrition and weather stress studies. - In nutrition under climatic stress. A symposium. (H. Spector and M.S. Peterson, editors). National Academy of Sciences. National Research Council, Washington, 130-156, 1954.
529. SARGENT, F. : Chronic dehydration as a source of error in estimating body fat by skinfold thickness method - Fed. Proc. 13, 126, 1954.
530. SARGENT, F. and JOHNSON, R.E.: Solute load and the renal parameters of chronically dehydrated men - J. Clin. Investigation, 33, 962, 1954 (Abstract).
531. SARGENT, F. and FLEISCHNER, J.R.: Effects of heat on the rat: a study of crossacclimatization - The American Journal of Physiology, 187, 3, 1956.
532. SCHAEFER, H.H. : Untersuchungen über den Einfluss von Abhärtungsmaßnahmen auf die Hauttemperatur - Diss.Med.Kiel, 1948.
533. SCHULZ, A. : Thermische Einflüsse auf die Erythem- und Pigmentbildung - Vortrag Lichttagung Westerland 1950. Strahlentherapie 86, 142, 1952.
534. SELLERS, E.A., YOU, S.S. and THOMAS, N.: Acclimatization and survival of rats in the cold: Effects of clipping, of adrenalectomy and of thyroidectomy - Am. J. Physiol., 165, 481-485, 1951.
535. STEIN, H.J., ELIOT, J.W. and BADER, R.A.: Physiological reactions to cold and their effects on the retention of acclimatization to heat - J. Appl. Physiol., 1, 575-585, 1949.
536. STRUZKA, V. : Temperature feeling and the cooling-power especially - Bisklimatische Beiblätter, 1, 1943 (Czech.).

537. TALBOTT, J.H. : Heat Cramps: A clinical and chemical study - J. Clin. Invest., 12, 3, 1933.
538. TALBOTT, J.H. : Heat cramps - Ann. Int. Med., 10, 270, 1936.
539. TALBOTT, J.H., DILL, D.B., EDWARDS, H.T., STUMME, E.H. and CONSOLAZIO, W.V.: The ill effects of heat upon workmen - J. Industr. Hygiene and Toxicology, 19, 258, 1937.
540. TALBOTT, J.H. : Ill effects of heat - Modern Med. Therapie in Gen. Practice. Williams and Wilkins Co., 1940.
541. TALBOTT, J.H. : The physiologic and therapeutic effects of hypothermia - New. Eng. J. Med., 224, 281, 1941.
542. TALBOTT, J.H., CONSOLAZIO, W.V. and PECORA, L.J.: Hypothermia - Arch. of Int. Med., 68, 1120, 1941.
543. TALBOTT, J.H. and TILLOTSON, K.J.: The effects of cold on mental disorders - Dis. of Nerv. System, 2, 1, 1941.
544. TALBOTT, J.H. : The effect of cold - Oxford System of Medicine, 1948.
545. TALBOTT, J.H. : Cold: Pathologic effects - Med. Physics, 2, 1950.
546. TALBOTT, J.H. : Heat: Pathologic effects - Med. Physics, 2, 1950.
547. TALBOTT, J.H. : Frostbite in Korea 1950-51 - Trans Assoc. Amer. Phys., 65, 316, 1952.
548. TALBOTT, J.H. : Cold Injury. - Transaction of the first Conference June 4-5, 1951 ; Second conference November 20-21, 1952; Third conference, February 22-25, 1954. Josiah Macy Jr. Foundation.
549. TREICHLER, R. and MITCHELL, H.H.: The influence of plane of nutrition and of environmental temperature on the relationship between basal metabolism and endogenous nitrogen metabolism subsequently determined - J. Nutrition, 22, 333-343, 1941.
550. VAUGHAN, D.A. and VAUGHAN, L.N.: Effect of a low environmental temperature on weight and food consumption of thiamine-deficient rats - Technical Report, 58-7, Alaskan Air Command, Arctic Aeromedical Labor., June 1958.
551. VAUGHAN, D.A., HANNON, J.P. and VAUGHAN, L.N.: Arctic Survival Rations. IV. Associated effects of diet, environmental temperature, and duration of exposure on the major constituents of the livers of rats - Alaskan Air Command, Arctic Aeromed. Lab., Ladd Air Force Base, Techn. Report 59, 5, 1959.
552. VAUGHAN, D.A. and VAUGHAN, L.N.: Thiamine load test studies in humans - Alaskan Air Command, Arctic Aeromed. Lab., Ladd Air Force Base, Techn. Report, 57, 58, 1959.
553. VERZAR, F. : Dauer-Akklimatisation an grosse Höhen - Bull. d. Schweiz. Akad. der Med. Wissensch., 7, 1, 26-38, 1951.
554. VERZAR, F. and FLOEKIGER, E.: Lack of adaptation to low oxygen pressure in aged animals - Journal of Gerontology, 10, 3, 1955.
555. WARMBT, W. : Akrale Hauttemperatur (Finger) und Klimaempfinden in ihrer Beziehung zu meteorologischen Faktoren - Abh. d. meteor. u. hydrologischen Dienstes d. D.D.R., 6, 41, 1956.
556. WARMBT, W. : Akrale Hauttemperatur (Finger) und Klimaempfinden, gleichzeitige Messungen an einem kälte- und wärmeempfindlichen Reaktionstyp - Zschr. f. Angew. Met., 2, 11, 323-334, 1957.
557. WEINER, J.S. and HUTCHINSON, J.C.D.: Hot humid environment: Its effect on the performance of a motor co-ordination test - Brit. J. Indus. Med., 2, 154, 1945.
558. WEINER, J.S. : The regional distribution of sweating - J. Physiol., 104, 32, 1945.
559. WEINER, J.S. : Renal function of men working in hot environments - XVIIth International Physiological Congress 1947, 124 (Abstracts).
560. WEINER, J.S. : The biological assessment of clothing for tropical service use - Brit. med. Bull., 5, 1016, 1947.
561. WEINER, J.S. and HARRISON, R.G.: Abdomino-testicular temperature gradients - J. Physiol., 107, 48p, 1948.
562. WEINER, J.S. : The physiological study of the effects of high environmental temperatures - Royal Soc. Empire Scient. Conf., report, 1, 442, 1948.
563. WEINER, J.S. and HARRISON, R.G.: Vascular patterns of mammalian testis and their functional significance - J. Exp. Biol., 26, 304, 1949.
564. WEINER, J.S. and HEYNINGEN, R. van: Lactic acid and sweat gland function - Nature, 164, 1949.
565. WEINER, J.S. : Observations on the working ability of Bantu mine workers with reference to acclimatisation to hot humid conditions - Proc. 4th Empire Mining and metal. Congress 1950. Brit. Med. Industr. Med., 7, 17.
566. WEINER, J.S. : Physiological studies of the effects of heat - London H.M.S.O., 1951.
567. WEINER, J.S. : A spectrophotometer for skin colour measurement - Man, 253, 1952.
568. WEINER, J.S. and HEYNINGEN, R.E. van: Salt losses of men working in hot environments - Brit. J. Industr. Med., 9, 56, 1952.

569. WEINER, J.S. and HEYNINGEN, R.E. van: Relation of skin temperature to salt concentration of general body sweat - J. Appl. Physiol., 4, 725, 1952.
570. WEINER, J.S. and HEYNINGEN, R.E. van: Observations on the lactate content of sweat - J. appl. Physiol., 4, 734, 1952.
571. WEINER, J.S. and HEYNINGEN, R.E. van: The effect of arterial occlusion on sweat composition - J. Physiol., 116, 395, 1952.
572. WEINER, J.S. and HEYNINGEN, R.E. van: A comparison of arm bag sweat and body sweat - J. Physiol., 116, 395, 1952.
573. WEINER, J.S. : Human adaptability to hot conditions of deserts 1953. - Symposium on the biology and productivity of hot and cold deserts. UNESCO and Institute of Biology.
574. WEINER, J.S. and HELLMAN, K.: Anit-diuretic substance in urine following exposure to high temperatures. J. appl. Physiol., 6, 194, 1953.
575. WEINER, J.S. : Man's adaptation to climate - The Times Science Review, 1954.
576. WEINER, J.S. : Nose shape and climate - Am.J. Phys.Anth., 12, 4, 1954.
577. WEINER, J.S., GRAY, C.H., HELLMAN, K. and LUNNON, B.: Excretion of adreno-cortical steroids in men under heat stress - J. Endocrinology, 10, 18, 1954.
578. WEINER, J.S. and LIND, A.R.: Working capacity in hot and humid conditions - The Manager, 23, 853, 1955.
579. WEINER, J.S., LIND, A.R., HELLMAN, R.F. and JONES, R.M.: Tolerance of men to work in hot saturated environments with reference to Mines Rescue Operations - Brit.J.Industry Med., 12, 296-303, 1955.
580. WEINER, J.S. : The diuretic response of Men working in hot and humid conditions - J. Physiol., 103, 36p, 1955.
581. WEISS, A.K. : Adaptation of rats to cold air and effects in tissue oxygen consumptions - Am.J.Physiol., 177, 201, 1954.
582. WEISS, A.K. and MOSS, W.G.: Effects of cold exposure on liver regeneration in the rat - Fed. Proc., 14, 104, 1955.
583. WEISS, A.K. and MOSS, W.G.: Effects of cold-exposure on liver oxygen consumption of subtotally hepatectomized rats - Fed.Proc., 14, 161, 1955.
584. WEISS, A.K. : Effects of cold-exposure on tissue oxygen consumptions of subtotally hepatectomized rats - Abstracts of communications, XX International Physiological Congress, Brussels, 30 July- 4 August 1956.
585. WEISS, A.K. : Tissue Oxygen consumption of rats adapted to cold - Fed. Proc., 15, 197, 1956.
586. WEISS, A.K. : Tissue responses in the cold-exposed rat - Am.J.Physiol., March 1957.
587. WEISS, A.K. and MOSS, W.G.: Effects of subtotal Hepatectomy on the rat's response to cold-exposure - J. Appl.Physiol., Jan. 1957.
588. WHITTOW, G.C. : The effect of different environmental and cold-bath temperatures on the cold pressor response in man - Proc.of the Physiological Soc., 15 - 16 July 1955. Journal of Physiology, 129, 1955.
589. WHITTOW, G.C. : Effect of Anti-histamine substances on cold vasodilatation in the finger - Nature, 176, 511-512, 1955.
590. WHITTOW, G.C. : Renal excretion of water and salt by healthy people in Singapore - Med.J. Malaya, 11, 126-133, 1956.
591. WILBER, C.G. : Water requirements of man - Med.Jnl. USAF, 8,8, 1121-1130.
592. WILL, E. : Die Perspiratio insensibilis in ihrem Verhalten zur Hauttemperatur - Diss.med. Kiel, 1950.
593. YOSHIMURA, H. : A new method of treatment of frost bite - Jap.Soc.of Snow and Ice, 11, 110-112, 1949.
594. YOSHIMURA, H. and IIDA, T.: Studies on the reactivity of skin vessels to extreme cold. I. A point test of the resistance to frost bite - Jap.J. of Physiol., 1, 147-159, 1950.
595. YOSHIMURA, H. and IIDA, T.: Studies on the reactivity of skin vessels to extreme cold. II. Factors governing the individual difference of reactivity, or the resistance to frost bite - Jap.J. of Physiol., 2, 177-185, 1952.
596. YOSHIMURA, H., IIDA, T. and KOISHI, H.: Studies on the reactivity of skin vessels to extreme cold. III. The effect of diets on the reactivity of skin vessels to cold - Jap.J. of Physiol., 2, 310-315, 1952.
597. YOSHIMURA, H. : Studies on the method of determination of temperature distribution of the human skin - J.Physiol.Soc.of Japan, 14, 517-524, 1952.
598. YOSHIMURA, H. et al.: Studies on the seasonal correlation between the metabolism and the thermal regulatory function. I. Studies on the seasonal variations of water and salts metabolism of the human body - J.Physiol. Soc.of Japan, 15, 47-57, 1953.

HUMAN BIOCLIMATOLOGY
Section E: World literature *

1. BARTNICKI, L., WIERZBICKI, Z.: A climatic outline of the Nowy Sącz valley and Krynica-Spa - Wiadomości Służby Hydrologicznej i Meteorolog, Warszawa, VI, 3, 1958.
2. BEMBROWSKI, B., HADUCH, S. : The problem of protecting the human organism against the harmful action of the ionised radiation - Lekarz Wojskowy, 3, 1957.
3. BOGUSZEWSKA, M. : Investigation of some reactions of the human organism working in cool environments - Prace Centralnego Instytutu Ochrony Pracy, 2, 24, 1958.
4. CHEŁCHOWSKI, W. : Measurements of cooling power by the graduation works of Ciechocinek Spa - Poznań, Wiadomości Uzdrowiskowe 3, 1958.
5. CHEŁCHOWSKI, W. : The intensity of solar radiation in Krynica-Spa (Carpathian Mountains) - Wiadomości Służby Hydrolog.- Meteorolog., Warszawa, VI, 5, 1958.
6. CHOBOT, H., TYCZKA, S. : The test of valuation of the spa-treatment of children in Rymanów-Zdrój (Carpathian Mountains)- Wiadomości Uzdrowiskowe, Poznań, 4, 1958.
7. CHOMICZ, K. : Atmospheric precipitation in Zakopane (Tatra Mountains) - Wiadomości Służby Hydrol.-Meteor., Warszawa, VI, 5, 1958.
8. GURBA, A. : The results of measurements of cooling at Zakopane (Tatra Mountains) - Wiadomości Służby Hydrolog.- Meteorolog., Warszawa, VI, 5, 1958.
9. KARASINSKI, T. : Atmospheric precipitation in the Tatra Mountains - Wiad. Służby Hydrol.- Meteorolog., Warszawa, VI, 5, 1958.
10. KĘPSKI, J. : The project of a regional spa for High Silesia industrial district - Biuletyn Służby Sanitarno- Epidemiologicznej Województwa Katowickiego, Katowice, 4, 1958.
11. KĘPSKI, J. : Health-Service indications for the programme concerning the use of the regional-spa at Ustroń-Wisła - Biuletyn Stacji Sanitarno- Epidemiologicznej województwa katowickiego, Katowice, 6, 1958.
12. KWIECIEM-LECHOWICZ, K. : The climate of Hel (peninsula) - Wiadomości Służby Hydrol.- Meteorolog., Warszawa, VI, 4, 1958.
13. LEWIŃSKA, J. : The winds of Rytno and Rymanów-Spa - Przegląd Geofizyczny, Warszawa, 1, 1958.
14. LEWIŃSKA, J. : The direction of low-level winds in the Polish Tatra Mountains - Wiad. Służby Hydrol.-Meteorolog., Warszawa, VI, 5, 1958.
15. LIMAN, S. : Some problems of health-resort climatology - Gazeta Obserwatora PIHM, Warszawa, 9, 1958.
16. LIMAN, S. : Microclimatological investigations in Rymanów-Spa. - Wiadomości Uzdrowiskowe, Poznań, 4, 1957.
17. MACKIEWICZ, S., STEFFEN, J. : Appearance of rheumatic illness in the region of Great Poland - Polskie Archiwum Medycyny Wewnętrznej, Warszawa, 3, 1957.
18. MARCINIAK, S. : The porosity of protective glasses to ultraviolet radiation depending on their temperature - Prace Centralnego Instytutu Ochrony Pracy, 2, 24, 1958.

* Publications on Human Bioclimatology in Poland, a supplement for 1957-1958, collected by mr. S. Tyczka, Instytut Balneoklimatyczny, Poznań, Poland.

19. MICHALCZEWSKI, J. : The cloudiness and duration of sunshine at Zakopane during the period 1924 - 1948 - Wiad. Służby Hydrol.- Meteorolog., Warszawa, VI, 5, 1958.
20. NOWAKOWSKI, B.; WOŹOSZYŃSKA, A.: The protection of man from atmospheric pollution - Biuletyn Służby Sanitarnej- Epidemiologicznej województwa katowickiego, Katowice, 4, 1958.
21. ORLICZ, M., ZYCH, S., PASZYŃSKI, J.: Climatological conditions of Zakopane and its surrounding regions - Wiadomości Służby Hydrol.- Meteorolog., Warszawa, 5, 1958.
22. PARCZEWSKI, W. : On air flowing over hills, especially the Tatra Mountains - Wiadomości Służby Hydrol.-Meteorolog., Warszawa, VI, 5, 1958.
23. PARCZEWSKI W. : On the interdependence of various kinds of clouds and the distribution of sunshine - Przegląd Geofizyczny, Warszawa, 4, 1957.
24. PASZEK-BARANOWSKA, M. : The problem of criteria of sultry weather - Gazeta Obserwatora PIHM, Warszawa, 8, 1957.
25. PILARSKA-MYCIELSKA, E. : Inversions of temperature in the area of Rabka-Spa, Obidowa and Nowy Targ (High Tatra Region) - Gazeta Obserwatora PIHM Warszawa, 9, 1957.
26. POL, W. : The change in eye-pupil width, visual acuity and the direction of the optical axis of the eyeballs under influence of anoxemia in man at a height of 7000 m. - Lekarz Wojskowy, Warszawa, 6, 1958.
27. PUŁAWSKI, Z. : Ultraviolet radiation, its importance in hygiene and methods of measurement - Ochrona Pracy, Warszawa, 7, 1957.
28. SIKORA, E. : Curative qualities at the maritime spa of Ustka - Wiadomości Uzdrawiskowe, Poznań, 2, 1958.
29. SŁOMKA, J., OTPIŃSKA, K. : The yearly cycle of air pollution in Wrocław - Prace Wrocławskiego Towarzystwa Nauk. s.B., PWN Wrocław, 92, 1957.
30. ŚMIAŁOWSKI, T. : Atmospheric pollution in the industrial district of High Silesia - Biuletyn Służby Sanitarnej-Epidemiologicznej województwa katowickiego, Katowice, 2, 1958.
31. SPIOCH, F. : The level of ascorbic acid in plasma and sweat of man under the influence of damp heat - Tygodnik Lekarski, Warszawa, 33, 1958.
32. TARANOWSKA, S. : Appearance of winds on the Polish Baltic seashore - Wiadomości Służby Hydrolog.-Meteorolog., Warszawa, VI, 1, 1957.
33. TRYBOWSKI, Cz. : The bioclimate of Rabka-Zdrój - Przegląd Geofizyczny, Warszawa, 1, 1958.
34. TRYBOWSKI, Cz. : Introductory results on cooling investigations in Rabka-Spa - Wiadomości Służby Hydrolog.-Meteorolog., Warszawa, VI, 5, 1958.
35. TRYBOWSKI, Cz. : An outline of the climate in Rabka-Spa - Wiadomości Uzdrawiskowe, Poznań, 2/3, 1957.
36. TRYBOWSKI, Cz. : Winds in the lower atmosphere in Rabka (Carpathian Mountains) - Wiadomości Służby Hydrologicznej i Meteorologicznej, Warszawa, 5, 1957.
37. TYCZKA, S. : The problems, research methods and development of human bioclimatology - Biuletyn Służby Sanitarnej- Epidemiologicznej województwa katowickiego, Katowice, 6, 1958.
38. TYCZKA, S. : Methods of bioclimatological investigations - Gazeta Obserwatora PIHM, Warszawa, 4, 1958.
39. WÓJTOWICZ, W. : International Geophysical Year and medical climatology - Polski Tygodnik Lekarski, Warszawa, 42, 1957.
40. WÓJTOWICZ, W. : Some problems of meteoropathology - Wiadomości Uzdrawiskowe, Poznań, 4, 1957.
41. ZMIJEWSKI, M. : Some remarks on atmospheric pollution in Gdynia - Wiadomości Służby Hydrolog.-Meteorolog., Warszawa, VI, 4, 1958.
42. ZYSK, J. : Some problems of physiopathology in connection with stratospheric flights - Lekarz Wojskowy, Warszawa, 7/8, 1957.

PART V

COSMIC BIOCLIMATOLOGY

(1959)

Section A : General cosmic bioclimatology

PALEO-BIOCLIMATOLOGY

Section A: General paleo-bioclimatology

THE IMPACT OF PALEOMAGNETISM ON PALEOCLIMATIC STUDIES

by

Dr. N.D. Opdyke ¹⁾

Throughout the history of geology there have been many efforts made to interpret the climates of the geological periods. The methods employed have varied widely and have met with varying success. As we might suppose the climate of the Cenozoic (2) is much better known than the climate of earlier geologic time. This is largely due to the fact that many existing flora and fauna have survived from the earlier Cenozoic or have closely related Tertiary forms; therefore, it is possible in many cases to infer the climatic conditions of the Cenozoic by extrapolation from the climatic zones occupied by these forms today.

However as we go back in time to the earlier geologic periods confusion reigns. This confusion is a direct result of finding climatic indicators in areas where they are not logically expected; such as coral reefs in Arctic areas, and the great evaporite (3) development in the Devonian of arctic and subarctic Canada. Also dramatically out of place are the tillites (4) of Africa, India, Brazil and Australia. These facts have lead to a number of contradictory theories concerning past climates. The hypothesis which is seen in many introductory texts on historical geology is the "uniformly warm earth theory" propounded to explain the occurrence of sediments indicative of warm conditions in the polar regions; however, in at least one period, the Permian, this hypothesis does not appear to be valid because climatic indicators show that while glaciers were in existence close to the equator it was warm in the Arctic regions. The only theory that has satisfactorily explained the apparent climatic distribution of the past is the theory of continental displacement first advocated by Taylor (1910) and Wegener (1912) and championed in recent years in a modified form by Dutoit, 1937; Carey, 1955; and King, 1953. This theory has not yet been accepted by the majority of geologists; however, it still remains as the only theory which satisfies most of the evidence.

In recent years a potentially powerful tool has been developed by geophysicists which promises to be of great value in solving the problems of Continental Drift and could be of great assistance to the paleoclimatologist. This new tool is paleomagnetism, which is the study of the magnetic field in the geologic past reconstructed by using the magnetic moments retained in certain igneous and sedimentary rocks; such as lava flows, red sandstones and shales. It has been shown by the workers in this field that certain rocks retain the direction and dip of the magnetic field which was active while they were being deposited. This allows the worker to determine the position of the magnetic pole relative to the locality where the specimens were collected. It has also been shown that over a length of time, long compared to the period of the secular variation, the magnetic pole will coincide with the geographic pole, thus allowing the determination of the latitude of the collecting site. (The writer will not attempt to describe in detail the techniques and assumptions used in measuring and in interpreting paleomagnetic results since excellent reviews are available. Runcorn, 1957). Thus it is possible to determine the approximate latitude for a continent for any geological period in which suitable rocks are available.

(1) Formerly Dept. of Geology, The Rice Institute, Houston, Texas, U.S.A.

At present Dept. of Geophysics, Research School of Physical Sciences, The Australian Nat. Univ., Box 4 G.P.O., Canberra, Australia.

(2) As several members of I.S.B.B. may not be acquainted with the geological terminology used in this article and other ones related to Paleo-bioclimatology, in the annex attached a summary is given of the most important stratigraphic names which have been introduced to indicate different geological periods in the past (Editorial note).

(3) EVAPORITES are sediments which originate from water in hot climates by evaporation of lake- or seawater.

(4) TILLITES are boulder clay deposits of Late Carboniferous or Lower Permian age which were formed during an Upper Carboniferous Glacial Period in the present Southern Hemisphere.

It follows then that if the latitude of a particular continent for a period is known, it should be possible to construct a hypothetical climatic picture for the continent, taking into account the present distribution and causes of climate.

It has been shown that paleomagnetic results are in general agreement with paleoclimatic indicators (Irving, 1956; Schove et al., 1958); that is to say, that when paleomagnetic results indicate a position close to the equator the associated sedimentary deposits usually show evidence of a warm or arid environment (e.g. evaporites, desert sandstones, bioherms (5), bauxites (6); also it has been shown that when high latitudes are indicated, glacial tillites are sometimes associated with them (e.g. South Africa and Australia during the Permo-Carboniferous). Thus there is gross agreement between paleomagnetic results and paleoclimatology.

The next step would seem to be the construction of the probable climatic zones of continents for certain periods given the rough paleogeography of the continent and its position relative to the poles and equator. While it should be possible to construct probable climatic zones for a geological period given the aforementioned conditions, it must be kept in mind that any attempt at absolute accuracy in drawing the boundaries of the climatic zones will lead to difficulty; one reason being that modern climatic zones are drawn rather arbitrarily when in fact the different zones grade imperceptibly into one another. Another reason is that a rigid interpretation will lead to erroneous results if we take into consideration the fact that climatic zones are known to expand and contract. Climatic changes of this type are known to have taken place since the retreat of the last glaciers in Europe and North America. Profound changes of climatic zonation will occur during any major glaciation and are well documented for the Pleistocene.

Certain generalities regarding climate should be constant throughout geological time if the following axioms are true:

1. that the earth has rotated throughout time in the same direction without a large change in the velocity of rotation;
2. that the inclination of the earth's axis has remained at or close to its present value of $23^{\circ} 26' 44.8''$ (Skilling and Richardson, 1947).

There is no reason to believe that these factors have significantly altered since the Cambrian because most of the other planets, whose direction of rotation and angles of inclination to the ecliptic are known, rotate on their axis counter clockwise, the same as the earth, and have comparable ecliptic angles (e.g. Mars is inclined to the ecliptic at an angle of $23^{\circ} 98''$) (7)

If these factors have remained more or less constant then the earth's relationship to the sun probably has remained the same through geologic time. This is important since the sun is the primary cause of the earth's atmospheric energy and climate. Since the earth possesses an atmosphere and is an oblate spheroid, it follows that more heat energy will always be available at the equator per unit area than at the poles. We have also seen that the earth's axis is inclined at an angle of $23\frac{1}{2}^{\circ}$ which is directly responsible for the seasons. This causes total darkness above $66\frac{1}{2}^{\circ}$ in December in the northern hemisphere and in June in the southern hemisphere. These facts cause the progressive lowering of the mean annual temperature from the equator to the poles. Furthermore, the sun through the heating of the atmosphere causes it to be set in motion. If the earth were not rotating a single large cell would be developed rising at the equator and settling at the poles. This picture does not develop because the rotation of the earth exerts a force (the Coriolis force) on the moving air masses which breaks the large cell down into the components which are observed today: the prevailing easterlies or trade winds, the mid-latitude westerlies and the polar easterlies. These factors coupled with the physical geography of the earth cause climatic patterns.

Therefore, certain broad generalities regarding the earth's climate should hold true through geologic time.

- A. THE PLANETARY WIND SYSTEM SHOULD REMAIN MUCH AS IT IS TODAY giving rise to an equatorial low pressure zone (0°), trade wind zone ($0 - 30^{\circ}$), subtropical high ($30 - 35^{\circ}$), zone of westerly winds ($35 - 60^{\circ}$), subpolar low pressure area (65°), polar easterlies ($65 - 90^{\circ}$) and a polar high pressure area ($80 - 90^{\circ}$).

-
- (5) BIOHERMS or ORGANIC REEFS: name introduced in 1928 by Cummings and Shrock to indicate reef-like, moundlike or lens shaped carbonate masses consisting chiefly of debris from organisms (coral, algae, crinoids etc.) and surrounded by impervious rocks.
 - (6) Under certain conditions in tropical climates all the silica of felspar minerals may be removed by weathering. The residue left is called BAUXITE, a mixture of two aluminous minerals ($Al_2O_3 \cdot H_2O$ and $Al_2O_3 \cdot 3H_2O$), often a valuable aluminium ore.
 - (7) URANUS is an exception having a very steep ecliptic angle of 98° and is rotating clockwise.

B. PRECIPITATION WOULD ALSO TEND TOWARD SOME TYPE OF ZONATION RESTING ON THE HIGH AND LOW PRESSURE AREAS:

1. The equatorial low pressure zone should produce high precipitation as it does today.
2. Deserts could be expected to develop between 15° and 35° north of the equator caused by the subtropical high. Mid latitude deserts would tend to develop in the east central portion of the westerly zone.
3. In the absence of mountain barriers the portions of the continents facing the prevailing westerlies should receive abundant moisture.

C. MONSOONAL CONDITIONS COULD BE EXPECTED TO DEVELOP ON THE SOUTHEASTERLY PORTIONS OF THE CONTINENTS (in the northern hemisphere).

D. WINTERS COULD BE EXPECTED TO BE MORE SEVERE ON THE POLEWARD SIDE OF CONTINENTS even in the absence of ice caps.

Although this general picture is probably valid there will always be modifications due to geographic factors such as the changing relations of land and sea, the rise of mountain chains and the influence of ocean currents. These factors can be very important and their effect on the present climate of the earth is well known; however, the effect of change in latitude must always be present and the basic pattern must persist even if modified.

The construction of the climatic zones for a hypothetical continent in a nonglacial period is shown in Fig. 1 (after Finch and Trewartha, 1949); therefore, if the approximate outline of a continent is known with its probable latitude, a reasonable reconstruction of its climatic zones is possible for a geological period. The paleogeography of some continents is known in its essential parts and the study of paleomagnetism gives the approximate latitude. Thus the major components are present for reconstructing past climates.

Using the procedure outlined above the writer has attempted to reconstruct the climatic zones of North America for the Pennsylvanian (Upper Carboniferous), Permian and Triassic and for the Permian of Europe. Relatively good paleogeographic maps exist for North America for the periods under study and for Europe during the Permian, Schuchert, 1955; Eardley, 1951; H. Termier and G. Termier, 1952. No attempt has been made to divide the periods into the different stages and the maps show the maximum extent of the epicontinental seas for these periods. The writer does not contend that these maps are extremely accurate but they do show the relative position of land and sea. The first map series (indicated with A1, A2 etc) shows the probable climatic zones for the different periods with the continents oriented as they are at the present and the second series (indicated with B1, B2 etc.) shows the climatic zones that follow from the orientation of the continents which is suggested by paleomagnetism.

On these maps have been placed the locations of occurrence of sediments which the writer believes to have paleoclimatic significance. These are:

1. Bauxite, tropical or subtropical climate with high temperature and possibly seasonal rainfall (Harder, 1949).
2. Evaporites, high temperature causing high rates of evaporation coupled with low precipitation.
3. Coal, adequate precipitation.
4. Bioherms, high water temperature, causing super saturation of sea water with CaCO_3 which allows rapid biochemical fixation of calcium carbonate (Rankama and Sahama, 1950).

Also shown on these maps are:

5. Direction of dominant wind determined by studies of eolian dune sands. Usually representing sand dune action in desert areas (Opdyke and Runcorn, 1959).
6. Direction of dominant wind predicted by the orientation of the continent.

The discussion of the different periods covering the reconstructions based on the two different orientations will start with the Permian of Europe.

Figure A.1 shows the hypothetical climatic zones for Europe during the Permian given its present latitude. If Europe lay in the same latitudes that it does at the present the continent would remain largely in the zone of the mid latitude westerlies. These westerly winds would tend to bring abundant moisture in from the Atlantic Ocean much as it does today. However, in the absence of ice caps the continent would probably be much warmer than it is at present. The subtropical zone may have extended farther north than the position shown on the map, but the fact remains that western Europe would remain in an area of abundant moisture.

Figure B.1 shows Europe oriented as indicated by paleomagnetism (Shove, Nairn and Opdyke, 1958). Most of Europe remains in the northern hemisphere; however, it is now almost entirely in the tropical zone. The hypothetical reconstruction of the climate is shown. It is immediately apparent that the most important difference between A.1 and B.1 would be the predicted appearance of a Permian desert in western Europe due to the shift of wind from the west to the east.

The climatic evidence from the Permian of Europe would seem to support the second orientation overwhelmingly:

1. The large evaporite basins of the Urals and northwestern Europe.
2. Development of reefs in England, Belgian and Russia, indicating tropical water temperatures.
3. The wind direction deduced from dune bedded sandstones in Great Britain agrees with the expected wind direction (Shotten, 1956).
4. Permian coals which would hardly have formed in an arid climate would reasonably occur in a position closer to the equator than the evaporites.

Next we turn to the Triassic of North America. Figure A.2 shows North America oriented in its present position. Most of North America would be dominated by the westerly winds bringing moisture to the west coast and, in the absence of a mountain barrier in Triassic times, would bring moisture into the interior. A mild monsoonal regime would probably be established in the southeast bringing a seasonal aspect to the precipitation in this part of the continent.

Figure B.2 shows North America oriented as indicated by paleomagnetism (Du Bois et al., 1957). Most of North America remains in the northern hemisphere but approximately 30° closer to the equator than it is at present. Thus most of North America would lie in the cone of the northeasterly trades. This would cause a belt of rainfall to the south with an arid climate developing in the west central portion of the continent.

The climates of the North American Triassic as deduced from the sedimentary indicators would seem to favor the climatic pattern presented in B.2:

1. The development of thick evaporites in Alberta (700 ft.) and thinner evaporites in the Wyoming shelf area (Hunt and Ratcliff, 1959; Krumbein, 1951).
2. Development of Triassic desert sand dunes in Arizona, Utah, and Colorado indicating a dominant northerly wind (Poole and Williams, 1956).
3. Triassic coals in Virginia and North Carolina indicating abundant moisture.
4. Existence of bioherms in Nevada indicating conditions of warm water (Twenhofel, 1950).
5. Evaporites in southern Mexico possibly indicating conditions of aridity south of the Triassic equator (Benavides, 1956).

The Permian of North America will be considered next. Figure A.3 shows the probable climatic zones based on the present orientation of North America. Conditions as shown are much the same as those given in Figure A.2 for the Triassic. Figure B.3 shows North America oriented as suggested by paleomagnetism (Runcorn, 1956). Most of North America would seem to lie in the tropical zone between 30°N and 30°S . The zonation corresponding to this position is given.

The climatic indicators from the Permian seem to indicate warm, arid conditions:

1. The Permian reef complex of the West Texas basins.
2. Evaporite deposition in Kansas, Oklahoma, Texas, New Mexico and Arizona.
3. Desert dune sandstone in Arizona and Wyoming indicating a north or northwest wind (Beiche, 1938).

It can be seen that neither of the proposed climatic schemes fit the distribution of the climatic indicators precisely. The wind direction deduced from the Coconino sandstone of Arizona could be a result of the northeasterly trade wind if the paleomagnetic data were slightly in error. This is a possibility because the results are mainly from one formation; e.g. the Supia formation (L. Perm.). Also the arid zone in the Southwest could possibly be a wind shadow effect of the mountains formed in the Appalachian Orogeny (8); however, it is concluded that A.3 and B.3 could both lead to the arid conditions which existed at this time in the western United States.

The Pennsylvanian paleogeography oriented in the present position of North America, Figure A.4, should lead to warm humid conditions penetrating deep into the middle of the continent under the influence of the prevailing westerlies with the subtropics extending farther toward the north than they do at the present.

The orientation of the continent suggested by paleomagnetism (Runcorn, 1956) on the other hand would place most of the continent again within the tropical zone with the equator passing diagonally across the continent from Newfoundland to the Gulf of California. This orientation would place most of the continent north of the equator in the path of the northeast trade wind and that portion of the continent remaining south of the equator would be influenced mainly by the southwest trades. A belt of high precipitation should run diagonally across the continent, being

-
- (8) The APPALACHIANS represent a mountain belt in the eastern U.S.A. and Canada along the Atlantic Coast, mainly formed during a period of mountain building at the end of the Carboniferous and during the Lower Permian (so called Hercynian folding period of Europe).

broad in the east and narrow in the west. Arid conditions could be expected to develop in the west central part of the country caused by the easterlies having to cross the bulk of the continent.

The sedimentary climatic indicators present in the North American Pennsylvanian are:

1. Large amounts of coal formed in the east and east central part of North America. Indicating abundant rainfall in these areas.
2. Bauxite occurs in Pennsylvania and Missouri indicating tropical conditions for these regions (Harder, 1949).
3. Bioherm development in Texas indicating conditions of warm water (Bergsenbeck and Terrierre, 1953).
4. Evaporites developed in the Black Hills region, the Gypsum basin Colorado and the Paradox basin Utah, the latter containing 4000 ft. of evaporites (Krumbein, 1951).
5. Dune bedded sandstones in Wyoming, northeast Utah and northwest Colorado show, on the basis of cross-stratification studies, that the depositing wind was dominantly a northeast wind (Opdyke and Runcorn, 1959).

We can see from a comparison of Figures A.4 and B.4 that the distribution is consistent with B.4 and not A.4. For B.4 predicts warm, humid conditions in eastern and northeastern North America and increasing aridity in the West, whereas A.4 does not indicate any aridity in the West. The direction of the dominant wind given above is also consonant with B.4. It might be noted that the areas of the greatest accumulation of evaporites in the Paradox and Gypsum basins would lie on the wind shadow of the Uncompagnian and Colorado Range positive areas if the wind was from the east but not if it was from the west. Thus being more consistent with the orientation derived from paleomagnetism.

CONCLUSIONS

If the theories concerning paleomagnetism are correct then this new branch of the earth sciences offers a means of determining lines of latitude independent of paleoclimatic study. Given these determinations in combination with paleogeographic studies it should be possible to reconstruct a hypothetical scheme of climatic zonation based on climatic principles known at the present time. This has been attempted in figures B.1 to B.4. It has been shown that these reconstructions fit the known distribution of certain climatic indicators in three of the four cases studied in detail. It is apparent that this method can be used to test whether the continents have remained stable in their present position or have moved relative to the poles as suggested by paleomagnetism. The results obtained support the paleomagnetically determined positions of the continents.

The author realizes that the reconstruction of the climatic belts herein presented can only be tentative but with an increase in the knowledge of paleogeography and more precise determinations of pole positions with the growth of paleomagnetism refinement in detail may be attempted. If accurate reconstructions of climate in the past are finally obtained it will be of great assistance to paleoecological and paleo-bioclimatological studies and may help us to explain the observed geographical distribution of plants and animals through geologic time. It is the hope that this paper will stimulate more thinking along these lines.

REFERENCES


1. BENAVIDES, L.G. : Notas sobre la Geologia Petrolera de Mexico - Sym. sobre yacimientos de Petroleo y Gas XX Cong. Geol. Inter., 3, 351-362, 1956.
2. BERGENBECK, R.E. and TERRIERRE, R.T.: Petrology and Petrography of the Scurry Reef, Scurry County, Texas - Amer. Assoc. Pet. Geol. Bull., 37, 5, 1014-1029, 1953.
3. CAREY, S.W. : The orocline concept in geotectonics, Part I - Roy. Soc. Tasmania, Pap. o Pr., 89, 255-288, 1955.
4. DUBOIS, P.M., IRVING, E., OPDYKE, N.D., RUNCORN, S.K. and BANKS, M.R.: The geomagnetic Field in Upper Triassic Times in the United States - Nature, 180, 1186-1187, 1957.
5. DUTOIT, A.L. : Our Wandering Continents - Oliver and Boyd, London, 366 p., 1937.
6. EARDLEY, A.J. : Structural Geology of North America - Harper & Bros., New York, 624 p, 1951.
7. FINCH, V.C. and TREWARTHA, G.T.: Physical Elements of Geography, 3rd Ed., Mc Graw Hill, New York, 536 p., 1949.
8. HARDER, E.C. : Stratigraphy and Origin of Bauxite Deposits - Geol. Soc. Amer. Bull., 60, 887, 1949.

9. HUNT, A.D. and RUTCLIFFE, J.D.: Triassic Stratigraphy, Peace River area Alberta and British Columbia, Canada - Amer. Assoc. Pet. Geol. Bull., 43, 3, 563 - 589, 1959.
10. IRVING, E. : Paleomagnetic and Paleoclimatological aspects of polar wandering - Geofisica Pura e Applicata, 33, 1-20, 1956.
11. KING, L.C. : The necessity for continental drift - Amer. Assoc. Pet. Geol. Bull., 37, 9, 2163-2177, 1953.
12. KRUMBEIN, W.C. : Occurrence and Lithologic Associations of Evaporites in the United States - Jour., 8 ed., Pet., 21, 2, 63-81, 1951.
13. OPDYKE, N.D. and RUNCORN, S.K.: Paleomagnetism and ancient wind directions - Endeavor, 18, 69, 26-34, 1959.
14. POOLE, F.G. and WILLIAMS, G.A.: Direction of transportation of the Sediment Constituting the Triassic and associated formations of the Colorado Plateau - Inter. Cong. on Peaceful Uses of Atomic Energy, U.N., 326-330, 1956.
15. RANKAMA, K. and SAHAMA, Th.G.: Geochemistry - Univ. of Chicago Press, 911, 1950.
16. REICHE, P. : An analysis of Crosslaminations in the Coconino Sandstone - Jour. Geol., 46, 905-932, 1938.
17. RUNCORN, S.K. : Paleomagnetic survey in Arizona and Utah, Preliminary results - Geol. Soc. Amer. Bull., 67, 301-316, 1956.
18. RUNCORN, S.K. : The sampling of rocks for Paleomagnetic comparisons between the continents - Adv. in Phy., 6, 22, 169-176, 1957.
19. SCHOVE, D.J., NAIRN, A.E.M. and OPDYKE, N.D.: The climatic geography of the Permian - Geog. Ann., 49, 3-4, 216-231, 1958.
20. SCHUCHERT, C. : Atlas of Paleogeographic Maps of North America - John Wiley & Sons, New York, 177 p., 1955.
21. SHOTTON, F.W. : Some aspects of the New Red Desert in Britain - The Liv. and Man. Geol. Jour., 1, 450-466, 1956.
22. SKILLING, W.T. and RICHARDSON, R.S.: Astronomy - Henry Holt and Co., New York, 692 p., 1947.
23. TAYLOR, F.B. : Bearing of the tertiary mountain belt on the origin of the earth's plan - Bull. Geol. Soc. Amer., 21, 2, 179-226, 1910.
24. TERMIER, H. and TERMIER, G.: Histoire Géologique de la Biosphère - Masson and Co., Paris, 721 p., 1952.
25. TWENHOFEL, W.H. : Coral and other organic reefs in the geological column, - Amer. Ass. Pet. Geol. Bull., 34, 182-202, 1950.
26. WEGENER, A. : Die Entstehung der Kontinente - Peterm. Mitt., 185-195, 253-256, 305 - 309, 1912.
27. WEGENER, A. : Die Entstehung der Kontinente and Ozeane - Sammlung Vieweg, Braunschweig, no. 23, 1915.

ABSTRACT

Paleomagnetism presents the paleo-bioclimatologist for the first time with an independent determination of latitude. In combination with paleogeography a reconstruction of climatic zones for the geologic periods can be attempted by taking into account the principles underlying the present distribution of climate. This was attempted for the Permian of Europe and the Triassic, Permian and Pennsylvanian (U. Carboniferous) of North America. An extension of this method can be of great importance in future paleo-ecological and bioclimatological studies if the underlying principles of Paleomagnetism prove to be correct.

Fig. 1




The climatic zonation of a hypothetical continent in a nonglacial period (Modified from Finch and Trewartha 1949)

Fig. A.1




The paleogeography and climatology of the Permian of Europe, oriented as it is at present. In the different figures the large arrow shows the presumed dominant wind direction, the small arrows show the dominant wind direction as deduced from cross-stratification studies of aeolian sandstones, the cross-hatching shows areas of evaporite deposition, the "C" designates the accumulations of coal and "X" represents reef development, "O" represents accumulations of bauxite.

Fig. B.1




The paleogeography and climatology of the Permian of Europe, oriented as indicated by paleomagnetism.

Fig. A.2




The paleogeography and climatology of the Triassic of North America oriented as it is at present.

Fig. A.3




The paleogeography and climatology of the Permian of North America oriented as it is at present.

Fig. B.2




The paleogeography and climatology of the Triassic of North America oriented as indicated by paleomagnetism.

Fig. B.3




The paleogeography and climatology of the Permian of North America oriented as indicated by paleomagnetism.

Fig. A.4



The paleogeography and climatology of the Pennsylvanian of North America oriented as it is at present.

Fig. B.4



The paleogeography and climatology of the Pennsylvanian of North America oriented as indicated by paleomagnetism.

C O S M I C B I O C L I M A T O L O G Y

Section A: General Cosmic Bioclimatology

BLOOD CLOTTING TIME UNDER METAL COVER (BIOLOGICAL P-TEST)

by

Prof. Shinji Itoh, Keisuke Tsujioka and Hiroo Saito (Japan)*

Inorganic colloid systems are greatly influenced by various environmental conditions. To determine extra-terrestrial factors which may affect the colloid reactions, Piccardi (1) proposed a test method entitled the "CHEMICAL TEST".**

For this purpose he used the formation of insoluble bismuth oxychloride which is produced by the hydrolysis of bismuth chloride. In his "P-test", the reaction is observed under two different conditions: one in a metal-free room and another in a room where it is completely covered by a thin copper sheet 1 mm thick. Each pair of experiments is carried out simultaneously inside and outside a copper box and the rate of precipitation compared. If the formation of the colloidal precipitation of bismuth is faster inside than outside of the box, it is recorded as "T". The result of ten measurements is expressed as T per cent. Piccardi also measured the clotting time of human blood inside and outside the copper box, and found that the time is always longer inside than outside (2). He called this clotting reaction under a copper sheet "BIOLOGICAL P-TEST". However, this conclusion may not be acceptable. If T per cent is high, clotting time inside the box should be shorter than that outside of the box. His experiments were carried out in Florence during August 1955, when T per cent throughout the month was below 50 per cent. (1) This might be the reason why he obtained such a one-sided result. To clarify this problem we measured the clotting time of rabbit blood by the method of Sahli and Fonio inside and outside a copper box simultaneously. The ratio of the time outside/inside was compared with T per cent which was determined at the same time. One hundred observations were carried out during a period from November 1958 to June 1959. The results are illustrated in Fig. 1.

Though individual values fluctuated considerably, the mean value of each group showed a consistent tendency: a low ratio of the clotting time when the T per cent was low and vice versa. That is, when the rate of colloidal precipitation of bismuth is slower in the copper box than outside of it, the clotting time of blood is also slower in the box. Statistical analysis indicated that the results are highly significant. Thus, the "BIOLOGICAL P-TEST" is proven to be correlated with the inorganic P-test.

REFERENCES:

1. PICCARDI, G. : Physikalische, chemische und biologische Effekte als Ausdruck von Weltraumeinflüssen - Wien, Med. Wschr., 106, 975-980, 1956.
2. PICCARDI, G. and DANTI, M. : Wirkung einer Metall-Abschirmung, ein biologischer Test - Wetter und Leben, 7, 7, 1955.

* Department of Physiology, Hokkaido University School of Medicine, Japan.

** For further details see Int. Journal of Biocl. and Biomet., vol. II, 1958, Part V, sect. A.




Fig. 1 Relationship between the ratio of the clotting time (outside/inside) and T per cent of P-test. White circles represent mean values of the clotting time ratio for each group of the T% values.

Section B : Special cosmic bioclimatology

Section C : World literature

PART VI

PALEO-BIOCLIMATOLOGY

(1959)

Section A : General paleo-bioclimatology

Section B : World literature

PART VII

MISCELLANEOUS BIOCLIMATOLOGICAL DATA

Scientific Committees of the Society

Symposia or Congresses

Bioclimatological Stations and Institutions

Requests from Members

Requests from non-members

Book Reviews

International Organizations

Advertisements

(1959)

Section A : Scientific committees of the Society

1. Reports

- a. Allergic diseases
- b. Ecological climatography
- c. Instrumentation
- d. Ionisation of the air
- e. Nautical bioclimatology (general-, cargo bioclimatology)
- f. Chemical tests
- g. Tropical bioclimatology
- h. Solar Radiation

2. Literature

Section B : Symposia or congresses of international
organizations and of national
bioclimatological societies (dates, programmes,
summaries of important lectures, decisions, etc.)

M I S C E L L A N E O U S B I O C L I M A T O L O G I C A L D A T A

Section B: Symposia of Bioclimatological Societies

SYMPOSIUM INTERNATIONAL SUR LES RELATIONS ENTRE PHENOMENES SOLAIRES

ET TERRESTRES EN CHIMIE-PHYSIQUE ET EN BIOLOGIE *

A l'initiative du PROFESSEUR G. PICCARDI, Directeur de l'Institut de Chimie physique de l'Université de Florence et sur l'invitation du PROFESSEUR P. BOURGEOIS de l'Université de Bruxelles, Directeur de l'Observatoire Royal de Belgique, un Symposium international s'est tenu à l'Observatoire Royal de Belgique, à Uccle, les 8, 9 et 10 octobre 1958.

Il était agréé par le Commissariat Général du Gouvernement près l'Exposition Universelle et Internationale de Bruxelles 1958.

I. L E B U T D U S Y M P O S I U M

Le but de ce Symposium était essentiellement un examen critique et une synthèse des observations effectuées depuis 1950 par la METHODE DIFFERENTIELLE ET STATISTIQUE DU PROFESSEUR PICCARDI.

Les hypothèses interprétatives des fluctuations périodiques, observées en fonction du temps, dans les résultats de divers tests physico-chimiques inorganiques, lient celles-ci à l'activité solaire et au mouvement de la Terre dans la Galaxie. Une interprétation relativiste d'une certaine fluctuation annuelle a été présentée comme une conséquence d'une théorie générale de la gravitation universelle.

Le rapprochement de ces observations avec celles d'ordre biologique ou médical qui mettent en cause l'influence de facteurs atmosphériques ou cosmiques ouvre une perspective nouvelle dans l'étude de ces facteurs et de leurs effets pathologiques ou sociaux.

L'exposé introductif du Professeur Piccardi situe la portée des problèmes abordés sur le plan philosophique et humain. Il souligne notamment l'importance essentielle d'ordonner en fonction du temps les résultats d'un grand nombre d'observations semblables. La prise en considération de la variable temps, peu habituelle dans les expériences physico-chimiques de laboratoire, a permis de donner un sens aux résultats statistiques non reproductibles d'un ensemble d'expériences apparemment identiques.

Les causes de cette non-réproductibilité apparaissent ainsi liées à la variabilité des conditions physiques de l'espace environnant, qui, dans la plupart des conditions expérimentales usuelles, échappe à tout contrôle.

Ceci met en question, aussi bien dans certains domaines de la physico-chimie et de la biologie que de la médecine et de la sociologie, des critères de validité admis sans conteste jusqu'ici, dans les méthodes expérimentales d'investigation scientifique.

II. O B J E T D E S C O M M U N I C A T I O N S

E T D I S C U S S I O N S

PROF. G. PICCARDI (Directeur de l'Institut de chimie physique de l'Université de Florence): "EXPOSE INTRODUCTIF"

Cet exposé introductif apporte, en dehors des considérations générales déjà indiquées, un aperçu des systèmes sensibles aux facteurs météorologiques ou cosmiques, dans un bref historique du développement des recherches.

Il souligne particulièrement l'importance biologique, parmi ceux-ci, de l'eau et des systèmes colloïdaux.

Il évoque finalement les liens qui rattachent le thème des discussions du Symposium à celui "....pour un monde plus humain", de l'Exposition Universelle et Internationale de Bruxelles 1958,

* Prepared by Prof. G. Piccardi, Director Istituto di Chimica Fisica, Via Gina Capponi 9, Firenze, Italy.

qui a accordé son patronage à la réunion. Il en conclut: "C'est seulement en sachant par quel mécanisme il est lié à la terre et au ciel, que l'homme pourra comprendre mieux sa position physique et psychique dans l'Univers d'aujourd'hui".

PROF. G. PICCARDI: "LES TESTS CHIMIQUES"

Le professeur Piccardi expose ensuite le principe de la méthode des "tests chimiques" et les résultats de toutes les observations faites par cette méthode à Florence. Ceux-ci mettent remarquablement en évidence, sur une réaction d'hydrolyse, une fluctuation annuelle superposée à une variation séculaire qui a pu être statistiquement rattachée à l'activité solaire.

Ils montrent aussi la grande sensibilité de l'eau et des systèmes colloïdaux instables à l'action de divers facteurs physiques naturels ou artificiellement appliqués, les conditions différentielles et statistiques propres à la méthode, excluant les effets directs bien connus de la température ou de la pollution des réactifs.

MME. C. CAPEL-BOUTE (Fac. des Sciences Appliquées de l'Université de Bruxelles): "OBSERVATIONS SUR LES TESTS CHIMIQUES DE PICCARDI EFFECTUEES A BRUXELLES DE 1950 à 1958"

Mme C. Capel-Boute apporte la confirmation du caractère général des facteurs influençant cette réaction d'hydrolyse, par les corrélations, étudiées statistiquement, entre des observations semblables faites simultanément à Bruxelles et à Florence. Celles-ci laissent prévoir cependant des effets météorologiques locaux ou des effets de latitude, variables selon la période du cycle de l'activité solaire.

Les possibilités et les limites de la méthode ont pu être discutées en fonction des conditions opératoires, celles-ci ayant été préalablement concrétisées pour les participants au Symposium, par la visite du laboratoire de Mme Capel, installé à l'Observatoire Royal de Belgique.

Dr. UDO BECKER (Frauenhofer Institut - Freiburg i.B.): "SONNENAKTIVITAET UND CHEMISCHE TESTS"

M.U. Becker présente l'étude des corrélations, statistiquement assurées, entre les résultats des tests chimiques obtenus à Florence et les caractères de l'activité solaire, taches et éruptions, dans leur variation séculaire. Cependant une périodicité annuelle dans les fluctuations des tests chimiques, apparaît sans relation avec l'importance des taches solaires. L'allure annuelle sinusoïdale de certains tests apparaît comme une variation saisonnière normalement liée à la position du Soleil tandis que l'allure annuelle moyenne, propre au test D, effectué dans une chambre faradisée en tôle de cuivre mince, exige une interprétation particulière.

PROF. G. PICCARDI: "UNE HYPOTHESE SOLAIRE"

Le Professeur Piccardi présente ensuite "l'hypothèse solaire" qu'il a proposée pour interpréter l'allure typique de la fluctuation annuelle du test D. Cette hypothèse repose sur la considération de la trajectoire hélicoïdale de la Terre dans la Galaxie, résultant de la combinaison de l'orbite elliptique de la Terre autour du Soleil, et de la translation du Soleil lui-même dans la Galaxie.

Un modèle matériel de ce mouvement résultant de la Terre a été présenté au Planétarium de Bruxelles dans la cadre de l'Exposition de 1958.

DR. D. QUILGHINI (Institut de Mécanique rationnelle de l'Université de Florence): "LES ELEMENTS CINEMATIQUES DU MOUVEMENT DE LA TERRE REFERE AUX ETOILES FIXES"

L'étude mathématique de cette trajectoire, faite par M. Quilghini, montre une importante variation périodique de la vitesse et de l'orientation de la Terre le long de cette trajectoire. Celles-ci présentent notamment aux équinoxes de mars et septembre, des caractères particuliers vis à vis du champ galactique.

La nouveauté de cette hypothèse solaire consiste sans doute à appeler l'attention sur l'importance que pourrait avoir dans les fluctuations de certains phénomènes terrestres, les facteurs de champs, de rayonnement ou de déviation de particules d'origine solaire ou cosmique.

Elle pourrait conduire ainsi à une conception nouvelle des phénomènes saisonniers, restes traditionnellement rattachés jusqu'ici à la simple orbite elliptique de la terre autour du soleil.

PROF. A. GIAO (Ing. Géophys. Lisbonne-Paris): "INTERPRETATION RELATIVISTE DE LA VARIATION ANNUELLE DU TEST D PHYSICO-CHIMIQUE DE PICCARDI ET SA SIGNIFICATION COSMOLOGIQUE"

Il appartient au Dr. Giso de confirmer l'importance qu'aurait la variation de la vitesse de la Terre dans son mouvement hélicoïdal dans le champ galactique. Il démontre, en effet, mathématiquement, que la variation annuelle du test D physico-chimique de Piccardi, peut apparaître comme une conséquence de la théorie générale de la gravitation universelle qu'il a proposée antérieurement.

Les liens possibles étant ainsi établis entre des actions physiques de l'espace cosmique et certains phénomènes physico-chimiques d'importance vitale, leurs conséquences d'ordre biologique ou médical peuvent être envisagées cette fois sous un angle nouveau.

PROF. H. BERG (Professeur de Météorologie et Géophysique à l'Université de Cologne): "SOLAR-TERRESTRISCHE BEZIEHUNGEN IN BIOLOGIE"

Le professeur Berg passe en revue nombre de corrélations tentées entre des observations biologiques ou médicales et des phénomènes solaires. Il en discute la valeur statistique souvent insuffisante, et critique différentes méthodes de corrélation employées dans ces recherches.

Il considère cependant la probabilité d'une influence directe ou indirecte de l'activité solaire sur certaines observations, notamment les effets biologiques liés à la coagulation du sang, comme sérieusement renforcée par les corrélations bien assurées établies entre l'activité solaire et les tests physico-chimiques de Piccardi.

Dr. F. VERING (Directeur du Bundesstaatliches Serumprüfungsinstitut de Vienne): "NACHWEIS EXTRA-TERRESTRISCHE EINFLUESSE AUF MIKROBIOLOGISCHE SYSTEME. EIN BIOLOGISCHER P-TEST"

Enfin, le Dr. Vering s'est efforcé de mettre en évidence l'influence de facteurs extérieurs spaciaux, en différenciant l'évolution d'un matériel biologique vivant, par l'action d'un écran métallique. La variabilité des résultats, soigneusement analysée en fonction des conditions opératoires pour discriminer les facteurs influençants, le conduit également à la nécessité d'une révision des conceptions expérimentales, et des principes de l'analyse causale, aussi bien dans certains domaines de la biologie, que de la médecine et de la sociologie.

Ces conceptions doivent tenir compte de la variabilité des facteurs incontrôlables qui déterminent l'irréversibilité du temps.

III. DISCUSSION GENERALE ET CONCLUSIONS

La discussion générale traite de l'importance des connaissances actuelles sur la structure de l'eau et de divers facteurs propres à la chimie colloïdale dans l'action de facteurs physiques sur l'eau et les solutions.

Elle souligne l'intérêt de recherches plus poussées sur la cinétique de coagulation et de sédimentation, prenant en considération la pureté des réactifs et les diverses actions physiques envisagées précédemment notamment celles de champs électromagnétiques de diverses fréquences, et celles d'écrans conducteurs ou non.

Elle porte encore sur l'importance relative des intensités de divers facteurs d'origine extérieure, par rapport à ceux mis en oeuvre au laboratoire, et sur la difficulté de réaliser des expériences "absolues" valables, plutôt que des essais différentiels et statistiques.

IV. COMPTE-RENDU DE LA REUNION DE LA SECTION EUROPEENNE DU COMITE INTERNATIONAL POUR L'ETUDE DES TESTS CHIMIQUES

La réunion du groupe européen du Comité International pour l'étude des tests chimiques, a examiné les conditions de développement des observations avec les tests chimiques aux diverses latitudes, au cours de l'Année Géophysique Internationale et après, grâce notamment à la collaboration active des institutions suivantes, du nord au sud:

BRUXELLES (Uccle)	: Laboratoire d'électrochimie et métallurgie de l'Université Observatoire Royal de Belgique.
JUNGFRAUJOCH	: Astronomisches Institut der Universität.
VIENNE	: Institut für Analytische Chemie der Technischen Hochschule Untertullnerbach (Oesterreich): Wiental-Wasserleitung.
GENES	: Istituto Geofisico e Geodetico dell'Università
TRIESTE	: Osservatorio Geofisico et Istituto Sperimentale Talassografico.
FLORENCE	: Istituto chimica fisica dell'Università
CASTELLANA GROTTA (di Bari)	: 2 stations (intérieur et extérieur de la Grotte)
LIBREVILLE	: Service Météorologique du Gabon.
LEOPOLDVILLE	: Service Météorologique du Congo Belge
FORT DAUPHIN	: Service Météorologique de Madagascar
ILES KERGUELEN	: Météorologie Nationale Française
SAPPORO	: Département de physiologie de l'Université de Hokkaido (Japon).
KUMAMOTO	: Département de physiologie et d'hygiène de l'Université (Japon).

Il s'y ajoute actuellement une station à la BASE ROI BAUDOUIN DES EXPEDITIONS BELGES DANS L'ANT-ARCTIQUE.

Les deux voeux suivants ont conclu la réunion:

1. Création d'un laboratoire central d'étude des "Tests chimiques" en vue de la recherche, de la mise au point et de la standardisation des conditions opératoires les plus appropriées à l'étude des différents facteurs physiques en cause.
2. L'organisation systématique dans les observatoires ou instituts météorologiques adéquats, d'observations régulières au moyen des tests les plus faciles, soit à des heures déterminées, soit continûment.

Florence, 7 octobre 1959

MISCELLANEOUS BIOCLIMATOLOGICAL DATA

Section B: Symposia of National Societies

THE EFFECTS OF POLLUTION ON LIVING MATERIAL

by

Dr. J.L. Cloudsley - Thompson (Gr. Britain) *

The subject of the Institute of Biology's symposium for 1958 was one of interest to many members of the International Society of Bioclimatology and Biometeorology. Held in the Lecture Hall of the Royal Geographical Society, London, on 25-26 September 1958, the symposium comprised fourteen papers on various biological aspects of pollution. The first six of these were concerned with various aspects of water pollution and therefore do not come within the scope of this Journal; but most of the remainder dealt with the effects of air pollution on living organisms.

Hazards and nuisances of air pollution from the burning of coal have been recognised in varying degree since the thirteenth century, but in spite of technological improvements the basic problems of preventing the emission of smoke and sulphur fumes and of obviating the damage they cause, have remained unsolved. As Dr. E.T. Wilkins (Department of Scientific and Industrial Research) pointed out in a paper on chemical and physical effects of air pollution on living material, the use of coal in Britain at the present time amounts to about 200 million tons annually and some 20 million tons of oil are also consumed. The emissions of dust smoke and sulphur dioxide resulting from the use of these fuels have been estimated at 0.8, 2 and 5 million tons a year respectively. In addition, the concentration of carbon monoxide in petrol-engine exhausts is several hundred times the toxic limit, although the rate of dilution after leaving the exhaust pipe is extremely rapid. Drivers and passengers of cars and other vehicles may inhale as much as 130 p.p.m. of carbon monoxide, if not more, the effect of which might well be to slow the driver's mental reactions and make him more accident-prone. With regard to "smog", it has been shown that the daily mortality follows closely and with little time lag, the daily changes in pollution as measured by smoke and sulphur dioxide.

Dr. J.K.A. Bleasdale outlined the results of pollution on plant growth and described experiments showing the effects of pollution in reducing the effective light period per day. Tarry deposits on leaves further reduce the light available and gaseous pollutants also have these effects. Sulphur compounds play an important part in regulating cell division and Dr. Bleasdale suggested that they might effect the rates of litter-production and leaf-formation by ryegrass.

The effects of atmospheric pollution on man were described by D.P.J. Lowther (Medical Research Council), who emphasised that the bewildering diversity and contradictions to be found in the vast literature relating to air pollution should not be allowed to diminish public anger and disgust at the tragic effects of the use of the air as a sewer.

In Britain, chronic fluorine poisoning or "fluorosis" occurs almost exclusively in animals in industrial areas, said Dr. Ruth Allcroft and arises from aerial contamination of herbage by the emission of fluorine-containing gases and dusts from industrial plants. The total economic loss occasioned is not great compared with that caused by other diseases, but is a matter of serious concern to farmers whose stock are affected. Dr. Allcroft illustrated her lecture with a film showing fluorine poisoning in cattle. The effects of this appeared to cause such acute agony and suffering to the wretched animals that surprise was expressed by members of the audience that such a disgraceful state of affairs should be tolerated.

The uptake of radioactive fission products by plants was discussed by Dr. H.J.M. Bowen, whilst their effect on aquatic organisms formed the subject of a paper by Mr. W.L. Templeton. The writer of this review was most impressed by the care applied by members of the Research and Development Branch of the United Kingdom Atomic Energy Authority to the establishment of safe levels of radioactivity in organisms that directly effect man.

Finally, a challenging and stimulating contribution by Dr. T.C. Carter (Medical Research Council's Radiobiological Research Unit) stressed the need for new experimental data on the nature of the radiation dose/mutation relationship in low dose ranges, in order that minimal criteria may be determined. At present this relationship is completely unknown in the case of mammals. Yet, if it is other than linear, the whole logical structure supporting present-day assessments of the genetical hazard to man of nuclear and allied radiations, collapses like a house of cards.

As in many previous Symposia of the Institute of Biology, the unfortunate dichotomy between purely scientific, economic and political motives was manifest. One could not help reflecting that the real difficulties are probably social and financial rather than biological, and present problems of ethics rather than of science.

The proceedings of the symposium will be published by the Institute of Biology, 41, Queen's Gate, London, S.W. 7.

* Dept. of Zoology, Kings Coll., Univ. of London, Strand, London W.C. 2, Gr. Britain.

M I S C E L L A N E O U S B I O C L I M A T O L O G I C A L D A T A

Section B: Reports of Congresses of National Bioclimatological Societies

REPORT ON THE BIOCLIMATOLOGICAL CONFERENCE IN CIECHOCINEK (Poland) *
(30 - 31 October 1958)

A Bioclimatological Conference was held in Ciechocinek-Spa; 30 - 31 October 1958, during the 3rd Polish Congress of Balneology. The purpose of the Conference was to discuss the problems of bioclimatology in towns, industrial centres and health-resorts; and St. Zych delivered a paper on this subject.

Papers of a more general character were presented by St. Rózański - on the need for a climatic classification of health-resort areas and by S. Tyczka - on the problems on bioclimatological research and methods, mentioning the latest world publications.

The climate of towns was discussed in one paper only - The Bioclimate of Łódź - by St. Rózański, M. Tarajkowska and St. Zych. Various methods of research carried out in the area of this town were concerned first with constant gaseous atmospheric pollution and its noxious influence on human health and secondly on the flora of the region. The research made it possible for the climate of the town to be classified approximatively and areas convenient for flat-, service- and industrial buildings to be designated.

The problems of climate in health-resorts and their surrounding regions and of the influence of climate on man were discussed in the majority of the papers. W. Wójtowicz presented an account of the climate of the Kłodzko-Valley. Based on classical climatological materials over a 50 years' period, the author tried to give a bioclimatological interpretation of the climatic relations of the region. Five climatic zones were differentiated differing distinctly from each other in regard to the value and dynamics of their climatic factors as well as in their importance in therapy. The indication of vertical increase in precipitation allowed areas of relative tranquillity to be excluded. In his paper "The local climate of Polanica-Spa (Kłodzko Valley)", in addition to a short outline of the climate, M. Kołodziejek discussed the local differences of microclimate based on data from the local meteorological station from 1950 until 1957 as well as on regional measurements. Microclimatic differences denoted the health-resort areas most advantageous for future building and those requiring amelioration of the climate.

The paper by J. Cieślak on the duration of sunshine in Polanica-Spa and Kudowa-Spa in the years 1951 - 1957 is a valuable supplement to the climatology of the Kłodzko Valley. T. Szczepańska presented the results of research on the ozone content of the air close to the earth's surface in the area of Ciechocinek-Spa and A. Gurba the results of research on cooling-power in the same Spa on the base of catathermometric measurements carried out at the meteorological station during a five-year period.

S. Liman described the result of microclimatic research in galleries and cavities of the salt-mine at Inowrocław, a possible site for the future treatment of asthmatic patients. The differences in temperature and humidity as well as in air movement are rather great, depending on distance from the shaft, on the presence of water and on the working of ventilators.

M. Kwaśniewska-Błaszczyk in her paper "The Groer-Hecht reaction in children during health resort treatment on the Baltic seashore (Kołobrzeg-Spa)" and J. Jez and H. Chobot in theirs "On the behaviour of the bacterial flora in the nasopharynx of children under the influence of thalassotherapy" tried to explain the influence of climatic conditions on the phenomena.

W. Gądzikiewicz considered air movement in some Polish health-resorts as an indispensable factor in denoting the "effective temperatures". In the papers of B. Kiełczewski and J. Bogucki the results are given of preliminary research on the influence of cosmo-meteorological factors on some physiological functions of the organism in rest and after activity tests.

Physical bioclimatology was discussed in several papers. J. Szymka delivered one on atmospheric extinction in Wrocław during the years 1950-1955. The author reached the conclusion that in an annual cycle the reduction in solar radiation in the atmospheric layer close to the earth's surface - even in the big towns - is insignificant in comparison with its reduction in the whole atmosphere. M. Kluge discussed the results of research on the influence of atmospheric pollution

* Report prepared by mgr Władysław Wójtowicz, Instytut Balneoklimatyczny, Poznań, Poland.

on the reduction in direct solar radiation in the industrial centre of High Silesia. J. Paszyński referred to the results of research on the transparency of the atmosphere carried out during a sea voyage from Vietnam to Poland. M. Cena and J. Słomka presented a map of bioclimatological dry and humid cooling power in Poland calculated with the empirical formulae of Hill and Lehmann on the basis of data from 50 Polish meteorological stations. The authors compared the results with those obtained by other authors in other areas and discussed from a biological and climatological point of view the value of the results obtained. W. Bogusławski presented insolation tables for Poland and discussed some biophysical aspects of colour vision.

TITLES OF PAPERS READ AT THE BIOCLIMATIC CONFERENCE IN CIECHOCINEK *

GENERAL THEMES:

1. St. ZYCH - Research into the bioclimates of towns and industrial centres and health resort treatment.
2. St. RÓŻAŃSKI - The necessity for climatic classification in health resort's areas.
3. S. TYCZKA - The problems, research methods and perspectives of development in human bioclimatology.

BIOCLIMATOLOGY OF TOWNS AND OF HEALTH RESORTS:

4. St. RÓŻAŃSKI
M. TARAJKOWSKA - The bioclimate of Łódź.
- St. ZYCH
5. W. WÓJTOWICZ - The climate of the Kłodzko valley.
6. M. KOŹDZIEJEK - The local climate of Polanica-Spa.
7. L. CIESLAK - The duration of sunshine in the spas of Kudowa and Polanica.
8. T. SZCZESNA - Research on the ozone content of the atmospheric layer close to the earth's surface in the area of Ciechocinek-Spa.
9. A. GURBA - Catatermometric cooling power in Ciechocinek-Spa.
10. W. GĄDZIKIEWICZ - Air movements and wind in Spas and Rest centres in Poland.
11. S. LIMAN - Microclimatic research in galleries and cavities of the salt - mine Solno I in Inowrocław.

MEDICAL BIOCLIMATOLOGY:

12. M. KWAŚNIEWSKA-BŁASZCZYK - The Groer-Hecht reaction in children during health-resort treatment on the Baltic Seas.
13. H. CHOBOT - On the behaviour of the bacterial flora in the naso-pharynx of children under the influence of thalassotherapy.
14. B. KIEŁCZEWSKI - The results of introductory ecological research on the influence of cosmo-meteorological factors on some physiological functions of human organism in rest and after activity tests.
15. J. BOGUCKI - Research on changes in physiological functions dependent on cosmo-meteorological factors based on research carried out in training camps of the Higher School for Physical Education, Poznań.

PHYSICAL BIOCLIMATOLOGY:

16. M. CENA
J. SŁOMKA - The bioclimatic cooling power in Poland.
17. J. SŁOMKA - Atmospheric extinction in Wrocław during the years 1950-55 (bioclimatic aspects).
18. M. KLUGE - The influence of atmospheric pollution on the effect of solar radiation in a High Silesian Industrial Centre.
19. J. PASZYŃSKI - Research into atmospheric transparency during oceanic voyage; introductory results of direct measurements of solar radiation.
20. W. BOGUSŁAWSKI - The iso-insolar table indexes of Poland.
21. W. BOGUSŁAWSKI - On some biophysical aspects of colour vision, decomposition of solar-radiation and colour vision.

* Compiled by mgr. Stefan Liman, Instytut Balneoklimatyczny, Poznań, Poland.

MISCELLANEOUS BIOCLIMATOLOGICAL DATA
Section B: Reports of Congresses of national bioclimatological societies

PROGRAM OF THE BIOCLIMATOLOGICAL CONFERENCE AT LIBLICE (CZECHOSLOVAKIA)
(3-5 November 1958) *

- PICKO, V. (Hygienisches Institut, Praha): Singularität und Morbidität (Unsere Erkenntnisse vom Einfluss und Manifestation der Singularität auf die Entwicklung und den Verlauf der Morbidität).
- TROMP, S.W. (Bioclimatological Research Centre, Leiden): Review of Bioclimatological studies carried out at the Bioclimatological Research Centre at Leiden (Netherlands).
- BARCAL, R. and BOBEK, K. (Interne Klinik der Medizinischen Fakultät in Plzen): Meteorotropismus plötzlicher Herz- und Gefässerkrankungen.
- KWASNIEWSKA-BASZCYK, M. (Instytut balneoklimatyczny, Poznań): Groer-Hecht's Reaktion bei Kindern während einer Sanatoriumskur an der Ostseeküste.
- NOVÁK, J. (I.Dermatologische Klinik der Medizinischen Fakultät in Praha): Periodizität der Entstehung und des Verlaufs einiger Krankheiten.
- SVORAD, D. and WELLNEROVÁ, J. (Physiologisches Institut der ČSAW, Praha): Einfluss einiger meteorologischer Faktoren auf den Rhythmus des Schlafes und des Wachens mit besonderer Berücksichtigung der Schlaflosigkeit bei Tieren.
- CUPCEA, S. and DELEANU, M. (Institut de igiena, Cluj): Experimentelle und klinische Forschungen über einige Einflüsse der Luftionen auf den Organismus.
- NOVÁK, J. (I.Dermatologische Klinik der Medizinischen Fakultät in Praha): Einfluss der Veränderungen des Geomagnetismus auf die Entstehung und den Verlauf einiger Krankheiten.
- TICHÝ, S. (Klinisches otorhinolaryngologisches Laboratorium der ČSAW, Praha): Einfluss der äusseren Umwelt auf das Krankenhaus-Mikroklima.
- BERJUSCHEW, K.G. (Institut für die Vervollkommnung von Aerzten, Moskau): Einfluss der Grünanlagen auf die Verbesserung der mikroklimatischen und hygienischen Bedingungen in den Siedlungen nach den Angaben der Forschungsarbeiten sowjetischer Hygieniker (Russisch)
- KAZDOVÁ, PhMr (Hygienisch-epidemiologische Bezirksdienststelle) and HAVLÍČEK, V. (Landwirtschaftliche Hochschule in Brno): Einfluss industrieller Exhalationen und meteorologischer Elemente auf die örtliche Verteilung des Staubfalles in Siedlungen.
- BÖHM, B. (Hydrometeorologische Anstalt, Praha): Antrag zur Vereinheitlichung der Methodik der katathermometrischen Messung.
- CHOBOT, H. and JEZOWA, L. (Instytut balneoklimatyczny, Poznań): Verhalten der Nasen-Rachen Flora bei Kindern nach einer Ostseebadkur.
- CERMÁK, K. (Forschungsanstalt für Forstwirtschaft, Zbraslav-Stirady): Einfluss der Witterung auf die Unfallshäufigkeit in der forstlichen Produktion.
- LIMAN, S. (Instytut balneoklimatyczny, Poznań): Mikroklimatische Untersuchungen in den Stollen und Kammern des Salzbergwerks von Inowrocław.
- VESELÝ, E. (Forschungsinstitut für Balneologie, Praha): Erwärmung im Wasser.
- WELLNEROVÁ, J. and SVORAD, D. (Physiologische Anstalt der ČSAW): Interindividuelle Veränderungen des emotionalen Verhaltens durch den Einfluss der Temperatur und Luftfeuchtigkeit.
- LIMAN, S. (Instytut balneoklimatyczny, Poznań): Ueber den Stand polnischer Forschungsarbeiten auf dem Gebiete der Bioklimatologie des Menschen.
- CUPCEA, S. and DELEANU, M. (Institutul de igiena, Cluj): Einige Probleme der geographischen Verteilung der Luftionisation in der Rumänischen Volksrepublik.

* Information concerning reports etc. can be obtained from Ing. V. Krečmer: Československa Akad. ved, 34 Zborovská, Praha 16, Czechoslovakia.

- PICHA, J. (Hydrometeorologisches Institut, Observatorium Neuköniggrätz (Nový Hradec Králové)) : Erfahrungen bei der Messung des bodennahen Ozons.
- STRUŽKA, V. (Anstalt für Hygiene, Praha): Methoden von Geländeuntersuchungen in bioklimatischer Richtung.
- MADE, A. (Forschungsinstitut für Agrarmeteorologie, Halle/Saale): Bericht über einige Erfahrungen auf dem Gebiete der meteorologischen Geländevermessung.
- KLEČKA, M. and STRÁDAL, V. (Forschungsinstitut für Agrarökonomik der ČSAL in Praha): Praktische Ausnützung der landwirtschaftlichen Bioklimatologie bei der Produktionsplanung in der Tschechoslowakei.
- HLAVÁČ, V. (Institut für Gebietsplanung, Praha): Beitrag zur Methodik der Standortverteilung der landwirtschaftlichen Produktion vom bioklimatologischen Gesichtspunkt (am Beispiel des Bezirkes České Budějovice).
- HESSE, W. (Institut für Agrarmeteorologie d. Karl-Marx-Universität, Leipzig): Ueber die Methodik der Bestandesklimamessungen.
- BALCAR, J. (Forschungsinstitut für Tabakindustrie, Báb): Einfluss des Klimas und des Bodens auf die Tabakpflanze in der Vietnamesischen demokratischen Republik.
- BERÉNYI, D. (Universität Debrecen): Ueber das Bestandesklima des Maises.
- KURPELOVÁ, M. (Hydrometeorologisches Institut, Bratislava): Beitrag zum Wachstum und zur Entwicklung einiger Fruchtarten im Zusammenhang mit der Witterung.
- ŠKULA, K. (Forschungsinstitut für Tabakindustrie, Báb): Einfluss der Witterung auf die Dauer der Wachstumsphasen beim Tabak.
- PEJML, K. (Hydrometeorologisches Institut Observatorium Doksany a/E): Beitrag zum Bestandesklima der Kartoffeln vom Gesichtspunkt der Entstehung der Kartoffelfäule.
- BERÉNYI, D. (Universität Debrecen): Ueber die in Weinkellern und Höhlen angewandten Beobachtungsmethoden.
- MAJERNÍK, O. (Biologisches Institut der SAW, Bratislava): Wirkung klimatischer Faktoren auf einige phyto gene Organismen im Zusammenhang mit ihrer Schädlichkeit für die Weinrebe und Aprikose.
- FOLTÝN, O. (Laboratorium für Pflanzenschutz der SAW, Ivanka pri Dunaji): Klimatische Einflüsse auf die Entstehung und Entwicklung der Perenospora der Weinrebe.
- KREČMER, V. and IZERA, V. (Forschungsanstalt für Forstwirtschaft, Zbraslav-Strnady): Neuer Typ eines Apparates zum Messen des täglichen Sonnenbogens.
- KESNER, B. (Laboratorium für Agrometeorologie der ČSAL, Praha): Neue Methode der Ermittlung der erforderlichen Lüftungintensität abgeschlossener Räume nach der CO₂ - Konzentration.
- SLAVÍK, B. (Biologisches Institut der ČSAW, Praha): Requivalente Verdunstung, ihr Höhengradient und seine ökologische Bedeutung.
- VELIKOVSKÝ, V. (Forschungsinstitut für Getreide, Kroměříž) and KRHOUNEK, V. (Hydrometeorologisches Institut, Praha): Ergebnisse der Messung in Winterroggenbeständen bei der Ermittlung der zwischen einigen klimatischen Faktoren und der Entwicklung des Schneeschimmels bestehenden Beziehung.
- TREFNÁ, E. (Hydrometeorologisches Institut, Praha) and MAREČEK, F. (Forschungsinstitut für pflanzliche Produktion, Praha): Einige aus Beobachtungen des Mikroklimas im ungeheizten Gewächshaus gezogene Schlüsse.
- HAVLÍČEK, V. (Hochschule für Landwirtschaft, Brno): Mesoklimatische Besonderheiten von Stauseen und Wasserflächen in ihrem täglichen Gang.
- SLAVÍK, B. and CATSKÝ, J. (Biologisches Institut der ČSAW, Praha): Methode der Ermittlung der CO₂ - Konzentration in der Luft.
- PASÁK, V. (Forschungsinstitut für Meliorationen der ČSAL, Praha): Mikroklimatische Verhältnisse bei der Aufforstung von Sandböden.
- TOMANEK, J. (Szkoła gówna gospodarstwa wiejskiego, Warszawa): Gestaltung der mikroklimatischen Verhältnisse in verschiedenen Lochschlagbetriebstypen.
- VOLNÁ, M. (Forstliche Hochschule, Brno): Beurteilung von Nestpflanzung und -Saat vom Gesichtspunkt ihrer mikroklimatischen Wirkung.
- VOLNÝ, S. (Forstliche Hochschule, Brno): Einfluss des gemeinen Wacholders auf einige mikroklimatische Prozesse der erdnahen Luftschichten und der oberen Bodenhorizonte als Voraussetzung einer bedeckenden Komponente bei der bodenschützenden Aufforstung.

- INTRIBUS, R. (Forstliches Laboratorium der SAW, Zvolen): Beitrag zum Mikroklima von Oedland in der Slowakei.
- BAUMGARTNER, A. (Meteorologisches Institut der forstlichen Forschungsanstalt, München): Die klimatischen Standortsfaktoren an einem Berg des Böhmerwaldes.
- KRECMER, V. (Forschungsinstitut für Forstwirtschaft, Zbraslav-Strnady): Grösse der Hiebessel in Kieferbeständen in Niederungslagen vom Gesichtspunkt der Mikroklimatologie
- KORTÜM, F. (Forstliche Hochschule, Eberswalde): Die Verdunstung von Waldböden und Freilandflächen.
- ZACHAR, D. (Forstliches Laboratorium der SAW, Zvolen): Einfluss des Reliefs auf den Wasserhaushalt des Bodens.
- VINŠ, B. (Forschungsinstitut für Forstwirtschaft, Zbraslav-Strnady): Witterung und forstlicher Holzartenzuwachs.
- KŘIVSKÝ, V. (Astronomisches Institut der ČSAW, Ondřejov): Säkulärer Trend von Witterung und Zuwachs.
- CHALUPA, V. (Forschungsinstitut für Forstwirtschaft, Zbraslav-Strnady): Neue Methoden der forstlichen laubholzbestand Phänologie.
- SLAVÍK, B. (Biologisches Institut des ČSAW, Praha): Das Eindringen des Regenwassers in den Laubholzbestand.
- KVĚT, J. (Geobotanisches Laboratorium der ČSAW, Práhonice): Bisherige Erfahrungen mit der Integrationsmethode der Messung von Bodentemperaturen mittels der Sacharoseinversion.
- KREČMER, V. and FOJT, W. (Forschungsanstalt für Forstwirtschaft, Zbraslav-Strnady): Niederschlagsmesser für mikroklimatologische und ökologische Studien.
- HOLOVSKÝ, M. (Institut für Wald- und Jagdwesen der ČSAL, Praha): Funktionall ökonomische Grünanlagen in der Industrie.
- HOLOVSKÝ, M. (Institut für Wald- und Jagdwesen der ČSAL, Praha): Dynamische Modifikation der Waldschutzgürtel in der Industrie.
- MATĚJKA, V. and CHROUST, M. (Forstliche Fakultät, ČVUT, Technische Hochschule, Praha): Mikroklimatische Forschungsarbeit auf dem Gebiete von Trockenkammern grundlegenden Typen in den Grossklimaanstalten.

M I S C E L L A N E O U S B I O C L I M A T O L O G I C A L D A T A
Section B: Reports of Congresses of national bioclimatological societies

PROGRAM BIOCLIMATOLOGICAL CONFERENCE ON CLIMATOLOGICAL AND
BALNEOLOGICAL HEALTH RESORTS AT E.BERLIN, D.D.R. (3-5 MARCH 1959) *

KURORTKLIMA

- | | |
|-----------------------------|---|
| W. BOËR (Potsdam) | : Zum Begriff des Lokalklimas |
| H. ZENKER (Bad Berka) | : Die Bedeutung des Lokalklimas von Kuranlagen - am Beispiel einer Tuberkulose-Heilstätte |
| W. WÓJTOWICZ (Posen) | : Die Auswertung der klassischen Klimadaten in der Kurortklimatologie - am Beispiel des Berglandes von Kłodzko |
| W. UNDT (Wien) | : Bioklimatische Arbeitsmethoden |
| G. HENTSCHEL (Berlin) | : Bioklimatische Arbeitsmethoden |
| L. TRAUNER (Zagreb) | : Klimakurort - Kurortklima |
| H. PFLEIDERER | : Bedeutung und Technik der Frigorigraphie |
| H. GRIMM (Berlin) | : Das Strandleben an der Ostseeküste in Abhängigkeit von Wärmestrahlung und Abkühlungsgrösse |
| H.P. MOCHMANN (Rostock) | : Leptospirosen, ein Beitrag zur Lokalbioklimatologie |
| A.v. KNORRE (Oschersleben) | : Ueber Asthmafragen |
| G. HENTSCHEL (Berlin) | : Untersuchungsergebnisse der Sterblichkeit unter verschiedenen lokalen Gegebenheiten |
| I. KÉRDÖ (Budapest) | : Komplexe Betrachtungsweise bei der Beurteilung von biologischen Wirkungen klimatischer Umweltfaktoren |
| K. MENYHÉRT (Budapest) | : Biometeorologischer Dienst und Forschung in Ungarn |
| H. MROSE (Dresden) | : Untersuchungsergebnisse über reduzierende Spurenstoffe in der Atmosphäre |
| U. JESSEL (Westerland/Sylt) | : Stickoxyde als Ursache von Fehlern in der chemischen Ozonometrie ? |
| F. PFEIFER (Leipzig) | : Die festen Schwebstoffe in der Atmosphäre und ihre natürliche Radioaktivität |
| R. NOACK (Berlin) | : Die lufthygienische Situation in einigen Kurorten und Erholungszentren der DDR sowie Hinweise für Neuanlagen von Kurzentren |
| L. RASSOW (Halle) | : Zum Problembereich Lufthygiene |
| J. KOLBIG (Potsdam) | : Abschätzung der Ausbreitung von Luftverunreinigungen |
| K. SEIFERT (Berlin) | : Staatliche Massnahmen zum Schutze des Klimas in Kurorten |
| F.E. CARL (Berlin) | : Bioklimatologie und Erholungsplanung |

KLIMATHERAPIE

- | | |
|--------------------|---|
| A. KUKOWKA (Greiz) | : Zum Problem der Luftelektrizität |
| PICHOTKA (Berlin) | : Die Abhängigkeit der Sauerstoffaufnahme von der Sauerstoffspannung, insbesondere bei geringen Erniedrigungen des Sauer- |

* Information concerning reports etc. can be obtained from Dr. Ortmeyer: Forschungsinstitut für Bioklimatologie, Hufeland-Krankenhaus, Haus 1, Karowerstrasse 11, Berlin-Buch, DDR.

stoffpartialdrucks

- | | |
|-------------------------------|---|
| K. LÜHR (Dresden) | : Objektivierung des Kurerfolges bzw. Kureffektes |
| E.G. SCHULTZE (Wyk auf Föhr) | : Ueber die Objektivierung der Kurergebnisse an der See |
| SCHIRGEL (Heiligendamm) | : Objektivierung des Gesamtkureffektes im Seeheilbad mit dem Leistungsprüfgerät nach Prof. Böhlau |
| SCHIRGEL (Heiligendamm) | : Akklimatisationsstudien im Seeheilbad Heiligendamm |
| H. WAGNER (Bad Elster) | : Die Beeinflussung des Organismus im Kurort - nachgewiesen am Verlauf der kollektiven Körpertemperatur |
| M. KALINOWSKI (Posen) | : Einfluss der Kurortbehandlung auf die Erhaltung der Leistungsfähigkeit alternder Menschen. |
| E.v. PHILIPSBORN (Oberstdorf) | : Indikationen für eine Klimakur im Gebirge |
| I. ILIJEW (Sofia) | : Physiologische Bemerkungen zur Methodik des Sporttrainings im Gebirge während der Akklimatisation |
| K. LINSER (Berlin) | : Die Klimatherapie bei Hautkrankheiten |
| F. FUCHS (Karl-Marx-Stadt) | : Klimatherapeutische Möglichkeiten für konstitutionelle Neurodermatiker auf dem Fichtelberg |
| K. HARNACK (Berlin) | : Auswertung unserer klimatischen Heilkuren bei Hautkranken |

M I S C E L L A N E O U S B I O C L I M A T O L O G I C A L D A T A
Section B: Reports of Congresses of national bioclimatological societies

"PROBLEME DER BIOKLIMATOLOGIE UND DES STALLKLIMAS
BEI DER RINDER- UND SCHWEINEHALTUNG"
(Program of Symposium to be held at Pössneck, Thüringen, D.D.R.,
on 28 and 29 May 1959)*

PROBLEME DER BIOKLIMATOLOGIE

Uebersichtsreferat

Die Bioklimatologie als Hilfsmittel der modernen Haltungsforschung (Methoden und Geräte, Ziele).

Die Konstitutionsforschung und die Bioklimatologie.

Die Veterinärhygiene und die Bioklimatologie in der Haustierhaltung.

Das Strahlungsklima und sein Einfluss auf die landwirtschaftlichen Nutztiere.

Probleme der Tierhaltung auf bioklimatischer Grundlage unter den Bedingungen des Kontinentalklimas.

Offene Fragen des Tierzüchters an die Haltung und die Bioklimatologie der Tiere unter mittteleuropäischen Verhältnissen.

ERGEBNISSE DER HALTUNGSFORSCHUNG BEI RINDERN UND SCHWEINEN

In der Tschechoslowakei

In Ungarn

In Dänemark

In Belgien

In der Bundesrepublik Deutschland

In der Deutschen Demokratischen Republik

STALLKLIMA UND STALLHYGIENE

Die physiologischen Grundlagen für die Berechnung des Stallklimas.

Wärmewirtschaftliche Unterlagen für die Berechnung des Stallklimas.

Welche Darstellungsform des Stallklimas ist notwendig ?

Ueber eine neue Berechnungsgrundlage des Stallklimas.

Welche Lüftungssysteme sind im Stallbau anzuwenden ?

Fragen der Baustoffprüfung.

Ergebnisse von Baustoffprüfungen in Schweden.

* Information concerning the Symposium can be obtained from Dr. Scholz, Forschungsstelle für Tierhaltung Knau, Knau (über Triptis i. Thür., Kreis Pöszneck).

Section C : Bioclimatological stations and institutions

MISCELLANEOUS BIOCLIMATOLOGICAL DATA

Section C: Bioclimatological stations and institutions

DIE BIOKLIMATOLOGIE DES MENSCHEN IN POLEN

von

Mgr. St. Liman (Poland)*

I. DIE POLNISCHE BIOKLIMATOLOGIE DES MENSCHEN
IN DER ZWISCHENKRIEGSZEIT

Auch in Polen sind seit einigen Jahren Probleme der Bioklimatologie des Menschen zum Gegenstand eingehender Forschungen geworden. Sie erregten das Interesse unserer Meteorologen, Klimatologen und Aerzte – und ganz besonders der Kurärzte. Es ist daher kein Zufall, wenn wir unter den Pionieren der polnischen Bioklimatologie nicht nur Namen von Geophysikern, sondern auch von Aerzten finden. Von den Bahnbrechern der gegenwärtigen Bioklimatologie des Menschen, deren wesentliche Entwicklung in die dreissiger Jahre unseres Jahrhunderts fiel, sind folgende Aerzte zu nennen: L. Korczyński, A. Sabatowski und M. Michałowicz. Ihren wissenschaftlichen Arbeiten auf dem Gebiete der medizinischen Bioklimatologie ist es zu verdanken, dass für diese Probleme ausser den Aerzten auch die Naturwissenschaftler gewonnen wurden. Unter den Meteorologen dagegen gebührt die grösste Anerkennung W. Gorczyński und E. Stenz. Ihre Forschungsarbeiten auf dem Gebiete der Aktinometrie führten zu einer genaueren Kenntnis des Solarklimas nicht nur Polens, sondern auch anderer Länder. Ein weiteres Verdienst von Stenz ist die Errichtung der ersten biometeorologischen Forschungsstation, die auf dem Kasprowy Wierch (polnische Hohe Tatra) im Jahre 1938 ihre Arbeit begann. Es wurden dort folgende Elemente des Bioklimas untersucht:

1. die Sonnenstrahlung unter Berücksichtigung des Sonnenspektrums im sichtbaren Lichte, Infrarot und Ultraviolett;
2. die atmosphärische Luftelektrizität (mit Israels Meszgerät);
3. die biologische Abkühlungsgrösze (mit dem Frigorimeter nach Thilenius-Dorno).

Das Arbeitsprogramm dieser Station berücksichtigte auch meteoro-physiologische Probleme, die der Hochgebirgs-Klimatherapie dienen. Die ersten Ergebnisse der vielversprechenden Forschungsarbeiten auf dem Kasprowy Wierch wurden leider durch den Ausbruch des zweiten Weltkrieges nebst seinen Folgen vernichtet.

Bei einer Bewertung des Standes der polnischen Forschung auf dem Gebiete der Bioklimatologie des Menschen in der Zwischenkriegszeit muss man feststellen, dass wir zwar im Vergleich mit den westlichen Ländern im Rückstand waren, jedoch diesen Wissenszweig entsprechend den damals gegebenen technischen und finanziellen Möglichkeiten entwickelten. Die polnische Bioklimatologie dieser Zeit kann zwar keine grössere Anzahl von wissenschaftlich-experimentellen Arbeiten aufweisen – erstens hauptsächlich deshalb, weil die Unterstützung gut eingerichteter Forschungsstellen fehlte – und zweitens, weil die Probleme, mit denen sich die bestehenden Arbeitskreise beschäftigten, durch eine gewisse Zufälligkeit bestimmt waren. Als eine bedeutende Errungenschaft dürfen jedoch zahlreiche Beiträge und populär-anschauliche Arbeiten angesehen werden. Auch die Herausgabe des von Korczyński verfassten zweibändigen 872 Seiten umfassenden Lehrbuches der medizinischen Klimatologie (heute schon teilweise veraltet) ist ohne Zweifel eine namhafte Errungenschaft der polnischen Bioklimatologie der Zwischenkriegszeit.

* Aus dem Forschungs-Institut für Balneologie und Klimatologie in Poznań (Poland); Direktor: Dr. J. Jankowiak.

II. DIE ENTWICKLUNG DER BIOKLIMATOLOGIE DES MENSCHEN IN DER NACHKRIEGSZEIT

A. ORGANISATION.

Der letzte Weltkrieg bedeutete eine Zwangspause in den Arbeiten unserer Bioklimatologen und verursachte einen grossen Verlust von den ohnehin nicht zahlreichen Fachleuten, Meteorologen, Klimatologen und Aerzten, die alle an der Entwicklung der Bioklimatologie des Menschen gearbeitet hatten. Daher bemühten sich unsere Wissenschaftler in den ersten Nachkriegsjahren vor allem um die Ausbildung eines jungen Nachwuchses. Im Vergleich mit der Vorkriegszeit ist das Interesse an bioklimatologischen Problemen nach dem Jahre 1945 in Volkspolen bemerkenswert gestiegen. Ohne Zweifel hängt dies mit dem intensiv fortschreitenden Umbau der wirtschaftlichen Struktur unseres Landes zusammen, den wiederum die Entwicklung unserer Industrie und speziell der grundlegenden Rohstoffbasis des Bergbaus und der Hüttenindustrie bewirkt hat. Aus diesem Grunde wurden vorzugsweise auf weite Sicht Pläne einer Raumbewirtschaftung von Städten und Siedlungen in Arbeit genommen. Da diese hauptsächlich auf einer Analyse der lokalen klimatischen Verhältnisse und der Lufthygiene basierten, erforderten sie die Zusammenarbeit mit Klimatologen. Gleichzeitig mit der fortschreitenden Industrialisierung machte sich eine Zunahme von Berufskrankheiten bemerkbar; dies verursachte zweifellos ein regeres Interesse an den Problemen der Sanitätshygiene und der medizinischen Bioklimatologie und trug zur Entwicklung der letzteren bei.

Von den augenblicklich auf dem Gebiete der medizinischen Bioklimatologie tätigen Arbeitskreisen kann man erwähnen:

1. die mikroklimatologische Forschungsstelle der Abteilung für Stadthygiene am Staatlichen Hygiene-Institut in Warszawa, die sich hauptsächlich mit sanitäts-hygienischen Problemen der Städte befasst;
2. die bioklimatologische Abteilung am Institut für Balneologie und Klimatologie in Poznań, die seit dem Jahre 1952 besteht und Probleme der medizinischen Bioklimatologie - vor allem der Kurortbioklimatologie - bearbeitet. Beide Stellen sind dem Gesundheitsministerium untergeordnet.
3. Als dritte Forschungsstelle ist die biometeorologische Abteilung am Staatlichen Hydro-Meteorologischen Institut in Warszawa zu verzeichnen. Ihr Forschungsprogramm umfasst ausgewählte Probleme der Meteorologie und der medizinischen Klimatologie.

Ausser den besprochenen drei Arbeitskreisen werden von einigen Lehrstühlen für Hygiene an den Medizinischen Akademien bzw. meteorologischen und klimatologischen Instituten der Universitäten neben dem allgemeinen Forschungsplan gewisse Probleme der medizinischen Bioklimatologie bearbeitet.

B. DIE INSTRUMENTELLE AUSSTATTUNG DER FORSCHUNGSSTELLEN.

Die in biometeorologischer Hinsicht am besten eingerichteten Beobachtungsstellen besitzt das Staatliche Hydro-Meteorologische Institut. Die biometeorologische Abteilung des obengenannten Instituts hat sowohl ein Aussenbeobachtungsnetz wie auch eine Zentralstation in Warszawa. Es werden dort vor allem das Solarklima (Sonnenstrahlung, Sonnenscheindauer), die Luftabkühlungsgrösze, sowie das Bioklima der Städte und der Industriezentren erforscht. Es ist auch bemerkenswert, dass der obengenannten Forschungsstelle des gesamte meteorologische Archivmaterial jederzeit zugänglich ist.

Das Institut für Balneologie und Klimatologie verfügt bisher über keine ständige biometeorologische Beobachtungsstelle. Das Institut ist jedoch im Begriff neue biometeorologische Forschungsstellen in Kurorten verschiedener physiographischer Regionen unseres Landes zu errichten. Für die Gebirgslandschaft sollten diese in Krynica bzw. Szczawnica und in Zakopane (Hohe Tatra) entstehen; für die mittelpolnische Tiefebene im Kurort Inowrocław und für die Küstenlandschaft im Ostseebad Kołobrzeg. Diese Beobachtungsstellen, die mit klima-physiologischen Laboratorien ausgestattet werden müssen, würden als Forschungsstellen für das Kurortklima jener Gegenden dem Institut für Balneologie und Klimatologie untergeordnet sein. Bis dahin ist das Institut gezwungen, in eigenen Forschungsarbeiten an ausgewählten Elementen des Kurortklimas sich vorwiegend auf stichprobenartige Geländebeobachtungen zu stützen. Augenblicklich werden seitens des Instituts für Balneologie und Klimatologie dauernde Messungen der Intensität der Sonnenstrahlung in Inowrocław, sowie Messungen der Sonnenscheindauer und Abkühlungsgrösze der Luft in zwei Kurorten Inowrocław und Rymanów-Zdrój, durchgeführt.

Von einer Gesamtzahl von 28 Kurorten, die der zentralen Kurortverwaltung unterstellt sind, besitzen 23 Kurorte meteorologische Stationen. Davon bestehen meteorologische Stationen I. Ordnung in 2 Kurorten, II. Ordnung in 6 und III. Ordnung in 15 Kurorten. Alle diese Beobachtungsstellen unterstehen dem Staatlichen Hydro-Meteorologischen Institut.

III. DIE FORSCHUNGSRICHTUNG UND ANGEWANDTE ARBEITSMETHODEN AUF DEM GEBIETE DER BIOKLIMATOLOGIE DES MENSCHEN

Die Forschungsarbeiten auf dem Gebiete der Bioklimatologie des Menschen gehen in zwei Richtungen: A. der biologischen, B. der physikalischen.

A. DIE BIOLOGISCHE RICHTUNG.

Ähnlich wie in anderen Ländern, gehen auch unsere meteoroklimabiologischen Forschungen in Richtung einer Korrelation der biologischen Prozesse des menschlichen Organismus mit den Wetterprozessen. Als Gegenstand speziellen Interesses sind meteoropathologische Probleme, alsdann der Jahreszeitrhythmus gewisser Krankheiten (z.B. Kreislaufstörungen, Krankheiten der Luftwege, Bewegungsstörungen und andere) zu verzeichnen. In dieser Richtung gehen eben auch die Arbeiten der bioklimatologischen Abteilung des Instituts für Balneologie und Klimatologie, sowie der biometeorologischen Abteilung des Staatlichen Hydro-Meteorologischen Institut.

Die klima-biologischen, bzw. meteoroblogischen Arbeiten werden bei uns mit den allgemein bekannten und angewandten Methoden durchgeführt; nämlich mit Hilfe der statistischen, experimentellen und dynamischen Methode.

Zur Vervollständigung eines allgemeinen Bildes der Thematik der durchgeführten Forschungen kann man beispielsweise einige der bearbeiteten Probleme anführen. So befasste man sich in den letzten Jahren mit dem Problem der Wärmeregulation des kindlichen Organismus in Beziehung zu balneoklimatischen Reizen, weiter mit biologischen und pharmakodynamischen Reaktionen, ebenfalls bei Kindern, bei denen die thalasso-therapeutische Heilmethode angewandt wurde.

Andere Arbeiten auf dem Gebiete der Meteoropathologie befassten sich mit Problemen, wie z.B. dem Einfluss des Wetters auf rheumatische Schmerzzustände, auf Erkrankungen überhaupt, und auf die Letalität.

KLIMATHERAPIE

In der klima-biologischen Problematik muss auch die Klimatherapie berücksichtigt werden. Dieses Problem ist bei uns fast ausschliesslich das Wirkungsgebiet des Instituts für Balneologie und Klimatologie. Die grösste Aufmerksamkeit schenkt man augenblicklich den Erfolgen der Klimatherapien bei allergischen Krankheiten des Kindesalters, aber auch bei Erwachsenen im See- und Gebirgsklima und letzters auch der klimatherapeutischen Behandlung in den Kammern des Salzbergwerkes in Wieliczka (hier unter wissenschaftlicher Aufsicht von Sabatowski und Obtułowicz).

B. DIE PHYSIKALISCHE RICHTUNG IN DER MEDIZINISCHEN BIOKLIMATOLOGIE.

Man befasst sich hier mit der Erforschung der einzelnen Elementen des Bioklimas: z.B. der Abkühlungsgrösze der Luft, der Sonnenstrahlung mit Berücksichtigung der U.V.-Strahlung, der Luftverunreinigung, der atmosphärischen Luftelektrizität, sowie dem Ozongehalt in der Grundschicht der Luft usw.

DIE ABKÜHLUNGSGRÖSZE

Das Problem der Abkühlungsgrösze der Luft ist Forschungsgegenstand mehrerer Anstalten. Besonders grosse Erfolge auf dem Gebiete der Abkühlungsforschung kann die biometeorologische Abteilung des Staatlichen Hydro-Meteorologischen Instituts aufweisen, die schon seit einigen Jahren systematisch katathermometrische Messungen an 31 Punkten der gesamten Landesfläche durchführt. In dieser Forschungsstelle hat man nicht nur für einzelne Orte (wie z.B. Ciechocinek oder Zakopane) das Material bezüglich der Luftabkühlungsgrösze bearbeitet, sondern auch für ganze Landesteile, z.B. für die südliche Ostseeküste, wobei man die Ergebnisse aus verschiedenen Orten des betreffenden Gebietes als Grundlage genommen hat.

DIE SONNENSTRAHLUNG

In der Erforschung des Solarklimas ist in Polen im Laufe der letzten Jahre ein grosser Fortschritt im Vergleich zur Vorkriegszeit zu verzeichnen. Das gegenwärtig bestehende aktinologische Netz umfasst 14 ständige Beobachtungsstellen, wovon 11 auf aktinographische und 3 auf aktinometrische Beobachtungsstellen entfallen. Das heliographische Netz dagegen stützt sich auf Beobachtungen, die in 60 Ortschaften registriert werden. Die Mehrzahl der obengenannten Beobachtungsstellen ist dem Staatlichen Hydro-Meteorologischen Institut untergeordnet.

Die Forschungsarbeiten über das Solarklima (Intensität der Sonnenstrahlung, Sonnenscheindauer) führen bei uns hauptsächlich: die biometeorologische Abteilung am Staatlichen Hydro-Meteorologischen Institut in Warszawa, dann die Beobachtungsstelle der Abteilung für Geophysik an der Polnischen Akademie der Wissenschaften in Belsk (bei Warszawa), die bioklimatische Forschungsstelle des Instituts für Zoologie an der Polnischen Akademie der Wissenschaften in Białyowieza,

das Observatorium des Instituts für Meteorologie und Klimatologie der Universität zu Wrocław, sowie das Institut für Balneologie und Klimatologie in Poznań, welches über Beobachtungsstellen in den Kurorten Inowrocław und Rymanów-Zdrój verfügt.

Ausserdem bestehen einige andere Forschungsstellen, wie z.B. die klimatologische Abteilung des Geographischen Instituts der Polnischen Akademie der Wissenschaften (Warszawa), dann die Lehrstühle für Meteorologie und Klimatologie der Universitäten in Toruń, Lublin und Wrocław, die sich mit periodischen aktinometrischen Messungen der direkten Sonnenstrahlung befassen zwecks Bestimmung des Energieverlustes, der durch Verunreinigung der Atmosphäre der Städte verursacht wird. Dasselbe Problem erforschen im oberschlesischen Industriegebiet die oben erwähnte klimatologische Abteilung des Geographischen Instituts der Polnischen Akademie der Wissenschaften gemeinsam mit dem Institut für Meteorologie und Klimatologie der Universität in Wrocław. Die Ergebnisse dieser Forschungen sind bereits in einem speziellen Bericht (Biulletin) veröffentlicht worden, der vom Komitee für die Angelegenheiten des oberschlesischen Industriegebietes an der Polnischen Akademie der Wissenschaften herausgegeben wurde. Diese Forschungsergebnisse wurden ausserdem im "International Journal of Bioclimatology and Biometeorology", vol I, 1957, part I, section D: "Rapporte des I. Internationalen Kongresses für Bioklimatologie und Biometeorologie - Wien, 23 - 27 IX. 1957", veröffentlicht.

Gegenwärtig führt die Mikroklimatische Abteilung der Sanitär-Epidemiologischen Station in Katowice in Oberschlesien profilartige Beobachtungen der Sonnenstrahlung durch.

Von anderen Arbeiten auf dem Gebiete der Sonnenstrahlung sind die studienartigen Arbeiten, die in der mikroklimatischen Forschungsstelle der Abteilung für Stadthygiene am Staatlichen Institut für Hygiene (Warszawa) und in der bioklimatischen Forschungsstelle des Instituts für Zoologie an der Polnischen Akademie der Wissenschaften in Białowieża geführt werden, zu nennen. Sie erforschen u.a. die Messmethoden der biologisch aktiven Ultraviolettstrahlung.

STAUBGEHALT DER LUFT

Das Verstaubungsproblem grosser Stadtzentren, vor allem gewisser Industriebezirke - wie des oberschlesischen Industriegebietes oder des Steinkohlenreviers von Wałbrzych (Niederschlesien) - war in den letzten Jahren Forschungsgegenstand unserer Bioklimatologen. In Oberschlesien führte das speziell für die Klimaangelegenheiten des oberschlesischen Industriegebietes berufene Komitee an der Polnischen Akademie der Wissenschaften in einer Zusammenarbeit mit dem Institut für Meteorologie und Klimatologie der Universität zu Wrocław sowie dem Staatsunternehmen "Energopomiar" (Gliwice) längere Messungsserien des Staubgehalts der Luft durch. Bei diesen Untersuchungen bediente man sich zweier Methoden:

1. der Sedimentationsmethode,
2. der konimetrischen Methode (Staubmesser von Ovens).

Die Forschungsergebnisse wurden in dem schon vorher erwähnten Bericht (Biulletin) des Komitees für die Angelegenheiten des oberschlesischen Industriegebietes an der Polnischen Akademie der Wissenschaften veröffentlicht.

Auch in anderen Städten wurde der Staubgehalt der Luft untersucht. So z.B. führte in Gdynia und Zakopane das Staatliche Hydro-Meteorologische Institut Messungen aus, in Kraków die Polnische Meteorologisch-Hydrologische Gesellschaft, in Warszawa die Sanitäts-Epidemiologische Station, in Wałbrzych das Physiographische Staatsunternehmen "Geoprojekt" und in Lublin, Łódź und Wrocław die sich an dortigen Universitäten befindlichen Institute für Klimatologie und Meteorologie.

Die in Wrocław durchgeführten Messungen führten zu folgenden Ergebnissen:

- a. Der Staubgehalt der Luft in dieser Stadt ist in kühlen Jahreszeiten (Heizperiode) fast vierfach grösser als in den warmen Jahreszeiten.
- b. Es wurde eine Korrelation zwischen dem Staubgehalt und einigen meteorologischen Elementen bestimmt.

Ausserdem erwiesen die gleichzeitig durchgeführten aktinometrischen Beobachtungen, dass das von der Stadt erzeugte Aerosol anders die Strahlung der Sonne und des Himmels beeinflusst, als das Aerosol der freien Atmosphäre.

DIE ATMOSPHERISCHE LUFTELEKTRIZITÄT

Die Probleme der atmosphärischen Luftelektrizität stellen in Polen gegenwärtig einen Aufgabenkreis zweier Forschungsstellen dar:

- a. des Geophysikalischen Observatoriums der Polnischen Akademie der Wissenschaften in Świdrig bei Warszawa,
- b. des Meteorologischen Instituts an der Agrarhochschule zu Poznań.

DIE RADIOAKTIVITÄT DER LUFT UND DER NIEDERSCHLÄGE

Diese Fragen werden in Polen an folgenden Forschungsstellen bearbeitet:

1. In der aerologischen Abteilung des Staatlichen Hydro-Meteorologischen Instituts in Legionowo bei Warszawa und in einigen Auszenkontrollmeszstellen, die der obengenannten Zentralstelle untergeordnet sind.
2. Am Lehrstuhl für allgemeine Physik an der Bergakademie in Kraków, der das Problem der Radioaktivität der Luft unter zwei Gesichtspunkten betrachtet und zwar:
 - (a) den Einfluss der klimatischen Verhältnisse auf die Radioaktivität der Luft,
 - (b) die radioaktiven Verunreinigungen als Konsequenz der nuklearen Explosionen.
3. In der chemischen Abteilung des Instituts für Balneologie und Klimatologie in Poznań. Man befasst sich hier mit dem Problem der Radioaktivität der Niederschläge.

OZON

An der Meszmethodik des bodennahen Ozons und der Konstruktion einer geeigneten Meszapparatur arbeiten:

1. die mikroklimatische Forschungsstelle der Abteilung für Stadthygiene am Staatlichen Institut für Hygiene (Warszawa),
2. die klimatische Abteilung des Geographischen Instituts der Polnischen Akademie der Wissenschaften (Warszawa), die schon einige Meszserien von Ozonkonzentration in der Grundschrift der Luft im Kurort Ciechocinek durchgeführt hat.

IV. DIE KLIMATOGRAPHIE DER KURORTE IN BIODYNAMISCHER HINSICHT

Gegenwärtig ist dieser Teil der Kurortklimatologie bei uns am meisten entwickelt. Man arbeitet jetzt an verschiedenen Forschungsstellen an einer Reihe von klimatologischen Monographien der Kur- und Erholungsorte bzw. Regionen, wozu man hauptsächlich das langjährige makroklimatische Material verwertet hat. So z.B. hat bereits die bioklimatische Abteilung des Instituts für Balneologie und Klimatologie eine umfangreiche Klimamonographie für den Talkessel und das Bergland von Kłodzko und eine weitere für das Schlesische Beskidenbergland fertiggestellt. In Bearbeitung befindet sich auch eine monographische Erfassung des Makroklimas der Westsudeten. Es sind ausserdem auch Klimacharakteristiken einzelner Kurorte z.B. von Ładek-Zdrój (Sudeten), Rymanów-Zdrój (Karpathenvorland), Inowrocław, Ciechocinek-Zdrój (mittel-polnische Tiefebene), Połczyn-Zdrój (Moränen-Hügellandschaft) und Kołobrzeg (Ostseeküste) erfasst worden.

Einige Klimacharakteristiken einzelner Kurorte stammen auch vom Arbeitskreis des Staatlichen Hydro-Meteorologischen Instituts. Dieser hat die Klimaverhältnisse von Zakopane - mit Berücksichtigung des polnischen Teiles der Hohen Tatra und des Vortatragebietes-, ferner die Klimacharakteristiken von Rabka Piwniczna (Karpathenbergland), Polanica (Sudeten) sowie der polnischen Ostseeküste bearbeitet.

In allen klimatographischen Charakteristiken schenkt man der Biodynamik einzelner Klimaelemente eine spezielle Aufmerksamkeit.

V. ARBEITEN AM LOKALKLIMA FÜR DEN BEDARF DER MEDIZIN UND DER STAEDTEBAUPLANUNG

DAS LOKALKLIMA DER KURORTE

Die Forschungsarbeiten am Lokalklima der Kurorte gehen in zwei Richtungen: 1. der wissenschaftlichen und 2. der praktischen:

1. In wissenschaftlicher Hinsicht bearbeiten das Lokalklima der Kurorte:
 - (a) das Institut für Balneologie und Klimatologie, das bereits Untersuchungen in Kołobrzeg, Miedzyzdroje, Krynica Morska und Rymanów-Zdrój durchgeführt hat, sowie
 - (b) die klimatische Abteilung des Geographischen Instituts der Polnischen Akademie der Wissenschaften, die eine Monographie des Lokalklimas vom Kurort Ciechocinek bearbeitet hat.
2. Für praktische Zwecke, z.B. für die Planung und Raumbewirtschaftung der Kurorte, werden klimatologische Gutachten vom speziell dazu berufenem Physiographischen Staatsunternehmen "Geoprojekt", welches dem Ministerium für Städte- und Siedlungsbau untersteht, bearbeitet.

DIE ARBEITEN AM LOKALKLIMA DER STAEDTE

Diese Arbeiten führt man für den Bedarf der kommunalen Wirtschaft, also der Planung und Raumbewirtschaftung. Sie werden angeregt und finanziert vom Forschungsinstitut für Städtebau und Architektur in Warszawa.

Die Studien am Klima der Städte und der Siedlungen werden von folgenden Forschungsstellen betrieben:

1. vom Lehrstuhl für Meteorologie und Klimatologie an der Universität in Łódź, der augenblicklich das Klima und Bioklima folgender Industriestädte bearbeitet: Łódź, Tomaszów Mazowiecki und Zgierz;
2. vom Lehrstuhl für Meteorologie und Klimatologie der Universität in Lublin, der das Lokalklima dieser Stadt bearbeitet;
3. vom Staatlichen Hydro-Meteorologischen Institut in Warszawa, welches das Stadtklima von Kraków bearbeitete und ausserdem sich an den Klimaforschungsarbeiten der Städte des oberschlesischen Industriegebietes beteiligte.
4. Das Lokalklima der Städte und Siedlungen gehört auch zu den Arbeitsproblemen des Lehrstuhls für Regionale Planung der Architektonischen Fakultät an der Technischen Hochschule zu Gdańsk. Aus diesem Institut stammt auch eine recht interessante Bearbeitung des Lokalklimas der Stadt Gdynia.
5. Gewisse Arbeiten an dem Lokalklima der Städte, jedoch ohne tiefer greifendewissenschaftliche Unterlage verfertigt auch das Physiographische Staatsunternehmen "Geoprojekt" in Warszawa.

An den Problemen der Lufthygiene in den Städten dagegen arbeiten speziell dazu berufene mikroklimatische Arbeitszellen an den städtischen und provinziellen Sanitäts-Epidemiologischen Stationen. Die wissenschaftliche Betreuung übernimmt das Staatliche Hygiene-Institut in Warszawa.

MISCELLANEOUS BIOCLIMATOLOGICAL DATA
Section C: Bioclimatological Institutions

UEBER DEN STAND DER AGRARMETEOROLOGISCHEN FORSCHUNG
IN DER DEUTSCHEN DEMOKRATISCHEN REPUBLIK

von

Prof. Dr. A.Mäde (D.D.R.)*

Mit diesem Bericht wird eine bibliographische Zusammenstellung vorgelegt, die agrarmeteorologische Beiträge aus den Jahren 1953 bis 1958 enthält. Bei der Zusammenstellung wurde Wert darauf gelegt, möglichst alle Veröffentlichungen zu erfassen, in denen über agrarmeteorologische Arbeitsergebnisse aus den wissenschaftlichen Institutionen in der Deutschen Demokratischen Republik berichtet. Bei einer Beschränkung auf die im Gebiet der DDR verlegten Druckereierzeugnisse wäre ein Ueberblick über den Umfang der Forschungsarbeiten auf dem Gebiete der Agrarmeteorologie nicht vollständig gewesen. Es muszten daher auch die Beiträge aufgenommen werden, die in anderen Publikationsorganen erschienen sind.

Stellt man der alphabetischen Ordnung nach dem Verfassernamen eine Gliederung nach den Institutionen gegenüber, in denen die Untersuchungen durchgeführt wurden, so erhält man einen Ueberblick über die Struktur der agrarmeteorologischen Forschung in der D.D.R.

Die Probleme der Agrarmeteorologie als einem Grenzgebiet zwischen der Biologie und der Meteorologie werden naturgemäß von Wissenschaftlern beider Disziplinen bearbeitet. Von Seiten der Meteorologie ist eine starke Förderung durch den Meteorologischen und Hydrologischen Dienst der D.D.R. zu verzeichnen. Ein Forschungsinstitut für Agrarmeteorologie dieses Dienstes arbeitet unter Leitung des Berichterstatters in Halle. Zu seinem Arbeitsbereich gehören die agrarmeteorologischen Forschungsstationen in Grosz-Lüsewitz, Müncheberg (Mark), Aschersleben, Dresden-Pillnitz, Marquardt und Potsdam-Bornim. Es gehörte eine zunächst weitere Station in Quedlinburg dazu, die im Jahre 1954 in das Institut für Pflanzenzüchtung der Deutschen Akademie der Landwirtschaftswissenschaften als Abteilung dieses Instituts übernommen wurde.

In diesem Netz agrarmeteorologischer Forschungsstützpunkte werden in enger Zusammenarbeit mit Instituten der Deutschen Akademie der Landwirtschaftswissenschaften Probleme bearbeitet, die von einer einzelnen Fachrichtung, der Meteorologie oder der Agrarwissenschaft, nur unvollkommen gelöst werden können.

Seit mehr als 10 Jahren wird an diesen Stellen die kollektive Arbeitsweise praktiziert. Veröffentlichungen aus diesem Bereich liegen von folgenden Mitarbeitern vor: Fusz, F., Hoffmann, K., Koitsch, R., Mäde, A., Raeuber, A., Schmidt, M., Schrödter, H.

Der Meteorologische und Hydrologische Dienst besitzt für den Sektor der humanen Bioklimatologie eine ähnliche Forschungseinrichtung. Ueberschneidungen zwischen Agrarmeteorologie und humaner Bioklimatologie sind nicht zu vermeiden. So liegen aus dem Bereich des Forschungsinstituts für Bioklimatologie in Berlin-Buch und seinen Forschungsstationen von Hentschel, G., Kühn, G., Mrose, H., Wagner, H., und Zenker, H. Arbeitsergebnisse vor, die mit in diese agrarmeteorologische Bibliographie aufgenommen werden können.

Aus anderen Institutionen des MHD sind Arbeiten vorgelegt worden, die für den Agrarmeteorologen ebenfalls von Interesse sind. Als Verfasser seien genannt: Gelbke, W., Hinzpeter, H., Matzke, H., Seyfert, F., Skeib, G. und Vilkner, H.

In einer Reihe von Instituten der Deutschen Akademie der Landwirtschaftswissenschaften wird ebenfalls zu agrarmeteorologischen Problemen Stellung genommen. Vor allem von den Instituten für Pflanzenzüchtung sind wertvolle Beiträge vorgelegt worden. Als Bearbeiter sind zu nennen: Athenstädt, H., Dahse, F., Engel, K.H., Junges, W., Krümmel, H., Löhrke, L., Müller, H.J., Pechert, H., Scholz, K., Sörgel, G., Unger, K., Vogel, G., Werner, K. Aus dem Bereich der von der Akademie betreuten Institute der ehemaligen Biologischen Zentralanstalt für Land- und Forstwirtschaft berichteten Hoffmann, G., Klemm, M., Kirchner, H.A., Müller, F.P. und Richter, G.

Im Bereich der Universitäten sind die meteorologischen und agrarmeteorologischen Institute durch Veröffentlichungen von Hesse, W., Koch, H.G., Pfeifer, F. und Stenz, S. vertreten. Von geographischer Seite haben sich Bauer, L., Reinhard, H. und Schulze, J.H. geäußert. In groszem

*) Agrarmeteorolog. Institut, Gr. Steinstrasse 81, Halle/Saale, Germany (D.D.R.).

Umfang legten die land- und forstwirtschaftlichen Institute der verschiedenen Universitäten Bearbeitungen agrarmeteorologischen Inhalts vor. An Mitarbeitern sind aus den einzelnen Universitäten zu nennen:

HUMBOLDT-UNIVERSITÄT BERLIN: Baumann, H., Casperson, G., Ehwald, E., Fröhlich, H., Göhre, K., Hennebo, D., Illner, K., Olbertz, M.H., Richter, P., Schulze, J., Schumann, A., Weber, H., Wilamowitz-Moellendorf und Zillmann, K.H.

MARTIN - LUTHER - UNIVERSITÄT HALLE-WITTENBERG: Bauckmann, H., Dörter, K., Friedrich, G., Meier, G.

FRIEDRICH - SCHILLER- UNIVERSITÄT JENA: Baltzer, H., Bergmann, W., Kretschmer, G., Matzner, F. und Rust, G.

KARL-MARX - UNIVERSITÄT LEIPZIG: Britz, L., Fröhlich, G., Geiler, H., und Quass, R.

UNIVERSITÄT ROSTOCK: Bauer, F. und Bochow, H.

PAEDAGOGISCHE HOCHSCHULE POTSDAM: Müller-Stoll, W.R. und Steubing, L.

Zu Fragen des Wasserhaushalts liegen Mitteilungen aus der TECHNISCHEN HOCHSCHULE DRESDEN (Busch, K.F.) und dem INSTITUT FÜR WASSERWIRTSCHAFT (Kalweit, H., und Täumer, F.) vor.

Ueberblickt man die Thematik der Publikationen, so ist festzustellen, dass den für die agrarmeteorologische Forschung grundlegenden methodischen Problemen grosse Beachtung geschenkt wurde. Dies gilt sowohl für den Sektor der Messtechnik als auch für den der Bearbeitung der Messwerte. In zunehmendem Masse werden dabei moderne statistische Prüfmethode angewandt.

Einen grossen Raum nehmen die Berichte über die Zusammenhänge zwischen Witterung und Pflanzenkrankheiten oder Pflanzenschädlingen ein. Das Ziel dieser Untersuchungen ist es, Unterlagen für den Aufbau eines Schädlingswarndienstes zu schaffen.

Dem Wasserhaushalt landwirtschaftlicher Kulturen und dem Einfluss der Witterung auf das Pflanzenwachstum und den Ertrag wird ebenfalls grosse Aufmerksamkeit gewidmet. Diese Untersuchungen streben in zunehmendem Umfang einen auch vom Standpunkt der Statistik her gesicherte Aussagewert der Ergebnisse an. Sie benutzen simultan phänometrische und meteorologische Beobachtungen, die korrelationsanalytisch bearbeitet werden.

BIBLIOGRAPHIE

1. ATHEENSTAEDT, H., SCHMADLAK, J. und ZAHN, F.P.: Ueber die Frostschäden an Obstgehölzunterlagen im Winter 1953/54 - Arch.f. Gartenbau, 4, 6-67, 1956.
2. BALTZER, H. : Untersuchungen über die mikroklimatischen Wirkungen der winterlichen Weideabdeckung - Wiss.Z.Univ.Jena, Math.-Nat. Reihe, 1, 49-60, 1952/53.
3. BAUCKMANN, M. : Untersuchungen über eine termingerechte Bekämpfung des Apfelwicklers unter Berücksichtigung des Falterfluges - Arch.f. Gartenbau, 4, 253-277, 1956.
4. BAUER, F. : Agrarmeteorologische Untersuchungen über die nächtliche Wiesenunterkühlung innerhalb norddeutscher Feldgemarkungen. Ein Beitrag zur Gäländeklimatologie des Flachlandes - Diss. Rostock 1956.
5. BAUER, L. : Hochwasserabfluss und Landschaftshaushalt - Herm. Haack, Geograph.-Kartograph.Anst. Gotha, 1956.
6. BAUMANN, H. und OLBERTZ, M.H. : Untersuchungen über die Entwicklungstendenzen und Sorptionsfähigkeiten langjährig genutzter Rieselböden im Berliner Abwasserverwertungsgebiet - Wiss.Z.Humboldt-Univ.Berlin, Math.-Nat. Reihe, 2, 43-61, 1952/53.
7. BAUMANN, H. : Ueber den Wasserhaushalt des kulturpflanzentragenden Standorttes - Sitz.-Ber.Dt.Akad.Landwirtschaftswiss. Berlin, 3, 1, 28, 1954.
8. BAUMANN, H. und KREIL, W. : Die Weideberechnung - D.Dt.Landwirtsch., 5, 496-500, 1954.
9. BAUMANN, H. : Zur Frage des Kartoffelabbaues in der Uebergangslage - D. Dt. Landwirtsch., 5, 570-578, 1954.
10. BAUMANN, H. und KLAUS, M.L. : Gefässversuche mit Sommergetreide zur Frage der konstitutionellen Anpassung der Pflanzen an die Wasserversorgung - Z. f. Acker- u. Pflanzenbau, 101, 35-52, 1956.
11. BAUMANN, H. und KLATT, F. : Berechnungszeitpunktversuche mit Hafer- und Winterweizensorten - Z.f. Acker- u. Pflanzenbau, 103, 380-396, 1957.
12. BERGMANN, W. : Ueber den Einfluss von Umweltfaktoren auf Wurzelwachstum und Ernteertrag - D.Dt. Landwirtsch., 5, 259-264, 1954.
13. BERGMANN, W. und RIETZEL, P. : Ein Beitrag zur Standortauswahl von Hopfengärten - D.Dt. Landwirtsch., 5, 511-514, 1954.

14. BERNBECK, O. : Wind und physiologische Tiefgründigkeit in ihrer Bedeutung für die Bodenkultur - Deutscher Bauernverlag, 100, 1954.
15. BOCHOW, H. : Der Einfluss der Witterung auf das Auftreten der Kraut- u. Knollenfäule (Phytophthora infestans Mont. de Bary) und die sich daraus ergebenden Möglichkeiten für die Einrichtung eines Warndienstes zur Durchführung prophylaktischer Massnahmen - Wiss. Z. Univ. Rostock, Math.-Nat. Reihe, 4, 47-66, 1954/55.
16. BOCHOW, H. und RAEUBER, A. : Untersuchungen über den Einfluss niederer Temperaturen auf die Krautfäuleerreger Phytophthora infestans de By - Nachr.-Bl. f. Dt. Pflanzenschutzdienst, 6, 120-123, 1956.
17. BRITZ, L. und EICHLER, Wd. : Stubenfliegenbekämpfung und Witterungsfaktoren - Wiss. Z. Univ. Leipzig, Naturwiss. Reihe, 2, 255-258, 1952/53.
18. BRITZ, L. und HOEHNE, W. : Temperaturschwankung und Entwicklungsgeschwindigkeit bei Anopheles atroparvus (Diptera; Culcidae) - Z. angew. Zool., 2, 209-234, 1955.
19. BUSCH, K.-F. : Das Wasserhaltevermögen des Bodens - Wasserwirtsch.-Wasser-techn., 6, 116-121, 1956.
20. CASPERSON, G. : Untersuchungen über den Einfluss von Windschutzanlagen auf den standörtlichen Wärmehaushalt - Angew. Met., 2, 339-351, 1957.
21. CASPERSON, G. : Erfassung des Wärmehaushalts in Windschutzgebieten - D. Dt. Landwirtschafts., 8, 537-538, 1957.
22. DAHSE, F. : Der Feuchtigkeitsverlauf im Getreide bei verschiedenen Ernteverfahren - D. Dt. Landwirtschafts., 7, 428-432, 1956.
23. DOERTER, K. : Die Steighöhe des Wassers in humusfreien Sanden in Abhängigkeit von der Hygroskopizität - Diss. Halle, Landw. Fak., 1954.
24. DOERTER, K. : Hinweise über die seitliche Wasserausbreitung und den seitlichen Strangabstand bei unterirdischer Bewässerung versch. Mineralböden - Wasser u. Boden, 8, 124-130, 1956.
25. EHWALD, E. : Aufgaben und Methoden der forstlichen Standortskunde - Sitz.-Ber. Dt. Akad. Landwirtschaftswiss. Berlin, 2, 4, 20, 1953.
26. EHWALD, E. : Vergleichende Forstklimatologie des Thür. Gebirges und benachbarter Mittelgebirge - Wiss. Abh., 11, Deutscher Bauernverlag 1956.
27. ENGEL, K.H., HINZPETER, H., RAEUBER, A. : Die Abhängigkeit der Fleischfarbenbonitierung an Kartoffeln von der Beleuchtung - Z. f. Pflanzenzüchtung, 37, 77-80, 1957.
28. FIEDLER, H. : Ein Beitrag zur elektrischen Taumessung - Z. f. Met., 10, 217-219, 1956.
29. FRIEDRICH, G. und SCHWOPE, D. : Zusatzbewässerung und Frostschtzberechnung im Obstbau - D. Dt. Gartenbau, 5, 104-107, 1958.
30. FROELICH, G. : Zur Biologie und Bekämpfung der Kohlschoten-Gallmücke (Dasynura brassicae Winn) - Nachr.-Bl. f. d. Dt. Pflanzenschutzdienst, 6, 123-128, 1956.
31. FROELICH, H. : Zu Fragen der Bodenheizung bei der Treibtomatenkultur - Die Gartenbauwissenschaft, 4, 491-506, 1957.
32. FUSZ, F., ERNST, A., LAUENSTEIN, A., JUNGES, W., BORCH, B., MUELLER, H., THIELE, K., KAMMEYER, H.F. : Die Frostnacht vom 19. zum 20. Mai 1952 und ihre Auswirkungen auf gärtnerische Kulturpflanzen in Sachsen. - Arch. f. Gartenbau, 2, 9-68, 1954.
33. FUSZ, F. : Untersuchungen zur Frage des Windschutzes in gärtnerischen Kulturen - Angew. Met., 2, 86-95 und 97-110, 1954/55.
34. FUSZ, F. : Zur Frage des Wachstumsnullpunktes der Temperatur bei Winterweizen unter Freilandbedingungen - Wiss. Z. Univ. Halle, Math.-Nat. Reihe, 5, 513-521, 1955.
35. FUSZ, F. : Lufttemperatur und Luftfeuchtigkeit in drei verschiedenen Schweineställen im Sommer - Wiss. Z. Univ. Halle, Math.-Nat. Reihe, 5, 523-531, 1955.
36. FUSZ, F. : Spätfrost und Spätfrostvorhersage - Arch. f. Gartenbau, 3, 65-73, 1955.
37. FUSZ, F. : Untersuchungen zur Frage der Messung von Knospendurchmessern - Angew. Met., 2, 151-155, 1955.
38. FUSZ, F. : Zum Einfluss des Windschutzes auf den Ernteertrag der Erdbeeren - Angew. Met., 2, 176-183, 1955.
39. FUSZ, F. : Ueber die Entwicklung von Weizen in Abhängigkeit von der Bodentemperatur - Angew. Met., 2, 253-256, 1956.
40. FUSZ, F. : Thermische Wirkungen der Bodenbedeckung auf verschiedenen Bodenarten - Dt. Gartenbau, 4, 162-163, 1957.
41. FUSZ, F. : Veränderungen des Bodenklimas durch Bodenbedeckung bei ausdauernden Kulturen - Angew. Met., 3, 74-76, 1958.

42. GEILER, H. : Ueber die Möglichkeit der Einrichtung eines Schädlingsswarndienstes in der D.D.R. - D.Dt.Landwirtsch., 4, 194-196, 1953.
43. GEILER, H. und SCHOTT, H. : Erfahrungen beim Aufbau eines Schädlingsbeobachtungs- und Schädlingsswarndienstes für die Landwirtschaft in Mitteldeutschland - Nachr.-Bl.f.d.Dt.Pflanzenschutzdienst, 9, 42-45, 1955.
44. GELBKE, W. : Untersuchungen zur Methodik der Taumessung an Hand dreijähriger Taumeszreihen in Greifswald - Abh.Met.u.Hydrolog.Dienst d. DDR, 5, 38, 104, 1955.
45. GELBKE, W. : Ergebnisse dreijähriger Taumessungen - Ann.d.Met., 7, 79-86, 1955/56.
46. GOEHRE, K. : Lysimeterversuche über den Wasserhaushalt von unbewachsenem und mit jungen Eichen bewachsenem Boden - Arch.f.Forstwesen, 3, 645-674, 1954.
47. GOEHRE, K. und LUETZKE, R. : Der Einfluss von Bestandsdichte und -struktur auf das Kleinklima im Walde - Arch.f.Forstwesen, 5, 487-572, 1956.
48. HENNEBO, D. und ILLNER, K. : Strömungsbedingte Erscheinungen an Windschutzpflanzungen und ihre landwirtschaftliche Bedeutung - D.Dt.Landwirtsch., 4, 603-607, 1953.
49. HENTSCHEL, G. : Phänologisch-singuläre Prognostik der Winterroggenblüte - Angew.Met., 1, 314-318, 1953.
50. HESSE, W. : Karte der Vegetationsperiode - Angew.Met., 1, 276-278, 1953.
51. HESSE, W. : Der Einfluss meteorologischer Faktoren auf die Transpiration der Pfefferminze (*Mentha piperita* L.) - Angew.Met., 2, 14-18, 1954.
52. HESSE, W. : Ergebnisse von Pflanzentranspirationsmessungen mit Kleinlysimetern im Zusammenhang mit meteorologischen Einflüssen - Angew.Met., 2, 65-82, 1954.
53. HESSE, W. : Die physikalisch-meteorologischen Bedingungen im Luv und Lee von Waldstreifen und feldschützenden Hecken - Wiss.Z.Univ.Leipzig, Math.Nat. Reihe, 4, 153-161, 1954.
54. HESSE, W. : Ergebnisse von Pflanzentranspirationsmessungen - Ann.d.Met., 7, 111-118, 1955/56.
55. HOEHNE, W. : Ein thermoelektrischer Feinwindmesser - Z.f.Met., 7, 215-218, 1953.
56. HOEHNE, W. : Häufigkeitsstatistische Untersuchungen an einer 50-jährigen Reihe von Minimaltemperaturen. Ein Beitrag zum Frostschadenproblem - Wiss.Z.Univ.Halle, Math.-Nat. Reihe, 4, 191-198, 1954.
57. HOEHNE, W. : Abschätzung der Frostgefährdung witterungsabhängiger Phasen der Pflanzenentwicklung - Wiss.Z.Univ.Halle, Math.-Nat. Reihe, 4, 199-201, 1954.
58. HOEHNE, W. : Experimentelle und mikroklimatische Untersuchungen an Kleingewässern - Abh.Met.u.Hydrolog.Dienst der DDR, 4, 26, 56, 1954.
59. HOEHNE, W. : Ueber die Weiterentwicklung des thermoelektrischen Feinwindmessers - Z.f.Met., 8, 243-247, 1954.
60. HOEHNE, W. : Windrichtungsschreiber für schwache Luftströmungen - Z.f.Met., 9, 135-143, 1955.
61. HOEHNE, W. : Zur Auswertmethodik von Punktregistrierungen bei mikroklimatologischen Messungen - Angew.Met., 2, 212-217, 1956.
62. HOEHNE, W., MAEDE, A. und SCHMIDT, M. : Kompensation des Einflusses der Vergleichstemperatur bei thermoelektrischen Messungen - Z.f.Met., 10, 131-136, 1956.
63. HOEHNE, W. : Theoretische Betrachtungen über die Verwendbarkeit von Halbleiter-Widerständen in der Mikrometeorologie und -klimatologie - Z.f.Met., 11, 143-156, 1957.
64. HOFFMANN, G. und SCHROEDTER, H. : Mikroklimatische Untersuchungen bei Kartoffelschorfinfektionen in Lochtöpfen unter Freilandbedingungen - Der Züchter, 24, 131-137, 1954.
65. HOFFMANN, K. : Untersuchungen zur Frage des Kornfeuchteverlaufs bei Winterweizen - Angew.Met., 3, 44-46, 1957.
66. ILLNER, K. : Ueber den Einfluss von Windschutzpflanzungen auf die Unkrautverbreitung - Angew.Met., 2, 370-373, 1957.
67. JUNGES, W. : Zur Unterscheidung von entwicklungsphysiologisch optimaler und ertragsmäßig optimaler Jarowisationsdauer bei der Buschtomate (*Lycopersicon esculentum* Mill. var. *finiens* Lehm) - Arch.f.Gartenbau, 2, 213-221, 1954.
68. JUNGES, W. : Die jährliche Niederschlagsverteilung als entscheidender Faktor bei der photoperiodischen Anpassung der Pflanze - Die Gartenbauwissenschaft, 4, 527-540, 1957.

69. JUNGES, W. : Beziehungen zwischen der photoperiodischen Angepasstheit der Pflanzen und der geogr. Breite ihrer Heimat - Planta, 49, 11-32, 1957.
70. KALWEIT, H. : Lysimeterbeobachtungen als Grundlage zur Berechnung des Wasserhaushaltes - Z.f. Met., 7, 35-42, 1953.
71. KALWEIT, H. : Beitrag zum Gebietsverhalten der Verdunstung in mitteldeutschen Landschaften - Z.f. Met., 7, 373-380, 1953.
72. KALWEIT, H. : Der Wasserhaushalt, Berechnungsverfahren und Anwendungen auf mitteldeutsche Flussgebiete - Verl. Technik, 1, 408 und 2, 279, 1953.
73. KALWEIT, H. : Der Einfluss der Steigerung des land- und forstwirtschaftlichen Hektarertrages auf den Wasserhaushalt - Wasser und Boden, 7, 66-69, 1955.
74. KIRCHNER, H.A. : Der Einfluss der Spätfröste auf den Raps - Nachr.-Bl.f.d. Dt. Pflanzenschutzdienst, 7, 175-177, 1953.
75. KLEIN, W. : Untersuchungen zur Bestimmung des Keimungsminimums landwirtschaftlicher Unkräuter - Z.f. Acker- und Pflanzenbau, 101, 395-430, 1956.
76. KLEMM, M. : Bio-phänologischer Kalender. Entwicklungsdauer einiger wichtiger Schädlinge im Acker-, Obst- und Gemüsebau - Nachr.-Bl.f.d. Dt. Pflanzenschutzdienst, 7, 130-132, 1953.
77. KLEMM, M. und MASURAT, G. : Schadgebiete der Phytophthora infestans in der DDR - Nachr.-Bl.f.d. Dt. Pflanzenschutzdienst, 7, 221-224, 1953.
78. Klimaatlas für das Gebiet der DDR, hrg.v. Met.u.Hydrol.Dienst der DDR - Akademieverlag 1953.
79. KOCH, F. : Die Auswirkungen der anormalen Trockenheit des Sommers 1952 auf die Ackerunkrautgemeinschaften deutscher Dauerdüngungsversuche - Z.f. Pflanzenbau u. Pflanzenschutz, 6, 32-40, 1955.
80. KOCH, H.G. : Durchmesserzuwacherhebungen an Waldbäumen zum Nachweis vom Geländeklimaeinflüssen - Arch.f. Forstwesen, 5, 673-689, 1956.
81. KOCH, H.G. : Zur Phänologie der Bienevölker - Angew.Met., 3, 33-43, 1957.
82. KOCH, H.G. : Jahresringauszählungen an Waldbäumen zum Nachweis witterungsbedingter Zuwachsschwankungen - Ann.d.Met., 8, 21-33, 1957.
83. KOITZSCH, R., RABSCH, B. : Ueber Erdbohrer zur Entnahme kleiner Volumenproben bei leichten Böden - Ann.d.Met., 2, 240-244, 1956.
84. KRUEHNE, G. : Karte der Maifrostschäden 1953 in Wernigerode/Harz - Z.f. Met., 8, 180-182, 1954.
85. KRUEMMEL, H. : Standortkartierung im Obstbau - Wiss.Z.Univ.Leipzig, Math.-Nat. Reihe, 5, 309-319, 1955/56.
86. KRETSCHMER, G. : Pflanzenwachstum und Temperatur - Wiss.Z.Univ.Jena, Math.-Nat. Reihe, 4, 634-638, 1954/55.
87. KRETSCHMER, G. : Messungen der vertikalen Volumänderung von Ackerböden - Wiss.Z.Univ.Jena, Math.-Nat. Reihe, 4, 640-645, 1954/55.
88. KRETSCHMER, G. : Das Verhalten gefrierender Bodenwassers im Acker - Angew.Met., 2, 193-211, 1956.
89. KRETSCHMER, G. : Die vom Frost verursachte Wasserbewegung im Boden und ihre Wirkung auf Pflanzen - D.Dt.Landwirtsch., 7, 199-200, 1956.
90. LOEHRKE, L. : Bodenfeuchtigkeit in Obstanlagen mit und ohne Gründungsanbau - Arch.f. Gartenbau, 4, 93-111, 1956.
91. MAEDE, A. : Ueber die Höhe des nächtlichen Taufalles im mitteldeutschen Trockengebiet - Wiss.Z.Univ.Halle, Math.-Nat.Reihe, 4, 185-189, 1954.
92. MAEDE, A. : Zur Methodik der Taumessung - Wiss.Z.d.Univ.Halle, Math.-Nat. Reihe, 5, 483-512, 1955.
93. MAEDE, A. : Ueber die Methodik der meteorologischen Gelände Vermessung - Sitz.-Ber.Dt.Akad.Landwirtschaftswiss. Berlin, 5, 25, 1956.
94. MATZNER, F. : Die Bedeutung einer Hagelschadenkarte für die Landwirtschaft und den Gartenbau - Wiss.Z.Univ.Jena, Math.-Nat.Reihe, 3, 299, 1954.
95. MATZNER, F. : Die Hagelgefährdung in den nördlichen Bezirken der Deutschen Demokratischen Republik - D.Dt.Landwirtsch., 9, 24-27, 1958.
96. MATZNER, F. : Die Hagelgefährdung der mittleren Bezirke der Deutschen Demokratischen Republik - D.Dt.Landwirtsch., 9, 125-129, 1958.
97. MATZNER, F. : Die Hagelgefährdung in den südlichen Bezirken der Deutschen Demokratischen Republik - D.Dt.Landwirtsch., 9, 339-343, 1958.
98. MATZKE, H. und GESELLE, P. : Ueber ein Gerät zur Registrierung der nächtlichen "effektiven Ausstrahlung" - Ann.d.Met., 7, 73-78, 1955/56.
99. MEIER, G. : Der Einfluss von Blütenfrosts Schäden auf den Ertrag verschiedener Obstarten, Sorten und Baumformen auf dem Obstversuchsfeld

- Prussendorf - Dt. Gartenbau, 5, 129-132, 1958.
100. MROSE, H. : Der Einfluss des Waldes auf die Luftfeuchtigkeit - Angew.Met., 2, 281-286, 1956.
 101. MUELLER, F.P. : Prognose des Massenauftritts von Blattläusen bei Berücksichtigung des Wirtswechsels - Nachr.-Bl.f.d.Dt. Pflanzenschutzdienst, 8, 206-209, 1954.
 102. MUELLER, H.J. und UNGER, K. : Ueber die Bedeutung der Zusammenhänge zwischen Witterung und Blattlausflug für die Probleme des Kartoffelabbaus - Forsch.u. Fortschr., 29, 229-238, 1955.
 103. MUELLER, H.J. : Die Saisonformenbildung von *Arachnia levana*, ein photoperiodisch gesteuerter Diapause-Effekt - D. Naturwiss., 42, 134-135, 1955.
 104. MUELLER, K. : Untersuchungen über die Abhängigkeit der Winterroggenreife von der Witterung - Veröff.d. Instit.f.Agrarmet.d.Univ.Leipzig, 1, 122-173, 1957.
 105. MUELLER-STOLL, W.R. und FREITAG, H. : Beiträge zur bestandsklimatischen Analyse von Wiesen- gesellschaften (Untersuchungen im Ober-Spree-Wald) - Angew.Met., 3, 16-30, 1957.
 106. OLBERTZ, M.H. : Ueber die am Standort des Kulturbodens erfassbaren Größen des Wasserhaushaltes - Wiss.Abh.d.Dt.Akad.d. Landwirtschaftswiss., 23, 109, 1957.
 107. PECHERT, H. : Diskussionsbeitrag zu: "Künstliches Licht in landwirtschaftlichen Bauten" von Dipl.-Ing.F.K. Mann in Heft 1/1956 - D.Dt. Landwirtsch., 7, 251-252, 1956.
 108. PFEIFER, F. : Eine Schwimmwaage zur Registrierung der Transpiration - Z.f. Met., 10, 22-28, 1956.
 109. PFEIFER, F. : Untersuchungen über die Transpirationsintensität von Erbsen (*Pisum sativum*) mittels einer registrierenden Schwimmwaage - Angew.Met., 2, 374-381, 1957.
 110. QUAAS, R. : Die unterirdische Zufuhr von Wasser und ihre Wirkung auf die Transpiration der Pflanzen - D.Dt.Landwirtsch., 4, 155-156, 1953.
 111. QUAAS, R. : Die Transpirationsintensität als Grundlage für die Ermittlung des Wasserbedarfs bei Wintergerste - D.Dt.Landwirtsch., 4, 299-300, 1953.
 112. RAEUBER, A. : Meteorologische Vergleichsmessungen zwischen Schuppenstall und Freiland in Grosz-Lüsewitz - Angew.Met., 2, 217-222, 1956.
 113. RAEUBER, A. : Untersuchung zur Witterungsabhängigkeit der Krautfäule der Kartoffel im Hinblick auf einen Phytophthora-Warndienst - Abh.Met. u. Hydrolog. Dienst d. DDR, 6, 40, 38, 1957.
 114. RAEUBER, A. : Kritische Tage und Krautfäuleaufreten an Kartoffeln in England und Mecklenburg 1950-1955 - Angew.Met., 3, 47-51, 1957.
 115. RAEUBER, A. und BOCHOW, H. : Ueber die Bedeutung des Sporenfluges von *Phytophthora infestans* für den Warndienst - Nachr.-Bl.f.d.Dt. Pflanzenschutzdienst, 11, 6-11, 1957.
 116. REINHARD, H. : Die Reifezeit des Winterroggens in Mecklenburg - Wiss.Z. Univ. Greifswald, Sonderheft 2, 1, 47-54, 1952/53.
 117. RICHTER, G. : Die Maikäferpopulation im Gebiete der Deutschen Demokratischen Republik - Nachr.-Bl.f.d.Dt. Pflanzenschutzdienst, 12, 21-35, 1958.
 118. RICHTER, P. : Das Wittenberger Maiblumenanbaugbiet und seine klimatischen und edaphischen Bedingungen als Grundlagen seiner wirtschaftlichen Entwicklung - Arch.f.Gartenbau, 5, 179-251, 1957.
 119. RUST, G. : Abhängigkeit der Erträge der Sojabohne vom Ablauf der Witterungsverhältnisse - Wiss.Z.Univ.Jena, Math.-Nat.Reihe, 4, 451-458, 1954/55.
 120. SCHMIDT, M. : Ueber die Berechnung von Tagesmitteltemperaturen der oberen Bodenschichten - Z.f.Met., 10, 52-56, 1956.
 121. SCHMIDT, M. : Der Einfluss der Temperatur auf Schlüpfzeitpunkt und Flugaktivität des Apfelwicklers (*Carpocapsa pomonella* L.) - Arch. f. Gartenbau, 6, 12-22, 1958.
 122. SCHMIDT, M. : Der Einfluss der Temperatur auf Beginn und Verlauf des Apfelwicklerfluges - Arch.f.Gartenbau, 4, 277-279, 1956.
 123. SCHOLZ, K. und PECHERT, H. : Staubentwicklung im Stall, ihre Feststellung und Masnahmen zu ihrer Minderung - D.Dt.Landwirtsch. 4, 326-331, 1953.
 124. SCHOLZ, K. und PECHERT, H. : Die Untersuchungen von Stallluft hinsichtlich ihres Gehaltes an Kohlendioxyd mit Hilfe des Interferometers von Zeiss - D. Dt. Landwirtsch., 7, 246-249, 1956.

125. SCHULTZE, J.H. : Die naturbedingten Landschaften und ihre Rolle in der Land- und Forstwirtschaft - D.Dt.Landwirtsch., 7, 26-30, 1956.
126. SCHULZE, J. : Die Kältetoleranz des Maises - D.Dt.Landwirtsch., 9, 111-113, 1958.
127. SCHROEDTER, H. und SCHEIDING, U.: Die Abhängigkeit der Aktivität des Kohlgallenrüsslers (*Ceuthorrhynchus pleurostigma* Marsh.) von klimatischen Faktoren - Nachr.-Bl.f.d. Dt. Pflanzenschutzdienst, 7, 143-148, 1953.
128. SCHROEDTER, H. : Agrarmeteorologische Untersuchungen im Rahmen der Antibiotica-Forschung in der Pflanzenpathologie - Angew.Met., 2, 23 - 26, 1954.
129. SCHROEDTER, H. : Weitere Untersuchungen über die Anwendung der Temperatursummenregel auf phänologische Phasen - Angew. Met., 2, 59-64, 1954.
130. SCHROEDTER, H. : Die Bedeutung von Massenaustausch und Wind für die Verbreitung von Pflanzenkrankheiten - Nachr.-Bl.f.d.Dt. Pflanzenschutzdienst, 8, 166-172, 1954.
131. SCHROEDTER, H. : Zur meteorologischen Problematik eines Warndienstes im Pflanzenschutz - D.Dt. Landwirtsch., 6, 236-239, 1955.
132. SCHROEDTER, H. : Meteorologische Probleme der Antibiotikaforschung - Ann. d. Met., 7, 8-11, 1955/56.
133. SCHROEDTER, H. : Die Antibiotikawirkung im Pflanzenschutz ist wetterabhängig - Umschau, 56, 114-116, 1956.
134. SCHROEDTER, H. : Phänometrisch.-statistische Untersuchungen zum Problem "Witterung und Pflanzenwachstum" - Ann.d.Met., 8, 1-6, 1957.
135. SEYFERT, F. : Die Bedeutung phänologischer Beobachtungen für die Imkerei - Angew.Met., 2, 155-157, 1955.
136. SEYFERT, F. : Eintritt phänologischer Phasen und auslösende Wetterlagen - Angew.Met., 2, 183-190, 1955.
137. SEYFERT, F. : Anleitung zur Durchführung phänologischer Beobachtungen. 2. erw.Aufl.- Akademie-Verlag Berlin, 139, 1953.
138. SEYFERT, F. : Die klimatischen Voraussetzungen des Zwischenfruchtanbaues nach Winterroggen und Wintertraps in der Deutschen Demokratischen Republik unter Berücksichtigung von biologischen und Bodenfaktoren - Abh.Met.u.Hydrolog.Dienst der DDR, 4, 25, 35, 1954.
139. SEYFERT, F. : Phänologische Tabellen 1947-1950 - Abh.Met.u.Hydrolog. Dienst der DDR, 5, 37, 366, 1957.
140. SKEIB, G. : Ein rotierender Strahlungssumsatzmesser mit induktiver Uebertragung der Meszspannung - Z.f.Met., 7, 167-171, 1953.
141. SKEIB, G. : Die Weiterentwicklung des rotierenden Strahlungssumsatzmessers mit induktiver Uebertragung der Meszspannung - Z.f.Met., 10, 33-38, 1956.
142. SKEIB, G. : Ueber das Verhalten von Strahlungssumsatzmessern im instationären Zustande - Z.f.Met., 10, 97-100, 1956.
143. SOERGEL, G., UNGER, K. : Ueber den Befallsverlauf von *Myeosphacarella pinodes* *Ascochyta pinodella* und *Ascochyta pisi* an Erbsen am natürlichen Standort - Der Züchter, 24, 56-68, 1954.
144. SOERGEL, G. : Zur Wirkung der Temperatur auf die Größe von Pilzsporen - D. Naturwiss., 42, 565-566, 1955.
145. STENZ, S. : Ueber den Einfluss von Witterung und meteorologischen Größen auf die Ertragsleistung von Obstgehölzen - Veröff.d.Inst. f. Agrarmet.d.Univ. Leipzig, 1, 77-114, 1956.
146. STEUBING, L. : Experimentelle Untersuchungen über die Veränderung der Standortfaktoren durch Windschutzanlagen - Oikos.Acta Oecol.Scand., 4, 118-147, 1952/53.
147. STEUBING, L. : Der Tau und seine Bedeutung für die Pflanze - Umschau, 53, 456-457, 1953.
148. STEUBING, L. : Studien über den Taufall als Vegetationsfaktor - Ber.d.Dt.Botan.Ges., 68, 55-70, 1955.
149. STEUBING, L. und MUELLER-STOLL, W.R.: Ueber die Beeinflussung der Standortfaktoren und der Ertragsbildung durch Windschutz-Streifen aus hochwüchsigen Nutzpflanzen - Angew. Bot., 29, 90-107, 1955.
150. STEUBING, L. : Beobachtungen an Lysimetern verschiedener Bauart - Ber.d. Dt. Botan.Ges., 69, 35-48, 1956.
151. TAEUMER, F. : Eine Methode zur Bestimmung der Gebietsverdunstung auf meteorologischer Grundlage - Veröff.d.Inst.f.Agrarmet.d.Univ.Leipzig, 1, 1-71, 1955.

152. UNGER, K. und FABIG, F. : Ueber den Einfluss der Temperatur auf die Erntezeit und Ertragssicherheit bei einigen Tomatensorten. Quedlinburger Beitr. z. Züchtungsforsch. Nr. 13 - Der Züchter, 23, 257-266, 1953.
153. UNGER, K. und MUELLER, H.J. : Studien über die Bedeutung von Witterung und Mikroklima für den Massenwechsel der Schwarzen Bohnenlaus (*Doralis fabae* Scop.) - Angew.Met., 1, 257-275, 1953.
154. UNGER, K. : Bodenfeuchtigkeit und Alsterwelke - Angew.Met., 1, 278-280, 1953
155. UNGER, K. : Thermoelementpsychrometer für mikroklimatische Messungen - Angew.Met. 1, 280-283, 1953.
156. UNGER, K. : Agrarmeteorologische Studien - Abh.Met.u.Hydrolog.Dienst der DDR, 3, 19, 27, 1953.
157. UNGER, K. : Zur Abschätzung von Verdunstungsunterschieden verschiedener Pflanzenbestände am natürlichen Standort - Angew.Met., 2, 1-14, 1954.
158. UNGER, K. : Zur klimatologischen Analyse der Kartoffelabbau- und Gesundheitslagen - Angew.Met. 2, 26-32, 1954.
159. UNGER, K. und MUELLER, H.J. : Ueber die Wirkung geländeklimatisch unterschiedlicher Standorte auf den Blattlausbeflug - Der Züchter, 24, 337-345, 1954.
160. UNGER, K. und SCHNEIDER, A. : Ueber Zusammenhänge zwischen der Reifeentwicklung von Gemüserbsen und bestimmten mikrometeorologischen Faktoren - Der Züchter, 26, 12-22, 1956.
161. UNGER, K. : Zum Einfluss der klimatischen Standortfaktoren auf die Blühzeit und den Blühverlauf einiger Gemüserbsensorten - Z.f. Acker- u.Pflanzenbau, 102, 69-80, 1957.
162. UNGER, K. : Zur Anwendung der Durchstrahlungsmethode für phänometrische Messungen mit Hilfe von radioaktiven Strahlungsquellen - Angew.Met., 3, 115-118, 1958.
163. UNGER, K. : Eine Thermobatterie mit kompensierter Vergleichstemperatur für mikrometeorologische Temperaturmessungen - Arch.f.Met., Geoph. u. Biokl., Serie B, 8, 378-381, 1958.
164. VILKNER, H. : Ein Geländeverdunstungsmessgerät - Z.f.Met., 8, 247-251, 1954.
165. VILKNER, H. : Eine Verdunstungs-Fernregistriereinrichtung mit einer neuen Form eines Abtropfrohrs - Z.f.Met., 9, 271-277, 1955.
166. VILKNER, H. : Ueber die Erdbodenfeuchtigkeit und ihre Messung mittels elektrischer Kapazität - Z.f.Met., 11, 49-58, 1957.
167. VOGEL, G. : Die Beeinflussung des Wärmehaushaltes des Bodens und der bodennahen Luftschicht durch Anwendung der Bodenheizung in Freiland - Dt. Gartenbau, 5, 5-7, 1958.
168. WAGNER, H. und DINGER, H.J. : Die Besonnung im oberen Vogtland und ihre Bedeutung für das Pflanzenwachstum - Angew.Met., 2, 122-125, 1955.
169. WAGNER, H. und HUNGER, W. : Ueber Besonnungsverhältnisse im Raume des Elsterer Kessels in ihrer Bedeutung für forstliche Standortfragen - Angew.Met., 2, 365-369, 1957.
170. WEBER, H. : Ueber Kondensationsvorgänge im gewachsenen Boden - Wiss.Z.Humboldt-Univ. Berlin, Math.-Nat. Reihe, 4, 541-580, 1954/55.
171. WERNER, K. : Der Einfluss der Höhenlage auf die Organisation der Bodenwirtschaft - Wiss.Z.Univ.Halle, Math.-Nat.Reihe, 3, 111-128, 1953/54.
172. WILAMOWITZ-MOELLENDORF und SCHUMANN, A. : Agrarmeteorologische Beratung im Bezirk Dresden - D.Dt.Landwirtsch., 8, 491-494, 1957.
173. ZENKER, H. : Singularitäten der Erdbodentemperatur in Halle (S) - Z.f. Met., 7, 237-245, 1953.
174. ZILLMANN, K.H. : Ueber den Wasserverbrauch einiger Kulturpflanzen am Standort in Abhängigkeit von den meteorologischen Faktoren und von der Bodenfeuchtigkeit - Z.f.Acker- u.Pflanzenbau, 104, 257-273, 1957.

VERZEICHNIS DER ABKÜRZUNGEN.

Abh.Met.u.Hydrolog.Dienst der DDR

Angew.Met.

Ann.d.Met.

Arch.f.Forstwesen

Arch.f.Gartenbau

Arch.f.Met., Geoph.u.Biokl.

Ber.d.Dt.Botan.Ges.

Der Züchter

Dt. Gartenbau

D.Dt.Landwirtsch.

Die Gartenbauwissenschaft

D.Naturwiss.

Forsch.u.Fortschr.

Nachr.-Bl.f.d.Dt.Pflanzenschutzdienst

Planta

Sitz.-Ber.Dt.Akad.Landwirtschaftswiss.Berlin

Umschau

Veröff.d.Inst.f.Agrarmet.d.Univ.Leipzig

Wasser u. Boden

Wasserwirtsch.-Wassertechn.

Wiss.Z.Humboldt-Univ.Berlin,Math.-Nat.Reihe

Wiss.Z.Univ.Greifswald,Math.-Nat. Reihe

Wiss.Z.Univ.Jena, Math.-Nat. Reihe

Wiss.Z.Univ.Leipzig, Math.-Nat. Reihe

Wiss.Z.Univ.Rostock

Z.f.Acker-u. Pflanzenbau

Z.angew.Zool.

Z.f.Met.

Z.f. Pflanzenbau u. Pflanzenschutz

Z.f. Pflanzenzüchtung

Abhandlungen des Meteorologischen und Hydrologischen Dienstes der DDR.

Angewandte Meteorologie

Annalen der Meteorologie

Archiv für Forstwesen

Archiv für Gartenbau

Archiv für Meteorologie, Geophysik und Bioklimatologie

Berichte der Deutschen Botanischen Gesellschaft

Der Züchter

Der Deutsche Gartenbau

Die Deutsche Landwirtschaft

Die Gartenbauwissenschaft

Die Naturwissenschaften

Forschungen und Fortschritte

Nachrichtenblatt für den Deutschen Pflanzenschutzdienst.

Planta - Archiv für wissenschaftliche Botanik

Sitzungsberichte der Deutschen Akademie der Landwirtschaftswissenschaften zu Berlin.

Umschau in Wissenschaft und Technik

Veröffentlichungen des Instituts für Agrarmeteorologie der Universität Leipzig

Wasser und Boden

Wasserwirtschaft - Wassertechnik

Wissenschaftliche Zeitschrift der Humboldt- Universität Berlin, Mathematisch-Naturwissenschaftliche Reihe.

Wissenschaftliche Zeitschrift der Universität Greifswald, Mathematisch - Naturwissenschaftliche Reihe.

Wissenschaftliche Zeitschrift der Universität Jena, Mathematisch-Naturwissenschaftliche Reihe.

Wissenschaftliche Zeitschrift der Universität Leipzig, Mathematisch-Naturwissenschaftliche Reihe.

Wissenschaftliche Zeitschrift der Universität Rostock.

Zeitschrift für Acker- und Pflanzenbau.

Zeitschrift für angewandte Zoologie

Zeitschrift für Meteorologie

Zeitschrift für Pflanzenbau und Pflanzenschutz.

Zeitschrift für Pflanzenzüchtung.

Section D : Requests from members and general information

Section E : Requests from non-members

Section F : Book reviews

MISCELLANEOUS BIOCLIMATOLOGICAL DATA

Section F: Book Reviews*

- L. AARIO AND H. JANUS : BIOLOGISCHE GEOGRAPHIE (DAS GEOGRAPHISCHE SEMINAR) - Georg Westermann Verlag, Braunschweig 1958, with 7 illustrations, 3 tables, 136 pages. Price 5.80 German marks.

This booklet is a recent issue of the series "Geographical Seminar" published by the Georg Westermann Verlag. It gives a summary of the geographical distribution of plants (by Prof. Aario, Helsinki, Finland) and of animals (by Dr. Janus, Stuttgart, Germany) and of the various factors affecting this distribution, one group being climatological factors. Although the booklet is primarily intended for students of Geography, Zoology and Botany, it provides interesting reading for every bioclimatologist, the more so as it discusses the possible effect of environment on permanent changes in the living organism.

- H. BERG : SOLAR-TERRESTRISCHE BEZIEHUNGEN IN METEOROLOGIE UND BIOLOGIE - Akademische Verlagsgesellschaft Geest & Portig K.-G., Leipzig 1957, With 77 diagrams, 200 references, 172 pages.

This book is published as volume 30 of the series "Probleme der Kosmische Physik" (Problems of Cosmic physics) by H. Berg, Professor of Meteorology and Geophysics at the University of Köln, Germany and Vice President of the Internat.Soc.Bioclim. Biomet.

Bioclimatologists should be grateful to Prof. Berg that he has been prepared to summarise the vast amount of publications on this subject of which a very large part lacks any scientific or statistical basis. As a result most scientists are inclined to reject automatically any new publication in this field. Prof. Berg's objective, but at the same time very critical study of the problem gives an excellent summary of all these publications, and clearly indicates which conclusions should be rejected or accepted and which problems require further study.

The book can be highly recommended to bioclimatologists interested in the problem of the possible relationship between cosmic events, weather, climate and biological phenomena.

- E. BUNNING : DIE PHYSIOLOGISCHE UHR - Springer-Verlag, Berlin, Göttingen, Heidelberg 1958, with 107 diagrams, 105 pages, price 24 German marks.

Many bioclimatological phenomena observed both in man and animal seem, at least partly, to be caused by the disturbing effect of weather and climate on the endogenous biological rhythm of living organisms. The problems of Biological Rhythms should therefore be included in many bioclimatological studies.

Many workers in the field of Biological Rhythms (united in the Int. Soc. of Biological Rhythm, Secretariat: Dr. A. Sollberger, Dept. of Anatomy, Karolinska Inst., Solnavägen 1, Stockholm 60, Sweden) are often only interested in rhythmic phenomena observed in their own specific field of research, either zoology, botany etc. It is often not sufficiently realized that many of these phenomena in various groups of organisms are closely related to each other.

Prof. Bünning, a well known botanist by profession, has tried to unite these various principles, in a booklet called "Physiological Time". The book is based on a series of lectures, held in the USA (1956) and London (1958). The many diagrams, extensive list of references and clear style make this booklet a valuable guide for bioclimatologists interested in this aspect of Bioclimatology.

- J.L. CLOUDSLEY-THOMPSON : SPIDERS, SCORPIONS, CENTIPEDES AND MITES (THE ECOLOGY AND NATURAL HISTORY OF WOODLICE, 'MYRIAPODS' AND ARACHNIDS) - Pergamon Press, I.R. Maxwell and Co. Ltd., London, 1959, price 50 shillings.

This book by Cloudsley-Thompson, member of the Executive Board of ISBB, discusses various aspects of Entomological Bioclimatology. Although there are a number of works on the morphology and systematics of many of these animals, there is no readily accessible account of their ecology and natural history. In these days of specialization the combination of physiology with ecology and a study of the influence of microclimates is an especially useful type of approach. This book consists of a number of chapters devoted to different groups of animals, the chapters themselves consisting of a brief outline of the classification and distribution of the group, followed by detailed accounts of their general behaviour, food and feeding habits, enemies (predators and

* Titles of reprints of articles received from members since 1956, will be given in the following issues of the Journal.

parasites), courtship and mating habits, reproduction and growth stages. It is documented with bibliographies that also contain references to publications that can be used for the identification of specimens and supplied with a glossary and indexes. Various tropical groups, such as scorpions and Solifugae not represented in temperate regions, are included not only for completeness, but because they are of interest for purposes of comparison and will make the book useful to university students, research workers and amateur naturalists in all parts of the world.

E. GAGEL : DIE SIEBER MEERE - Georg Westermann Verlag, Braunschweig 1958, with 41 photographs, 105 diagrams, 18 tables, 236 pages.

This well illustrated book on marine hydrology does not pretend to be a textbook. It gives in simple language a good summary of various aspects of modern marine hydrology, although in the section on submarine canyons we missed the modern theories and experiments on turbidity currents. It discusses the effect of weather and climate on sea currents and their biological effects. The book can be recommended to bioclimatologists, in particular those interested in general problems of marine bioclimatology.

F. LAUSCHER, M. ROLLER, G. WACHA, M. GRAMMER, E. WEISS AND J.W. FRENZEL : WITTERUNG UND KLIMA VON LINZ - Sonderheft VI, Zeitschrift "Wetter und Leben", vol. X, 1958, with 21 illustrations, 30 tables and 235 pages.

This publication was prepared at the request of the "Kulturamt" of the city of Linz and represents a complete analysis of the meteorological and climatological data of the city which are required for the study of various problems in relation to urban bioclimatology. For example, airpollution is a very important problem in this industrialized city with its days of heavy fog. Members working in the field of Urban Bioclimatology are recommended to study this interesting publication.

LOCUSTA NO. 6 : BULLETIN OF THE INTERNATIONAL AFRICAN MIGRATORY LOCUST ORGANIZATION 1959.

The latest issue received contains three interesting articles by:

G.B. Popov : Ecological studies on oviposition by *Locusta migratoria migratorioides* (R. & F.) in its outbreak area in the French Sudan.

V.M. Dirsh : The early stages of *Gastrimargus nigericus* Uvarov 1926 (Acridoidea, Orthoptera).

A. Shulov and M.P. Pener : A contribution to knowledge of the development of the egg of *Locusta migratoria migratorioides* (R. & F.).

Members interested in this aspect of Entomological Bioclimatology are also recommended to study previous issues:

vol. 1 (July 1954) : B. Zolotarevsky - Historique de l'Organisation internationale de lutte préventive contre le Criquet migrateur africain.

vol. 2 (October 1954) : G. Remaudière - Etude écologique de *Locusta migratoria migratorioides* Rch. & Frm. (Orth., Acrididae), dans la zone d'inondation du Niger en 1950.

vol. 3 (February 1955) : J.T. Davey - A preliminary note on seasonal movements of the African Migratory Locust in the Solitary Phase.

vol. 4 (November 1956) : F.O. Albrecht - Limitation des effectifs chez un Acridien : Influence de la sécheresse du sol sur les oeufs de *Nomadacris septemfasciata* (Serv.). J.S.B. Choudhuri - Experimental studies on the selection of oviposition sites of *Locusta migratoria migratorioides* (R. & F.).

vol. 5 (December 1957) : J.T. Davey - The African Migratory Locust (*Locusta migratoria migratorioides* Rch. & Frm., Orth.) in the Central Niger delta Part One, Climate and Vegetation.

S.W. TROMP : THE GEOGRAPHICAL DISTRIBUTION OF ARTERIOSCLEROTIC HEART DISEASES IN THE NETHERLANDS (PERIOD 1936-1945) - Bioclimatological Research Centre, Leiden, vol. III, 20 November 1958, with 7 maps, 3 graphs, 8 tables, 12 pages.

On the basis of mortality data in the different municipalities in the Netherlands (standardized for age and number of inhabitants) a number of geographical distribution maps were prepared which indicate high mortality in the 5 coastal provinces and relatively low values in the Eastern provinces. The inhabitants of these coastal provinces do not differ specifically in food habits from those of the Eastern provinces. These and various other considerations make it unlikely that differences in food could explain the observed geographical distribution. The possible influence of weather and climatic differences between the Eastern and coastal provinces are discussed and their possible effect on arteriosclerotic diseases.

S.W. TROMP : THE GEOGRAPHICAL DISTRIBUTION OF CANCER OF THE LUNG IN THE NETHERLANDS (PERIOD 1936-1952) - Bioclimatological Research Centre, Leiden, vol. IV, 6 January 1959, with 5 maps, 15 tables, 10 pages.

The geographical distribution of lung cancer mortality in a standardized population is represented

ted in a number of maps. A very peculiar geographical distribution is found which cannot be explained by differences in smoking habits or air pollution. The areas with high lung cancer mortality are roughly opposite to those with high Bronchitis-Pneumonia mortality. The boundary, a S.W.-N.E. line in the Netherlands, roughly coincides with a climatic boundary; i.e. west of this line (nearer to the coast) low Bronchitis-Pneumonia mortality occurs, east of this line high mortality values are observed. This boundary is neither a racial or food boundary, or related to different medical facilities east and west of it. The influence of climate seems therefore possibly the cause of the observed distribution.

S.W. TROMP : BIOMETEOROLOGICAL ANALYSIS OF DAILY AND MONTHLY VARIATIONS IN RESTLESSNESS AND ILL-TEMPEREDNESS OF MENTAL PATIENTS IN THE WESTERN PART OF THE NETHERLANDS (PERIOD 1 NOVEMBER 1956 - 1 DECEMBER 1958).
First report: Psychiatric Institute "Hulp en Heil" (at Leidschendam)- Bioclimatological Research Centre, Leiden, vol. V, 5 April 1959, with 25 charts, 28 pages.

This publication represents the first report on data collected in one of seven large Psychiatric Institutes in the Netherlands cooperating in this project. Apart from monthly and pseudoseasonal variations, it was found that restlessness of Schizophrenic patients increases with influx of warm air masses; in Epileptic patients a decrease was observed. In Epileptic patients both general motoric restlessness and number of epileptic attacks increases with influx of cold airmasses. A mathematically significant correlation was found with cold front passages. Electrophoretic serum protein studies of mental patients showed characteristic deviations in Schizophrenics which were less pronounced or absent in other groups of mental patients.

H. WILHELMY : KLIMAMORPHOLOGIE DER MASSENGESTEINE - Georg Westermann Verlag, Braunschweig 1958, with 137 illustrations, 385 references and 238 pages, price 35.80 German marks.

Prof. Wilhelmy of the Dept. of Geography of the Technical University of Stuttgart has prepared an interesting book with many original photographs on the influence of the microclimate on soil and topography. Whereas around 1925 geomorphologists were inclined to overestimate the effect of movements of the earthcrust (tectonics) on the topography of the earthcrust, in recent years the great importance of weather and climate on soil topography is becoming appreciated more and more.

R. Geiger in his wellknown book "Das Klima der bodennahen Luftschicht" discussed mainly the effect of soil topography on the microclimate of an area. The effect of this microclimate on soil is generally a less common field of study. Although strictly speaking this subject does not belong to bioclimatology sensu stricto, it is felt that it is essential for agro- and forest meteorologists to be acquainted with the effect of microclimate in soil. For this group of bioclimatologists particularly Wilhelmy's book is an easy guide to the various problems in this field. The printing of many less known pictures of rock morphology is refreshing as many textbooks continue to reproduce only the same internationally known geomorphological points of interest.

GUIDE BOOK TO RESEARCH DATA FOR ARID ZONE DEVELOPMENT (ARID ZONE RESEARCH VOL. IX) - Unesco Publication (Edited by B.T. Dickson), 1957, 191 pages.

This excellent summary of research data required for arid zone development projects consists of the following parts and chapters:

Part I: Physical and Biological factors.

- Chapter 1 : Introduction (by B.T. Dickson, Botanist, former Chief Division of Plant Industry, CSIRO, Australia).
- Chapter 2 : Mapping (by F. George, Agricultural Officer, FAO, Rome).
- Chapter 3 : Climatology (by E.G. Bilham, formerly Deputy Director of the Meteorological Office, London).
- Chapter 4 : Geology and geophysics (by E. Sherbon Hills, Professor of Geology and Mineralogy, University of Melbourne, Australia).
- Chapter 5 : Geomorphology (by C.S. Christian, Officer-in-charge, Land Research and Regional Survey Section, CSIRO, Australia; J.N. Jennings, Reader, Dept. of Geography, Australian National University, Australia; C.B. Twidale, Geomorphologist, Land Research and Regional Survey Section, CSIRO, Australia).
- Chapter 6 : Hydrology (including irrigation) (by J. Tison, Secretary General of the International Association of Hydrology).
- Chapter 7 : Soil Resources (by Herbert Greene, Adviser on Tropical Soils, Colonial Office, United Kingdom).
- Chapter 8 : Vegetation (by W.G. McGinnies, Director of the Central States Forest Experiment Station, Columbus, Ohio, USA).
- Chapter 9 : Forests (by M. Leloup, Head, Forestry Division, FAO, Rome).

Part II: Human factors.

- Chapter 1 : Population (by E. Sargent).
- Chapter 2 : Land tenure (by E.H. Jacoby, Land Tenure Specialist, FAO, Rome).
- Chapter 3 : Animal Husbandry (by R.O. Whyte, Agriculture Division, FAO, Rome).
- Chapter 4 : Nutrition, health and sanitation (by WHO).
- Chapter 5 : Sources of energy (by E.W. Golding).

Other publications in the same arid zone series are:

- vol. I : Reviews of Research on Arid Zone Hydrology.
- vol. II : Proceedings of the Ankara Symposium on Arid Zone Hydrology.
- vol. III : Directory of Institutions Engaged in Arid Zone Research.
- vol. IV : Utilization of Saline Water. Reviews of Research.
- vol. V : Plant Ecology. Proceedings of the Montpellier Symposium.
- vol. VI : Plant Ecology. Reviews of Research.
- vol. VII : Wind and Solar Energy. Proceedings of the New Delhi Symposium.
- vol. VIII : Human and Animal Ecology. Reviews of Research.
- vol. X : Climatology. Reviews of Research.

This volume contains the eight following reports, together with a large bibliography: "Evaporation and the water balance", by E.L. Deacon, C.H.B. Priestly and W.C. Swinbank; "Climatic factors in arid zone animal ecology", by F.S. Bodenheimer; "Radiation and the thermal balance", by A.J. Drummond; "Climates and vegetation", by A. Vernet; "Proprioclimates of man and domestic animals", by D.H.K. Lee; "The modification of microclimates", by E.M. Fournier d'Albe; "The chemical climate and saline soils in the arid zones", by E. Eriksson; "Climatological observational requirements in arid zones", by M. Gilead and N. Rosenan.

- vol. XI : Climatology and microclimatology: proceedings of the Canberra Symposium.
This volume contains the 50 papers presented to the symposium: each paper is given in its original language, followed by a summary in the other language.
The papers are grouped in the following eight sections, each of which concludes with an analytical summary of the discussions: "Evaporation and the water balance"; "Radiation and the thermal balance"; "Interrelationships of climatic factors and flora"; "Interrelationships of climatic factors and fauna"; "Microclimate of man and domestic animals"; "Modification of microclimates"; "Salting and chemistry of rainwater"; "Climatological observational requirements in arid zones".

**Section G : International Organizations (WMO, FAO,
WHO, etc.)**

MISCELLANEOUS BIOCLIMATOLOGICAL DATA

Section G: International Organizations (Unesco)

RECENT ARID ZONE PROJECTS

1. UNESCO-SPAIN SYMPOSIUM ON PLANT-WATER RELATIONSHIPS IN ARID AND SEMI-ARID CONDITIONS:

A Symposium on Plant-Water Relationships in Arid and Semi-Arid Conditions, organized jointly by the Spanish Government and Unesco, will be held in Madrid from 24 to 30 September 1959. This meeting should provide an opportunity for a number of scientists to present and discuss scientific papers on original research.

PROGRAMME:

Methodology of water relation studies on plants.
Water sources for plants.
Water balance of plants under arid and semi-arid conditions.
Drought-resistance of plants.
Practical applications to agronomy.

2. ECOLOGICAL MAP OF THE MEDITERRANEAN AREA

At the proposal of the Food and Agriculture Organization's Forestry Division, the Unesco Advisory Committee on Arid Zone Research recommended, at its thirteenth session, the organization of a meeting of experts on plant ecology for the purpose of studying the preparation of an ecological map for the Mediterranean area. Following this recommendation, an FAO/Unesco Study Group was set up and met in Paris from 5 to 9 January 1959.

While maintaining "Ecological Map of the Mediterranean Area" as the title of the project, the Study Group decided to consider an area extending beyond the Mediterranean zone proper, and to map regions located approximately between the twentieth and forty-eighth parallels, excluding however the mountains of the northern portion, and stretching from the Atlantic to the Indus river. This delimitation would have the advantage of including almost the entire region in which activities under Unesco's major project are concentrated.

The Study Group, taking into account the international aim pursued, the number of maps available and financial considerations, decided to select a scale of 1/5,000,000. On this small scale it is possible to obtain very accurate overall perspectives for the study of the development of the countries concerned. Furthermore, this scale has the advantage of expediting map compilation. Because of the scale selected, the group proposed to draw up two maps - a bioclimatic map and a vegetation map.

BIOCLIMATIC MAP. The bioclimatic map will be drawn up by the ombrothermic curve method described by Mr. Gaussen, taking into account, if possible, the number of rainless days. The sites covered by this map will be selected on the basis of data furnished by Mr. Emberger's method, and the combination of these two methods will make a more accurate subdivision of the bioclimatic regions possible. This general map will be supplemented by bioclimatic maps on a smaller scale for the regions of Australia, South Africa and America with analogous climatic conditions. This work, undertaken by Messrs. Gaussen and Emberger, will be completed by a documentation reference notice giving the rainfall and temperature curves, the absolute values used and an explanation of the methods employed.

VEGETATION MAP. This map will represent the climatic vegetation, indicating by different colours the fundamental forest stages and the various types of steppe formations. Against this coloured background will be shown, in so far as possible, the condition of present-day vegetation, considering particularly the most valuable vegetation species or groups of species. As in the case of the bioclimatic map, this map will be supplemented by small reduced-scale maps showing the climatic vegetation of the homologous regions of the other parts of the world.

Preliminary work. While awaiting the drawing up of the complete bioclimatic map, the Study Group requested Messrs. Gaussen and Emberger to prepare immediately a preliminary general map on a scale of 1/5,000,000 accompanied, as example, by the final maps (bioclimatic and vegeta-

tion) of Morocco. This work will serve as a basis for discussion of the final complete map.

The Study Group was perfectly aware of the obviously limited possibilities of a map on a scale of 1/5,000,000 and recommended to FAO and Unesco that they frame a programme for the preparation of maps on a scale 1/1,000,000 against the international background of the region considered, in conjunction with projects for compiling other maps, particularly soil maps, on a scale of 1/1,000,000. These 1/1,000,000 scale maps are absolutely essential for adequate knowledge of actual and potential land use conditions.

A convincing illustration of the value of such 1/1,000,000 maps is given by the plant cover map of Tunisia, which was presented to the Study Group with the accompanying report drawn up by Messrs. Gausson and Vernet.

This map, printed over the international 1/1,000,000 scale map, shows the following:

1. Present permanent or temporary crops, with statistical indications enabling basic information to be shown on the map (for instance, one symbol representing 25,000 olive trees, etc.), are shown by means of contour lines established with the aid of aerial photographs.
2. Natural vegetation of three types is also shown by contour lines drawn from aerial photographs: forests shown in flat tint, scrub vegetation in lines, and herbaceous vegetation by dotted lines.
3. The basic idea being that of the "vegetation successional stage" (beech stage, holm - oak stage, etc., named according to the climax formation), and the successional stage being dependent on ecology, each stage is designated by a colour which thus represents an ecological value.
4. Alongside the principal map are the following small maps, on a scale of 1/5,000,000, giving numerous supplementary details: a hypsometric map; a geological or soil map; a botanical map indicating the manner in which the vegetation would evolve if men were eliminated; an administrative map; a climatic map; a land use map.

Many details are given in an appendix, as is also an analysis of the requisites for the development of the region involved.

It must be noted that, in order to cover on a scale of 1/1,000,000 the regions of the Mediterranean area defined above, some 45 different maps would have to be published for the vegetation alone. Such a programme could be carried out only as the result of considerable efforts by the countries concerned, in conjunction with wide-scale international action such as that envisaged by the new United Nations Special Fund for Economic Development.

3. FAO-UNESCO DESERT LOCUST ECOLOGICAL SURVEY

The first meeting of the Inter-Agency Working Group established to advise on the conduct of the FAO/Unesco Desert Locust Ecological Survey was held at the Anti-Locust Research Centre, London, from 11 to 13 February 1958. That meeting outlined the general problem, the objectives of the survey and planned the first survey in the summer breeding areas of the desert locust in the Sudan and Chad Territory.

The second meeting was convened at Unesco Headquarters in Paris on 29 and 30 December 1958 to review the results of the first survey and to plan further operations.

The meeting received an outline report of the first survey which was conducted in Sudan, Chad Territory and Northern Ethiopia from 18 June to 21 November 1958. This report will be circulated to all interested governments and organizations as soon as it is completed.

After examining possible areas for spring/summer surveys, the meeting recommended that the next survey should be undertaken in the Republics of Mauritania, Sudan and Niger.

This selection was influenced by the offer of Messrs. Frézal and Mallamaire to submit to their respective administrations proposals with a view to securing their full cooperation and assistance, particularly in loaning vehicles, equipment and technical staff to accompany the survey. The Survey Team assembled at Dakar at the end of February to prepare for proceeding to the field at the beginning of March. It is envisaged that the survey, commencing in Mauritania and later working eastwards, will continue its work in West Africa until about September. Owing to the importance of the Arabian peninsula in the desert locust problem, the meeting considered that this area should be the next objective for a survey during the winter and spring of 1960 if it were found practicable. It was also felt that it might be desirable at that time for the plant ecologist to return to an area previously visited in order to study the state of the vegetation at different times of the year.

4. UNESCO-WMO EXPERT MEETING ON SOLAR RADIATION PROBLEMS

Following on the recommendations of the Advisory Committee on Arid Zone Research at its fourteenth session, Unesco and WMO organized a meeting of experts on solar radiation in order to set up detailed plans for the following two projects:

1. A climatological survey of solar radiation in arid zones.
2. The development of a simple, robust and cheap solar radiation recorder for the utilization of solar energy.

The meeting was held at the WMO Secretariat in Geneva from 25 to 27 February 1959, under the chairmanship of Dr. W. Mörikofer, Director of the Physical Meteorological Observatory, Davos.

The meeting first studied in general terms the role of radiation measurement in the context of research activities on arid zone problems. Although the immediate interests of such measurements in arid zones would be in applied research, the group agreed that basic research on climatological problems of radiation should have priority within the Unesco programme, since a thorough knowledge of radiation and its related problems was essential to any balanced programme of research on arid zones.

These general considerations were followed by a more detailed discussion of the following radiation problems which are of evident importance for applied research: (a) utilisation as a source of energy; (b) effects on soil conditions; (c) effects on plant cover; (d) effects on humans and animals; (e) heat balance (short and long-wave radiation).

The group went on to discuss details of the proposed survey of existing radiation data. It agreed that the survey should cover the whole region stretching from North Africa and the Middle East to South Asia where the major project activities are concentrated. Where data are scanty, the materials from stations in adjacent and climatologically similar areas will also be used.

Recommendations were made as to the kind of data necessary for the survey and the possible sources. It was agreed that the work will be done mainly at the Davos Observatory by a young research worker with experience in radiation measurements and an assistant, the whole project being under the supervision and guidance of Dr. Mörikofer. Close contact with the Secretariat of WMO will be maintained, especially since it is through WMO that most of the material from the files of meteorological institutes will be collected, including those resulting from observations during the International Geophysical Year.

5. A REGIONAL COURSE ON PLANT ECOLOGY IN SOUTH ASIA

In the framework of the major project, a regional training course on plant ecology, organised by the Unesco Science Co-operation Office for South Asia with the assistance of the Indian authorities, took place at Jaipur (India), from 3 to 21 December 1958. It was similar to the course organized earlier in 1958 in the U.A.R. (Egypt). This course was intended to inform research workers and specialists in the governmental services concerned about the latest developments of ideas and practical methods of plant ecology. The course was placed under the direction of Professor L. Emberger, Director of the Botanical Institute of the University of Montpellier, France.

6. TRAINING ARID ZONE SPECIALISTS

A regional course on microclimatology, jointly organized by the Pakistan Meteorological Service and the Unesco Science Co-operation Office for South Asia, will be given this summer at Quetta, at the Geophysical Research Institute. This course, which will last one month, will comprise a series of lectures on the various aspects of microclimatology in arid zones, and practical work in measurement and interpretation of microclimatological data. It will be directed by Mr. R.O. Slatyer, Land Research and Regional Survey Division, CSIRO, Canberra (Australia), with the assistance of Mr. Hoyle, expert of the World Meteorological Organization, who is at present in Iran, and of specialists of the Pakistan Meteorological Service and the other appropriate services. It is expected that approximately fifteen participants from Pakistan and fifteen from the neighbouring countries will be invited to attend this course, which will be a follow-up, on the regional level, of the symposium organized in Canberra in 1956.

7. GENERAL INFORMATION

Prohuza (French Study and Information Centre for Human Problems in the Arid Zones) organized Saharan Medico-Social Information Days in Paris from 9 to 11 April 1959: the proceedings took place in the Unesco Conference Hall. The subjects discussed were biological and psycho-social adaptation of man to the desert; hygiene and disease prevention; housing and acclimatization; labour and economy. Numerous French scientists, administrators, doctors and engineers, as well as a number of foreign experts took part in these discussions.

An interesting experiment in irrigation is now being carried out by Messrs. H. and E. Boyko, of the Agricultural Research Station of the Israeli Ministry of Agriculture. Based on observations taken on coastal dunes, this experiment shows that soils of similar high permeability make it possible to cultivate certain salt-resistant species of economic value even when they are irrigated with water which has a high concentration of sodium chloride. The species used in this first experiment, begun in July 1957, are *Juncus maritimus* Lam. (a valuable cellulose plant) and *Agropyrum junceum* (fodder plant with a high nutritive value). The success of this method might render it possible to make barren sand dunes and similar soils productive by irrigating them with water having a high salt content, or even with sea water.

8. PUBLICATIONS RECENTLY RECEIVED BY UNESCO

CLIMATE AND MAN IN THE SOUTH-WEST. Proceedings of a symposium, edited by Terah L. Smiley. Tucson, Arizona, The University Press, 1958, 84 pp.

COMMENT CONSTRUIRE AU SAHARA. Cahier du Centre Scientifique et Technique du Bâtiment, 4, avenue du Recteur-Poincaré, Paris-16^e, July 1958, 83 pp.

A climatic, architectural and economic study of the problems of building living and office accommodation in a hot dry climate; technical annexes and bibliography.

ECOLOGIE ET PEUPLEMENT ENTOMOLOGIQUE DES SABLES VIFS DU SAHARA NORD-OCCIDENTAL, by F. Pierre. Centre National de la Recherche Scientifique, 13, quai Anatole-France, Paris-7^e, 1958, 332 pp. This publication describes research carried out between 1947 and 1953, mainly in the Beni-Abbes region. The first part deals with the origin and distribution of the sands, ecological study, microclimates and biotic factors. The second discusses the entomological population, its varying composition according to the types of sand, its special features and groupings.

FLORE DU SAHARA SEPTENTRIONAL ET CENTRAL, by Paul Ozenda. Centre National de la Recherche Scientifique, 13, quai Anatole-France, Paris-7^e, 1958, 488 pp., 16 plates, octavo.

The first part of this book, "Introduction to the botany of the Sahara", begins with a general description of desert conditions, followed by studies on conditions in the Sahara, the composition and origin of the flora of the Sahara, the biology of desert plants, plant communities in the Sahara, and the relation between vegetation and human occupation. The second part is an analytical flora for identifying vascular plants in the geographical area covered. It can also be used for most of the species found in the Southern Sahara. A number of drawings and about 15 photographic plates help readers in consulting the book.

HOMMES ET DESERTS: EVALUATION BIOCLIMATOLOGIQUE DES SITES D'INDUSTRIALISATION AU SAHARA, by Dr. G. Lambert. Centre d'Etudes et d'Information des Problèmes Humains dans les Zones Arides, 31, rue de Bassano, Paris-8^e, 1958.

HOMMES ET DESERTS: INFLUENCE DU CLIMAT TROPICAL SUR LE COMPORTEMENT PHYSIOLOGIQUE DE L'HOMME, by R. Lemaire. Centre d'Etudes et d'Information des Problèmes Humains dans les Zones Arides, 31, rue de Bassano, Paris-8^e, 1958.

M I S C E L L A N E O U S B I O C L I M A T O L O G I C A L D A T A

Section G: World meteorological organization

WORLD METEOROLOGICAL ORGANIZATION AND INTERNATIONAL
SOCIETY OF BIOCLIMATOLOGY AND BIOMETEOROLOGY TO COLLABORATE

by

Frederick Sargent, II, M.D. (U.S.A.)

President I.S.B.B.

In June of this year the president of this Society received a letter from the Secretary-General of the World Meteorological Organization informing him that the Executive Committee of WMO had, in Resolution 15 (EC-XI), directed the Secretary-General to (1) collaborate with this Society and (2) investigate the most proper and efficient manner in which bioclimatology and biometeorology of man may be integrated into the activities of WMO. The resolution, the letter from the Secretary-General to the President, and the President's reply are reproduced below.

Some comments on the background of this resolution may be of interest. Since the formal founding of this Society in August, 1956, at UNESCO H.Q. in Paris excellent relations have been maintained with WMO. Observers from WMO have attended the congresses of I.S.B.B. and likewise observers from I.S.B.B. have attended the meetings of the Commissions on Climatology and Agricultural Meteorology. At the 2nd Congress of the Commission on Climatology, Washington, D.C., January, 1957, the president of I.S.B.B. urged WMO to collaborate with I.S.B.B. This Society could act in an advisory capacity and keep WMO informed on current developments in bioclimatology and biometeorology. Similar suggestions have been made by Mr. P.M.A. Bourke at more recent meetings of the Commission on Agricultural Meteorology at Warsaw and Geneva and by the Secretary-General of I.S.B.B. at the recent world conference of W.M.O. itself at Geneva. Now these suggestions have born fruit. Just what form the collaboration will finally take cannot be foretold.

The aims of the Executive Board are to integrate the efforts of the many UN agencies engaged or interested in bioclimatology and biometeorology, to avoid unnecessary duplication of work, and encourage closer collaboration among these agencies. It may be noted in this regard that progress is also being made toward active collaboration with the World Health Organization. The Board hopes to be able to report further developments along these lines by September, 1960.

WORLD METEOROLOGICAL ORGANIZATION

Resolution 15 (EC-XI)

BIOCLIMATOLOGY AND BIOMETEOROLOGY OF MAN

THE EXECUTIVE COMMITTEE,

NOTING,

1) Recommendation 6 (CAGM-II);

2) The wish expressed by Third Congress that more attention should be given to bioclimatology and biometeorology;

DIRECTS the Secretary-General,

1) To collaborate with and to support the International Society of Bioclimatology and Biometeorology;

2) To investigate, in consultation with the Presidents of the Technical Commissions concerned, the most proper and efficient manner in which bioclimatology and biometeorology of man may be integrated into the activities of WMO.

ORGANISATION METEOROLOGIQUE MONDIALE

WORLD METEOROLOGICAL ORGANIZATION

SECRETARIAT: AVENUE DE LA PAIX, CAMPAGNE RIGOT-GENEVE

no. 6.683/59/DT/CL/BI
Annexes: 1

GENEVE 11 June 1959

Dear Sir,

I take pleasure in sending you herewith a copy of Resolution 15 (EC-XI), passed by the recent eleventh session of the Executive Committee of the World Meteorological Organization and entitled "Bioclimatology and Biometeorology of Man".

I have approached the Presidents of two of the WMO Commissions concerned namely the Commission for Climatology (CCl) and the Commission for Agricultural Meteorology (CAgM), suggesting that the former might wish to establish a working group to deal with this important and many-sided problem.

I shall not fail to keep you informed on further developments in this matter. In the meantime I should welcome any comment or suggestion you might wish to make in this connexion.

Yours sincerely,

(D.A. Davies)
Secretary-General

The President,
International Society of
Bioclimatology and Biometeorology,
Dept. of Physiology,
524 Biology Building,
University of Illinois,
URBANA, Ill.,
U.S.A.

cc. Mr. R.G. Veryard, President of CCl, London, England.
Mr. P.M.A. Bourke, President of CAgM, Dublin, Ireland.
Mr. M.S. Kulik, Vice-President of CAgM, Moskva, U.S.S.R.

INTERNATIONAL SOCIETY OF BIOCLIMATOLOGY AND BIOMETEOROLOGY

June 23, 1959

Dr. D.A. Davies
Secretary-General
World Meteorological Organization
Avenue De La Paix
Campagne Rigot - Geneva.

Dear Doctor Davies:

Re: 6.683/59/DT/CL/BI

I was truly gratified to receive your letter of 11 June in which you informed me that the Executive Committee of the World Meteorological Organization had resolved to develop a program related to human bioclimatology and collaborate with this Society. I am confident that much useful work can be accomplished by this collaboration, and I look forward to hearing from Mr. Veryard and Mr. Bourke on this matter.

I should like to pass on a few thoughts regarding the program which might be adopted. Insofar as a formal working group is concerned, I think that that would represent unnecessary duplication of the activities of this society. I would propose rather that there be a liaison committee comprised of representatives of the society who might formally report to WMO regarding the work of this society and who would assist you in answering queries addressed to WMO on bioclimatological matters. The liaison committee might have four members, one from human bioclimatology, one from agricultural bioclimatology, one from animal bioclimatology, and one from entomological bioclimatology.

The long-range goals of the society should perhaps become the focus of the program of collaboration. These goals, as I see them, are (1) to achieve a critical synthesis of what is actually solid fact and what is fancy speculation and assumption in the area of bioclimatology; (2) to document the past and current literature of bioclimatology; (3) to work with existing abstract agencies with a view to enhancing their coverage of the literature; (4) to encourage investigation of basic and applied problems urgently in need of solution; and (5) to stimulate graduate training in this complex interdisciplinary subject. We hope to achieve these ends through our journal, our congresses, and our technical committees. If active collaboration with WMO can focus on these broad goals, their more rapid attainment should be assured.

This society will hold its next congress in September, 1960. It is my sincere hope that by that time, I can report to the society that this WMO resolution has borne fruit. For the moment, please be assured of my most enthusiastic support for the ideas represented in your letter.

cc. Dr. S.W. Tromp
Mr. R.G. Veryard
Mr. P.M.A. Bourke

Yours sincerely,

Frederick Sargent, II, M.D.
President.

M I S C E L L A N E O U S B I O C L I M A T O L O G I C A L D A T A

Section B: Reports of Congresses of National Bioclimatological Societies

P R O G R A M O F S Y M P O S I U M O N B I O C L I M A T O L O G Y *

during the 176th National Meeting of the American Meteorological Society held jointly with the section of Meteorology, American Geophysical Union at WASHINGTON, D.C., U.S.A. ON 5 MAY 1959 AT THE NATIONAL ACADEMY OF SCIENCES

Chairman: Prof. Konrad J.K. Buettner

BIOCLIMATIC WORK IN THE U.S. WEATHER BUREAU: by H.E. LANDSBERG, Office of Climatology, U.S. Weather Bureau, Washington, D.C.

ROLE OF BIOCLIMATOLOGY IN THE ARMED FORCES: by DOUGLAS H.K. LEE, M.D., U.S. Quartermaster Research & Engineering Centre, Natick, Mass.

CHANGES IN IDEAS ON CLIMATIC ORIGIN OF DISEASE: by FREDERICK SARGENT, II, M.D., Dept. of Physiology, University of Illinois, Urbana, Ill.

THE EMISSION AND DISPERSION OF RAGWEED POLLEN: by JAMES B. HARRINGTON, Jr. and A. NELSON DINGLE, Meteorological Laboratory, University of Michigan, Ann Arbor, Mich.

ELECTRIC SPACE CHARGES AND HUMAN HEALTH: by IGHO HART KORNBLUEH, M.D., The Graduate Hospital, University of Pennsylvania, Philadelphia, Pa.

WEATHER IN PUBLIC HEALTH RESEARCH: by NICHOLAS E. MANOS, Air Pollution Program, Dept. of Health, Education and Welfare, Washington, D.C.

CONDENSATION DROPLETS AND IONIZATION PATTERN IN SUBMARINE ATMOSPHERE: by K.E. SCHAEFFER, M.D., U.S. Naval Medical Research Laboratory, New London, Conn.

HOW GOOD OR BAD IS WEATHER FOR NORMAL AND SICK PEOPLE?: by KONRAD J.K. BUETTNER, Dept. of Meteorology and Climatology, University of Washington, Seattle, Wash.

* Copied from "Bulletin of the American Meteorological Society", vol.40, no.4, pp. 193 - 208, April 1959.

Information concerning reports etc. can be obtained from Prof. Konrad J.K. Buettner, Professor of Meteorology and Climatology, University of Washington, Seattle, Washington, U.S.A.

MISCELLANEOUS BIOCLIMATOLOGICAL DATA

Section G: International Organizations

THIRD SESSION OF THE WORLD METEOROLOGICAL CONGRESS

at

the Palais des Nations, Geneva, Switzerland

from 1 - 28 April 1959

On 22 August 1958 the I.S.B.B. received an invitation from the Secretary-General of the World Meteorological Organization, Mr. D.A. Davies, to designate an observer who would represent the I.S.B.B. at the third session of the W.M.O. at Geneva.

During the first part of the Congress Prof. Dr. H. Berg (Univ. of Köln) attended the Congress during 2 days as an observer of I.S.B.B. The Secretary of I.S.B.B., Dr. S.W. Tromp, attended the Congress from 15 - 18 April 1959.

At the second session of the Commission for Agricultural Meteorology of the World Meteorological Organization at Warsaw (Poland) (29 Sept.-17 Oct. 1958) (see report of Dr. Zofia Pieslak, I.J.B.B. vol. III, 1959, part VII, sect. G), a resolution was accepted (Recommendation 6 CAgM, Cg-III/Doc.27, Appendix E) in which it was suggested that "the W.M.O. Congress in Geneva should consider the most appropriate manner of incorporating Bioclimatology and Biometeorology of man into their activities". In adopting this recommendation CAgM expressed the view that the terms of reference of CAgM itself should not be extended to cover this new field, as the Commission already has a very heavy programme.

In the Report to the third World Meteorological Congress in Geneva by Mr. P.M.A. Bourke, President of the Commission for Agricultural Meteorology (Cg-III/Doc.51, 16-1-1959), on page 6, point 8.2, the following statement was made:

"The Commission considered that the subject of meteorology as applied to man, in health and disease, should be given a place amongst the activities of W.M.O., and recommended that Congress should consider the most appropriate way in which this might be done. It was considered, in view of the heavy programme of CAgM, that the terms of reference of this Commission could not be extended to include responsibility for meteorological aspects of mankind, in addition to plants and animals, without serious repercussions on its present field of activity".

In document Cg-III/Doc.112 (10-4-1959), submitted to the third W.M.O. Congress by the Secretary-General, letters were published from F.A.O. and Unesco concerning proposals for joint action with W.M.O. in the field of Agroclimatology and Bioclimatology.

These various documents were discussed on 18 April 1959 after the observer of I.S.B.B. had been welcomed by the Chairman of the Commission for Agricultural Meteorology and an opportunity was given to the Secretary-General of I.S.B.B. to explain to the Delegates of the Third Congress of W.M.O. the purpose, the organization and the various activities of the I.S.B.B.

On behalf of I.S.B.B. Dr. Tromp concluded his statement by expressing the desire of the members of I.S.B.B., that one of the means of having a more permanent close contact with W.M.O. would be the creation of a permanent working group on Bioclimatology and Biometeorology in W.M.O. As a result of these discussions the Chairman of the Committee on Technical Questions prepared document Cg-III/Doc.160 (23-4-1959) to the Plenary.

In paragraph 2 of this document the following statement was made:

"With regard to the question of human biometeorology dealt with in paragraph 8.2 of Cg-III/Doc. 51, the President of CAgM stressed the need for close liaison with the International Society for Bioclimatology and Biometeorology (I.S.B.B.) and suggested that it might be useful to organize a joint symposium with W.H.O. in conjunction with the I.S.B.B. on relations between meteorology and health. The representative of I.S.B.B., Dr. Tromp, informed the Committee that the Society had been founded in 1956 and it already had a membership of about 500 scientists, of which approximately one-third were meteorologists from 49 different countries. The members served in an individual capacity and all the work of the Society was done voluntarily. I.S.B.B. had set up a number of committees whose work was closely related to that of CCl and CAgM. He felt that it would be useful to have a Working Group on Bioclimatology and Biometeorology within W.M.O."

In paragraph 3 of this document "the President of CCl (Commission of Climatology) pointed out that the terms of reference of his Committee included the application of climatological data to man, his comfort and his activities. CCl was, therefore, very interested in the work of I.S.B.B.

A great deal of knowledge had been acquired in several countries on bioclimatology and biometeorology and the important thing was that in these fields meteorologists must work with scientists in other disciplines, such as physiologists, agriculturalists and physicians".

The Committee on Technical Questions recommended to Plenary that the following text be included in the General Summary of the work of the Third Session of the Congress of W.M.O.:

"Congress reviewed the activities of the Commission for Agricultural Meteorology outlined in the report of the President of the Commission. It was noted with satisfaction that all the working groups established at the second session of the Commission had been given specific terms of reference with well-defined tasks. Congress agreed that it was very desirable that a session of the Working Group on the Guide to Agricultural Meteorological Practices should be arranged, at which the Chairmen of the other working groups concerned with the preparation of the Guide should be present.

Congress learned with interest of the activities of the International Society for Bioclimatology and Biometeorology and considered that this Society had done some very useful work in the short time which had elapsed since its creation. W.M.O. should continue to collaborate with and to support the International Society of Bioclimatology and Biometeorology in every possible way".

M I S C E L L A N E O U S B I O C L I M A T O L O G I C A L D A T A
Section G: International Organizations

REPORT OF THE SECOND SESSION OF THE
COMMISSION FOR AGRICULTURAL METEOROLOGY OF THE
WORLD METEOROLOGICAL ORGANIZATION
AT WARSAW (Poland)*
(29 Sept. - 17 Oct. 1958)

The second Commission for agricultural meteorology of the World Meteorological Organization was held in Warsaw from 29. IX to 17. X 1958. Thirty-two countries from all parts of the world sent their delegates and experts. The Session was opened by the Minister of Navigation and Water Administration who directs the Polish Hydro-meteorological Service. At the inauguration were also present the Vice-Minister, members of diplomatic corps, representatives of particular health-resorts and guests. Dr. J.J. Burgos, President of the Commission for Agricultural Meteorology, was Chairman of the Commission. The General Secretariat of WMO was represented by Dr. O. Ashford. At the Session were also present representatives of international scientific associations, the International Geographical Union, International Society of Soil Science and International Society of Bioclimatology and Biometeorology.

When the official part ended, national representatives grouped into two committees, which worked simultaneously during the subsequent days and discussed a great many essential problems of agricultural meteorology. These included the necessity for meteorological observations in agriculture and the instruments and methods of measurement. It is worth while stressing the fact that the soil climate and especially the water conditions of agricultural soils awoke a particular interest. A topic of animated discussion was the method for measuring the water balance of soil and plant, the ways for ensuring that plants obtain the necessary quantity of soil moisture investigation of the influence of soil moisture on the wear of agricultural machines, on fuel consumption, and so on.

That long hours were sacrificed to the problems of agro-hydrology - embracing extremely important problems of agriculture and practical forestry - can be explained by the fact that this branch of agro-meteorology is not yet sufficiently developed in many countries.

Matters of information and forecasting, with proper classification of meteorological forecasts, were broadly discussed; the first was concerned with a synoptic approach, the second was of interest to agro-meteorologists. A great deal of time was also devoted to agro-climatic classifications and to world agro-climatic atlas.

The problems of teaching agricultural meteorology aroused an animated and lengthy discussion. The urgent necessity for the training of cadres of young agro-meteorologists as well as the matter of textbooks was stressed. The lack of textbooks can be felt, and it is well known that most of the textbooks of meteorology appearing on the market are adjusted to the needs of the farmer and only seldom to those of agricultural meteorology. Although a model textbook, produced under the auspices of the WMO would be welcome, it has been decided to give priority to the preparation of an agro-meteorological guide. Published by WMO, this will contain technical details of instruments and methods of agro-meteorological measurement. The contents of the guide will most probably in future form a part of the above-mentioned textbook on agricultural meteorology. The countries concerned have themselves to start work on this.

Among other points the Session considered the significance of biometeorology and bioclimatology as parts of established meteorology and hence the necessity for closer collaboration between WMO and ISBB. As a result of the discussion, it was decided to ask the next Congress of WMO to examine the possibility of spreading the scope of activity of WMO by extending its responsibility to studies in the domain of the bioclimatology and biometeorology of man. The best course seems to be the establishment of a special technical commission to deal with these problems.

Some of the subjects discussed at the Session, on account of their importance and difficulty, were transferred to the attention of specially created study-groups. These working groups will deal with the problems entrusted to them during the intersessional period.

* Report prepared by Dr. Zofia Piślak, agro-meteorological section of the Instytut Hydrologiczno-klimatologiczny, Ul. Podlśna 61, Warszawa 32, Poland.

At the last plenary meeting of the Session a long speech was made by Dr. J.J.Burgos, the retiring President of the Commission of Agricultural Meteorology, who had held office for two terms over a period of seven years. At the close of this speech, Dr. Burgos expressed his thanks to all who had helped him up to the very last day, in the performance of his difficult and responsible duties. Dr. M.A. Bourke (Meteorological Service of Ireland) was elected President of the Commission for the period of the new term, Dr. H. Geslin, the retiring Assistant Vice-President was replaced by Mr. M.S. Kulik (Central Institute of Forecasts - U.R.S.).

The delegates present at the Second Session of the Commission of agricultural meteorology made an excursion to Białowieża and Mazury, which gave them an opportunity to become acquainted with some of the stations of the Polish Hydro-meteorological Service. The delegates also visited the agro-meteorological research station at Brwinów near Warsaw.

Section H : Advertisements

DYNA-BILT
A PRODUCT OF
BURTMAN IRON WORKS, Inc
BOSTON, MASS. 02137